Martin Krã, yer Rasmussen

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/3300224/publications.pdf

Version: 2024-02-01

60 papers

1,424 citations

331670 21 h-index 361022 35 g-index

61 all docs

61 docs citations

61 times ranked

1704 citing authors

| # | Article | IF | CITATIONS |
|----|--|--------------|--------------|
| 1 | Effects of Calcium Source, Inulin, and Lactose on Gutâ€Bone Associations in an Ovarierectomized Rat Model. Molecular Nutrition and Food Research, 2022, 66, e2100883. | 3.3 | 19 |
| 2 | Time-dependent regulation of hepatic cytochrome P450 mRNA in male liver-specific PGC-1 \hat{l}_{\pm} knockout mice. Toxicology, 2022, 469, 153121. | 4.2 | 1 |
| 3 | Curcumin and quercetin modify warfarin-induced regulation of porcine CYP1A2 and CYP3A expression and activity <i>inÂvitro</i> . Xenobiotica, 2022, 52, 435-441. | 1.1 | 1 |
| 4 | Brain foods - the role of diet in brain performance and health. Nutrition Reviews, 2021, 79, 693-708. | 5.8 | 21 |
| 5 | Primary hepatocytes isolated from human and porcine donors display similar patterns of cytochrome p450 expression following exposure to prototypical activators of AhR, CAR and PXR. Current Research in Toxicology, 2021, 2, 149-158. | 2.7 | 9 |
| 6 | Tissue-specific expression and activity of cytochrome P450 1A and 3A in rainbow trout (Oncorhynchus) Tj ETQqC | 0.8gBT | /Ogerlock 10 |
| 7 | Bovine Satellite Cells Isolated after 2 and 5 Days of Tissue Storage Maintain the Proliferative and Myogenic Capacity Needed for Cultured Meat Production. International Journal of Molecular Sciences, 2021, 22, 8376. | 4.1 | 14 |
| 8 | Hepatic PGC- $1\hat{1}\pm$ is not essential for fasting-induced cytochrome p450 regulation in mouse liver. Biochemical Pharmacology, 2020, 172, 113736. | 4.4 | 5 |
| 9 | Differentially expressed marker genes and glycogen levels in pectoralis major of Ross308 broilers with wooden breast syndrome indicates stress, inflammation and hypoxic conditions. Food Chemistry Molecular Sciences, 2020, 1, 100001. | 2.1 | 4 |
| 10 | Porcine cytochrome P450 3A: current status on expression and regulation. Archives of Toxicology, 2020, 94, 1899-1914. | 4.2 | 9 |
| 11 | Background Diet Influences TMAO Concentrations Associated with Red Meat Intake without Influencing Apparent Hepatic TMAO-Related Activity in a Porcine Model. Metabolites, 2020, 10, 57. | 2.9 | 21 |
| 12 | Sex dictates the constitutive expression of hepatic cytochrome P450 isoforms in Göttingen minipigs. Toxicology Letters, 2019, 314, 181-186. | 0.8 | 11 |
| 13 | Supplementation of sows with L-Arginine during gestating and lactation affects muscle traits of offspring related with postnatal growth and meat quality: From conception to consumption. Meat Science, 2019, 152, 58-64. | 5 . 5 | 15 |
| 14 | Porcine cytochrome 2A19 and 2E1. Basic and Clinical Pharmacology and Toxicology, 2019, 124, 32-39. | 2.5 | 13 |
| 15 | End-product inhibition of skatole-metabolising enzymes CYP1A, CYP2A19 and CYP2E1 in porcine and piscine hepatic microsomes. Toxicology Letters, 2019, 303, 67-71. | 0.8 | 5 |
| 16 | Effects of Multi-Component Mixtures from Sewage Treatment Plant Effluent on Common Carp (Cyprinus carpio) under Fully Realistic Condition. Environmental Management, 2019, 63, 466-484. | 2.7 | 18 |
| 17 | 7-Hydroxylation of warfarin is strongly inhibited by sesamin, but not by episesamin, caffeic and ferulic acids in human hepatic microsomes. Food and Chemical Toxicology, 2018, 113, 14-18. | 3.6 | 4 |
| 18 | Skeletal Muscle Interleukin-6 Regulates Hepatic Cytochrome P450 Expression: Effects of 16-Week High-Fat Diet and Exercise. Toxicological Sciences, 2018, 162, 309-317. | 3.1 | 20 |

| # | Article | IF | Citations |
|----|--|-----|-----------|
| 19 | The effects of sewage treatment plant effluents on hepatic and intestinal biomarkers in common carp (Cyprinus carpio). Science of the Total Environment, 2018, 635, 1160-1169. | 8.0 | 23 |
| 20 | Classification of wooden breast myopathy in chicken <i>pectoralis major</i> by a standardised method and association with conventional quality assessments. International Journal of Food Science and Technology, 2018, 53, 1744-1752. | 2.7 | 58 |
| 21 | In vitro effects of rebaudioside A, stevioside and steviol on porcine cytochrome p450 expression and activity. Food Chemistry, 2018, 258, 245-253. | 8.2 | 11 |
| 22 | Co-treatment with indole-3-carbinol and resveratrol modify porcine CYP1A and CYP3A activities and expression. Xenobiotica, 2018, 48, 232-240. | 1.1 | 6 |
| 23 | Impact of fasting followed by short-term exposure to interleukin-6 on cytochrome P450 mRNA in mice. Toxicology Letters, 2018, 282, 93-99. | 0.8 | 21 |
| 24 | In vitro differentiation of progenitor cells isolated from juvenile pig hearts – expression of relevant gene and protein markers. Scandinavian Cardiovascular Journal, 2018, 52, 34-42. | 1.2 | 0 |
| 25 | Activation of the aryl hydrocarbon receptor decreases rifampicin-induced CYP3A4 expression in primary human hepatocytes and HepaRG. Toxicology Letters, 2017, 277, 1-8. | 0.8 | 35 |
| 26 | Whole Milk Increases Intestinal <i>ANGPTL4</i> Expression and Excretion of Fatty Acids through Feces and Urine. Journal of Agricultural and Food Chemistry, 2017, 65, 281-290. | 5.2 | 6 |
| 27 | Constitutive expression and activity of cytochrome P450 in conventional pigs. Research in Veterinary Science, 2017, 111, 75-80. | 1.9 | 22 |
| 28 | Induction of cytochrome P450 mRNA in porcine primary hepatocytes cultured under serum free conditions: Comparison of freshly isolated cells and cryopreserved. Experimental Cell Research, 2017, 360, 218-225. | 2.6 | 11 |
| 29 | Taste receptors in the gut – A new target for health promoting properties in diet. Food Research International, 2017, 100, 1-8. | 6.2 | 29 |
| 30 | High resolution magic angle spinning NMR spectroscopy reveals that pectoralis muscle dystrophy in chicken is associated with reduced muscle content of anserine and carnosine. Food Chemistry, 2017, 217, 151-154. | 8.2 | 47 |
| 31 | Comparison of xenobiotic-metabolising human, porcine, rodent, and piscine cytochrome P450. Toxicology, 2017, 375, 10-27. | 4.2 | 68 |
| 32 | Skatole (3-Methylindole) Is a Partial Aryl Hydrocarbon Receptor Agonist and Induces CYP1A1/2 and CYP1B1 Expression in Primary Human Hepatocytes. PLoS ONE, 2016, 11, e0154629. | 2.5 | 50 |
| 33 | Constitutive expression of cytochrome P450 in foetal and adult porcine livers—Effects of body weight. Toxicology Letters, 2016, 258, 87-92. | 0.8 | 10 |
| 34 | Angiotensin l–converting enzyme–inhibitory peptides from bovine collagen: insights into inhibitory mechanism and transepithelial transport. Food Research International, 2016, 89, 373-381. | 6.2 | 70 |
| 35 | Chestnut wood extract in boar diet reduces intestinal skatole production, a boar taint compound. Agronomy for Sustainable Development, 2016, 36, 1. | 5.3 | 13 |
| 36 | Tissue-specific regulation of CYP3A by hydrolysable tannins in male pigs. Xenobiotica, 2016, 46, 591-596. | 1.1 | 4 |

| # | Article | IF | Citations |
|----|---|-----|-----------|
| 37 | Gender-related differences in the formation of skatole metabolites by specific CYP450 in porcine hepatic S9 fractions. Animal, 2015, 9, 635-642. | 3.3 | 11 |
| 38 | Immunocastration of Male Pigs – Situation Today. Procedia Food Science, 2015, 5, 324-327. | 0.6 | 21 |
| 39 | <i>In Vitro</i> Gender-Dependent Inhibition of Porcine Cytochrome P450 Activity by Selected Flavonoids and Phenolic Acids. BioMed Research International, 2015, 2015, 1-7. | 1.9 | 29 |
| 40 | Does dexamethasone affect hepatic CYP450 system of fish? Semi-static in-vivo experiment on juvenile rainbow trout. Chemosphere, 2015, 139, 155-162. | 8.2 | 12 |
| 41 | Regulation of Porcine Hepatic Cytochrome P450 — Implication for Boar Taint. Computational and Structural Biotechnology Journal, 2014, 11, 106-112. | 4.1 | 30 |
| 42 | Regulation of cytochrome P450 mRNA expression in primary porcine hepatocytes by selected secondary plant metabolites from chicory (Cichorium intybus L.). Food Chemistry, 2014, 146, 255-263. | 8.2 | 32 |
| 43 | Regulation of 3β-hydroxysteroid dehydrogenase and sulphotransferase 2A1 gene expression in primary porcine hepatocytes by selected sex-steroids and plant secondary metabolites from chicory (Cichorium intybus L.) and wormwood (Artemisia sp.). Gene, 2014, 536, 53-58. | 2.2 | 11 |
| 44 | Comparable constitutive expression and activity of cytochrome P450 between the lobes of the porcine liver. Toxicology in Vitro, 2014, 28, 1190-1195. | 2.4 | 4 |
| 45 | Regulation of 3β-Hydroxysteroid Dehydrogenase/Δ5-Δ4 Isomerase: A Review. International Journal of Molecular Sciences, 2013, 14, 17926-17942. | 4.1 | 65 |
| 46 | Expression and activities of hepatic cytochrome P450 (CYP1A, CYP2A and CYP2E1) in entire and castrated male pigs. Animal, 2012, 6, 271-277. | 3.3 | 34 |
| 47 | Dried chicory root modifies the activity and expression of porcine hepatic CYP3A but not 2C – Effect of in vitro and in vivo exposure. Food and Chemical Toxicology, 2012, 50, 4175-4179. | 3.6 | 12 |
| 48 | Improvac does not modify the expression and activities of the major drug metabolizing enzymes cytochrome P450 3A and 2C in pigs. Vaccine, 2012, 30, 3515-3518. | 3.8 | 6 |
| 49 | Feeding dried chicory root to pigs decrease androstenone accumulation in fat by increasing hepatic $3\hat{l}^2$ hydroxysteroid dehydrogenase expression. Journal of Steroid Biochemistry and Molecular Biology, 2012, 130, 90-95. | 2.5 | 24 |
| 50 | Expression of hepatic $3\hat{l}^2$ -hydroxysteroid dehydrogenase and sulfotransferase 2A1 in entire and castrated male pigs. Molecular Biology Reports, 2012, 39, 7927-7932. | 2.3 | 13 |
| 51 | Comparison of cytochrome P450 concentrations and metabolic activities in porcine hepatic microsomes prepared with two different methods. Toxicology in Vitro, 2011, 25, 343-346. | 2.4 | 50 |
| 52 | In vivo effect of dried chicory root (Cichorium intybus L.) on xenobiotica metabolising cytochrome P450 enzymes in porcine liver. Toxicology Letters, 2011, 200, 88-91. | 0.8 | 43 |
| 53 | In vitro inhibition of porcine cytochrome P450 by 17β-estradiol and 17α-estradiol. Interdisciplinary Toxicology, 2011, 4, 78-84. | 1.0 | 5 |
| 54 | <i>In Vitro</i> Cytochrome P450 2E1 and 2A Activities in the Presence of Testicular Steroids. Reproduction in Domestic Animals, 2011, 46, 149-154. | 1.4 | 22 |

| # | Article | lF | CITATIONS |
|----|---|-----|-----------|
| 55 | Genderâ€related Differences in Cytochrome P450 in Porcine Liver – Implication for Activity, Expression and Inhibition by Testicular Steroids. Reproduction in Domestic Animals, 2011, 46, 616-623. | 1.4 | 61 |
| 56 | Exercise-induced regulation of muscular Na ⁺ -K ⁺ pump, FXYD1, and NHE1 mRNA and protein expression: importance of training status, intensity, and muscle type. American Journal of Physiology - Regulatory Integrative and Comparative Physiology, 2011, 300, R1209-R1220. | 1.8 | 14 |
| 57 | Na+–K+ pump location and translocation during muscle contraction in rat skeletal muscle. Pflugers Archiv European Journal of Physiology, 2008, 456, 979-989. | 2.8 | 30 |
| 58 | Exerciseâ€induced regulation of phospholemman (FXYD1) in rat skeletal muscle: implications for Na ⁺ /K ⁺ â€ATPase activity. Acta Physiologica, 2008, 194, 67-79. | 3.8 | 40 |
| 59 | Reply to Bishop and Schneiker. American Journal of Physiology - Regulatory Integrative and Comparative Physiology, 2007, 293, R1460-R1460. | 1.8 | 1 |
| 60 | Effect of two different intense training regimens on skeletal muscle ion transport proteins and fatigue development. American Journal of Physiology - Regulatory Integrative and Comparative Physiology, 2007, 292, R1594-R1602. | 1.8 | 171 |