

Hirohito Metoki

List of Publications by Citations

Source: <https://exaly.com/author-pdf/3302657/hirohito-metoki-publications-by-citations.pdf>

Version: 2024-04-26

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

265
papers

9,796
citations

44
h-index

93
g-index

290
ext. papers

11,510
ext. citations

3.6
avg, IF

5.35
L-index

#	Paper	IF	Citations
265	Association of estimated glomerular filtration rate and albuminuria with all-cause and cardiovascular mortality in general population cohorts: a collaborative meta-analysis. <i>Lancet, The</i> , 2010 , 375, 2073-81	40	2495
264	Prognosis of "masked" hypertension and "white-coat" hypertension detected by 24-h ambulatory blood pressure monitoring 10-year follow-up from the Ohasama study. <i>Journal of the American College of Cardiology</i> , 2005 , 46, 508-15	15.1	459
263	Prognostic significance for stroke of a morning pressor surge and a nocturnal blood pressure decline: the Ohasama study. <i>Hypertension</i> , 2006 , 47, 149-54	8.5	327
262	Ambulatory blood pressure and 10-year risk of cardiovascular and noncardiovascular mortality: the Ohasama study. <i>Hypertension</i> , 2005 , 45, 240-5	8.5	320
261	Day-by-day variability of blood pressure and heart rate at home as a novel predictor of prognosis: the Ohasama study. <i>Hypertension</i> , 2008 , 52, 1045-50	8.5	313
260	Prognostic value of the morning blood pressure surge in 5645 subjects from 8 populations. <i>Hypertension</i> , 2010 , 55, 1040-8	8.5	205
259	How many times should blood pressure be measured at home for better prediction of stroke risk? Ten-year follow-up results from the Ohasama study. <i>Journal of Hypertension</i> , 2004 , 22, 1099-104	1.9	201
258	Kidney dysfunction as a risk factor for first symptomatic stroke events in a general Japanese population--the Ohasama study. <i>Nephrology Dialysis Transplantation</i> , 2007 , 22, 1910-5	4.3	155
257	The Tohoku Medical Megabank Project: Design and Mission. <i>Journal of Epidemiology</i> , 2016 , 26, 493-511	3.4	141
256	Global Cardiovascular and Renal Outcomes of Reduced GFR. <i>Journal of the American Society of Nephrology: JASN</i> , 2017 , 28, 2167-2179	12.7	127
255	Cardiovascular outcomes in the first trial of antihypertensive therapy guided by self-measured home blood pressure. <i>Hypertension Research</i> , 2012 , 35, 1102-10	4.7	126
254	Prediction of stroke by home "morning" versus "evening" blood pressure values: the Ohasama study. <i>Hypertension</i> , 2006 , 48, 737-43	8.5	124
253	Ambulatory arterial stiffness index and 24-hour ambulatory pulse pressure as predictors of mortality in Ohasama, Japan. <i>Stroke</i> , 2007 , 38, 1161-6	6.7	115
252	White-coat hypertension as a risk factor for the development of home hypertension: the Ohasama study. <i>Archives of Internal Medicine</i> , 2005 , 165, 1541-6		114
251	Prediction of stroke by self-measurement of blood pressure at home versus casual screening blood pressure measurement in relation to the Joint National Committee 7 classification: the Ohasama study. <i>Stroke</i> , 2004 , 35, 2356-61	6.7	102
250	Control of blood pressure as measured at home and office, and comparison with physicians' assessment of control among treated hypertensive patients in Japan: First Report of the Japan Home versus Office Blood Pressure Measurement Evaluation (J-HOME) study. <i>Hypertension Research</i> , 2004 , 27, 755-63	4.7	100
249	Home blood pressure variability as cardiovascular risk factor in the population of Ohasama. <i>Hypertension</i> , 2013 , 61, 61-9	8.5	95

248	Outline of Definition and Classification of Pregnancy induced Hypertension (PIH) <i>Hypertension Research in Pregnancy</i> , 2013 , 1, 3-4	0.7	87
247	Isolated uncontrolled hypertension at home and in the office among treated hypertensive patients from the J-HOME study. <i>Journal of Hypertension</i> , 2005 , 23, 1653-60	1.9	84
246	Association of (pro)renin receptor gene polymorphism with blood pressure in Japanese men: the Ohasama study. <i>American Journal of Hypertension</i> , 2009 , 22, 294-9	2.3	75
245	Detection of carotid atherosclerosis in individuals with masked hypertension and white-coat hypertension by self-measured blood pressure at home: the Ohasama study. <i>Journal of Hypertension</i> , 2007 , 25, 321-7	1.9	75
244	Enhanced radial late systolic pressure augmentation in hypertensive patients with left ventricular hypertrophy. <i>American Journal of Hypertension</i> , 2006 , 19, 27-32	2.3	72
243	Prognostic value of home heart rate for cardiovascular mortality in the general population The Ohasama study. <i>American Journal of Hypertension</i> , 2004 , 17, 1005-1010	2.3	69
242	Prognostic significance of night-time, early morning, and daytime blood pressures on the risk of cerebrovascular and cardiovascular mortality: the Ohasama Study. <i>Journal of Hypertension</i> , 2006 , 24, 1841-8	1.9	65
241	Prognostic value of home heart rate for cardiovascular mortality in the general population: the Ohasama study. <i>American Journal of Hypertension</i> , 2004 , 17, 1005-10	2.3	64
240	Ambulatory blood pressure, blood pressure variability and the prevalence of carotid artery alteration: the Ohasama study. <i>Journal of Hypertension</i> , 2007 , 25, 1704-10	1.9	61
239	Characteristics of resistant hypertension determined by self-measured blood pressure at home and office blood pressure measurements: the J-HOME study. <i>Journal of Hypertension</i> , 2006 , 24, 1737-43	1.9	61
238	Predictive value of ambulatory heart rate in the Japanese general population: the Ohasama study. <i>Journal of Hypertension</i> , 2008 , 26, 1571-6	1.9	59
237	Gene expression of (pro)renin receptor is upregulated in hearts and kidneys of rats with congestive heart failure. <i>Peptides</i> , 2009 , 30, 2316-22	3.8	58
236	Risk stratification by self-measured home blood pressure across categories of conventional blood pressure: a participant-level meta-analysis. <i>PLoS Medicine</i> , 2014 , 11, e1001591	11.6	57
235	Long-Term Stroke Risk Due to Partial White-Coat or Masked Hypertension Based on Home and Ambulatory Blood Pressure Measurements: The Ohasama Study. <i>Hypertension</i> , 2016 , 67, 48-55	8.5	55
234	Ambulatory versus home versus clinic blood pressure: the association with subclinical cerebrovascular diseases: the Ohasama Study. <i>Hypertension</i> , 2012 , 59, 22-8	8.5	54
233	Relative risks of chronic kidney disease for mortality and end-stage renal disease across races are similar. <i>Kidney International</i> , 2014 , 86, 819-27	9.9	52
232	Stroke risk and antihypertensive drug treatment in the general population: the Japan arteriosclerosis longitudinal study. <i>Journal of Hypertension</i> , 2009 , 27, 357-64	1.9	51
231	Prevalence of masked uncontrolled and treated white-coat hypertension defined according to the average of morning and evening home blood pressure value: from the Japan Home versus Office Measurement Evaluation Study. <i>Blood Pressure Monitoring</i> , 2005 , 10, 311-6	1.3	51

230	Day-to-day variability in home blood pressure is associated with cognitive decline: the Ohasama study. <i>Hypertension</i> , 2014 , 63, 1333-8	8.5	50
229	Factors associated with day-by-day variability of self-measured blood pressure at home: the Ohasama study. <i>American Journal of Hypertension</i> , 2010 , 23, 980-6	2.3	49
228	Use of 2003 European Society of Hypertension-European Society of Cardiology guidelines for predicting stroke using self-measured blood pressure at home: the Ohasama study. <i>European Heart Journal</i> , 2005 , 26, 2026-31	9.5	49
227	Cost-effectiveness of the introduction of home blood pressure measurement in patients with office hypertension. <i>Journal of Hypertension</i> , 2008 , 26, 685-90	1.9	48
226	Association of arterial stiffness with silent cerebrovascular lesions: the Ohasama study. <i>Cerebrovascular Diseases</i> , 2011 , 31, 329-37	3.2	47
225	Seasonal trends of blood pressure during pregnancy in Japan: the babies and their parents' longitudinal observation in Suzuki Memorial Hospital in Intrauterine Period study. <i>Journal of Hypertension</i> , 2008 , 26, 2406-13	1.9	47
224	Angiotensin-converting enzyme I/D polymorphism and hypertension: the Ohasama study. <i>Journal of Hypertension</i> , 2002 , 20, 1121-6	1.9	47
223	Fruit and vegetable consumption and the risk of hypertension determined by self measurement of blood pressure at home: the Ohasama study. <i>Hypertension Research</i> , 2008 , 31, 1435-43	4.7	45
222	Prediction of ischaemic and haemorrhagic stroke by self-measured blood pressure at home: the Ohasama study. <i>Blood Pressure Monitoring</i> , 2004 , 9, 315-20	1.3	45
221	Reproducibility of nocturnal blood pressure assessed by self-measurement of blood pressure at home. <i>Hypertension Research</i> , 2007 , 30, 707-12	4.7	44
220	The economic impact of the introduction of home blood pressure measurement for the diagnosis and treatment of hypertension. <i>Blood Pressure Monitoring</i> , 2006 , 11, 257-67	1.3	43
219	Cohort Profile: Tohoku Medical Megabank Project Birth and Three-Generation Cohort Study (TMM BirThree Cohort Study): rationale, progress and perspective. <i>International Journal of Epidemiology</i> , 2020 , 49, 18-19m	7.8	43
218	Aldosterone synthase gene (CYP11B2) C-334T polymorphism, ambulatory blood pressure and nocturnal decline in blood pressure in the general Japanese population: the Ohasama Study. <i>Journal of Hypertension</i> , 2001 , 19, 2179-84	1.9	41
217	Stroke risk in treated hypertension based on home blood pressure: the Ohasama study. <i>American Journal of Hypertension</i> , 2010 , 23, 508-14	2.3	40
216	Psychological distress during pregnancy in Miyagi after the Great East Japan Earthquake: The Japan Environment and Children's Study. <i>Journal of Affective Disorders</i> , 2016 , 190, 341-348	6.6	39
215	Association of microalbuminuria with brachial-ankle pulse wave velocity: the Ohasama study. <i>American Journal of Hypertension</i> , 2008 , 21, 413-8	2.3	39
214	Plasma fibrinogen, ambulatory blood pressure, and silent cerebrovascular lesions: the Ohasama study. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2007 , 27, 963-8	9.4	39
213	Evidence-based guidelines for treatment of uterine body neoplasm in Japan: Japan Society of Gynecologic Oncology (JSGO) 2009 edition. <i>International Journal of Clinical Oncology</i> , 2010 , 15, 531-42	4.2	38

212	Association of (pro)renin receptor gene polymorphisms with lacunar infarction and left ventricular hypertrophy in Japanese women: the Ohasama study. <i>Hypertension Research</i> , 2011 , 34, 530-5	4.7	37
211	Association of environmental tobacco smoke exposure with elevated home blood pressure in Japanese women: the Ohasama study. <i>Journal of Hypertension</i> , 2010 , 28, 1814-20	1.9	37
210	Predicting stroke using 4 ambulatory blood pressure monitoring-derived blood pressure indices: the Ohasama Study. <i>Hypertension</i> , 2006 , 48, 877-82	8.5	37
209	Masked hypertension determined by self-measured blood pressure at home and chronic kidney disease in the Japanese general population: the Ohasama study. <i>Hypertension Research</i> , 2008 , 31, 2129-35	4.7	36
208	Increased expression of (pro)renin receptor in the remnant kidneys of 5/6 nephrectomized rats. <i>Regulatory Peptides</i> , 2010 , 159, 93-9		35
207	Pre-hypertension as a significant predictor of chronic kidney disease in a general population: the Ohasama Study. <i>Nephrology Dialysis Transplantation</i> , 2012 , 27, 3218-23	4.3	35
206	Efficacy and duration of action of the four selective angiotensin II subtype 1 receptor blockers, losartan, candesartan, valsartan and telmisartan, in patients with essential hypertension determined by home blood pressure measurements. <i>Clinical and Experimental Hypertension</i> , 2005 , 27, 477-89	2.2	35
205	Usefulness of assessing masked and white-coat hypertension by ambulatory blood pressure monitoring for determining prevalent risk of chronic kidney disease: the Ohasama study. <i>Hypertension Research</i> , 2010 , 33, 1192-8	4.7	34
204	Association between tooth loss and cognitive impairment in community-dwelling older Japanese adults: a 4-year prospective cohort study from the Ohasama study. <i>BMC Oral Health</i> , 2018 , 18, 142	3.7	33
203	Serum magnesium, ambulatory blood pressure, and carotid artery alteration: the Ohasama study. <i>American Journal of Hypertension</i> , 2010 , 23, 1292-8	2.3	33
202	Blood pressure measured in the clinic and at home during pregnancy among nulliparous and multiparous women: the BOSHI study. <i>American Journal of Hypertension</i> , 2013 , 26, 141-8	2.3	31
201	The association between masked hypertension and waist circumference as an obesity-related anthropometric index for metabolic syndrome: the Ohasama study. <i>Hypertension Research</i> , 2009 , 32, 438-43	4.7	31
200	High fruit intake is associated with a lower risk of future hypertension determined by home blood pressure measurement: the OHASAMA study. <i>Journal of Human Hypertension</i> , 2011 , 25, 164-71	2.6	31
199	Increased expression of adrenomedullin 2/intermedin in rat hearts with congestive heart failure. <i>European Journal of Heart Failure</i> , 2008 , 10, 840-9	12.3	31
198	CYP11B2 polymorphisms and home blood pressure in a population-based cohort in Japanese: the Ohasama study. <i>Hypertension Research</i> , 2004 , 27, 1-6	4.7	31
197	Association Between Amplitude of Seasonal Variation in Self-Measured Home Blood Pressure and Cardiovascular Outcomes: HOMED-BP (Hypertension Objective Treatment Based on Measurement By Electrical Devices of Blood Pressure) Study. <i>Journal of the American Heart Association</i> , 2018 , 7,	6	30
196	Association of kidney dysfunction with silent lacunar infarcts and white matter hyperintensity in the general population: the Ohasama study. <i>Cerebrovascular Diseases</i> , 2010 , 30, 43-50	3.2	30
195	Practice and awareness of physicians regarding home blood pressure measurement in Japan. <i>Hypertension Research</i> , 2010 , 33, 428-34	4.7	30

194	Accumulation of common polymorphisms is associated with development of hypertension: a 12-year follow-up from the Ohasama study. <i>Hypertension Research</i> , 2010 , 33, 129-34	4.7	30
193	Night-time blood pressure is associated with the development of chronic kidney disease in a general population: the Ohasama Study. <i>Journal of Hypertension</i> , 2013 , 31, 2410-7	1.9	29
192	Associations between day-by-day variability in blood pressure measured at home and antihypertensive drugs: the J-HOME-Morning study. <i>Clinical and Experimental Hypertension</i> , 2012 , 34, 297-304	2.2	29
191	Genome-wide response to antihypertensive medication using home blood pressure measurements: a pilot study nested within the HOMED-BP study. <i>Pharmacogenomics</i> , 2013 , 14, 1709-21	2.6	28
190	Seasonal variation in self-measured home blood pressure among patients on antihypertensive medications: HOMED-BP study. <i>Hypertension Research</i> , 2017 , 40, 284-290	4.7	28
189	Animal protein intake is associated with higher-level functional capacity in elderly adults: the Ohasama study. <i>Journal of the American Geriatrics Society</i> , 2014 , 62, 426-34	5.6	27
188	Breastfeeding leads to lower blood pressure in 7-year-old Japanese children: Tohoku Study of Child Development. <i>Hypertension Research</i> , 2013 , 36, 117-22	4.7	27
187	Home blood pressure level, blood pressure variability, smoking, and stroke risk in Japanese men: the Ohasama study. <i>American Journal of Hypertension</i> , 2012 , 25, 883-91	2.3	27
186	Health behaviors as predictors for declines in higher-level functional capacity in older adults: the Ohasama study. <i>Journal of the American Geriatrics Society</i> , 2011 , 59, 1993-2000	5.6	26
185	Practice and awareness of physicians regarding casual-clinic blood pressure measurement in Japan. <i>Hypertension Research</i> , 2010 , 33, 960-4	4.7	26
184	Stroke risk of blood pressure indices determined by home blood pressure measurement: the Ohasama study. <i>Stroke</i> , 2009 , 40, 2859-61	6.7	26
183	Plasma renin activity and the aldosterone-to-renin ratio are associated with the development of chronic kidney disease: the Ohasama Study. <i>Journal of Hypertension</i> , 2012 , 30, 1632-8	1.9	24
182	Study Profile of the Tohoku Medical Megabank Community-Based Cohort Study. <i>Journal of Epidemiology</i> , 2021 , 31, 65-76	3.4	24
181	Predictive value for mortality of the double product at rest obtained by home blood pressure measurement: the Ohasama study. <i>American Journal of Hypertension</i> , 2012 , 25, 568-75	2.3	23
180	Daily serial hemodynamic data during pregnancy and seasonal variation: the BOSHI study. <i>Clinical and Experimental Hypertension</i> , 2012 , 34, 290-6	2.2	22
179	Current usage of diuretics among hypertensive patients in Japan: the Japan Home versus Office Blood Pressure Measurement Evaluation (J-HOME) study. <i>Hypertension Research</i> , 2006 , 29, 857-63	4.7	22
178	Outline of the new definition and classification of Hypertensive Disorders of Pregnancy (HDP) a revised JSSHP statement of 2005. <i>Hypertension Research in Pregnancy</i> , 2018 , 6, 33-37	0.7	22
177	Optimal cutoff point of waist circumference and use of home blood pressure as a definition of metabolic syndrome: the Ohasama study. <i>American Journal of Hypertension</i> , 2008 , 21, 514-20	2.3	21

176	Stroke risk in systolic and combined systolic and diastolic hypertension determined using ambulatory blood pressure. The Ohasama study. <i>American Journal of Hypertension</i> , 2007 , 20, 1125-31	2.3	21
175	Drug Use before and during Pregnancy in Japan: The Japan Environment and Children's Study. <i>Pharmacy (Basel, Switzerland)</i> , 2017 , 5,	2	20
174	Genotypes of the betaENaC gene have little influence on blood pressure level in the Japanese population. <i>American Journal of Hypertension</i> , 2002 , 15, 189-92	2.3	20
173	Association between social capital and the prevalence of gestational diabetes mellitus: An interim report of the Japan Environment and Children's Study. <i>Diabetes Research and Clinical Practice</i> , 2016 , 120, 132-41	7.4	19
172	Salt-inducible kinase 1 influences Na(+),K(+)-ATPase activity in vascular smooth muscle cells and associates with variations in blood pressure. <i>Journal of Hypertension</i> , 2011 , 29, 2395-403	1.9	19
171	Influence of alcohol intake on circadian blood pressure variation in Japanese men: the Ohasama study. <i>American Journal of Hypertension</i> , 2009 , 22, 1171-6	2.3	19
170	Aldosterone-to-renin ratio as a predictor of stroke under conditions of high sodium intake: the Ohasama study. <i>American Journal of Hypertension</i> , 2012 , 25, 777-83	2.3	19
169	Reference values and associated factors for Japanese newborns' blood pressure and pulse rate: the babies' and their parents' longitudinal observation in Suzuki Memorial Hospital on intrauterine period (BOSHI) study. <i>Journal of Hypertension</i> , 2016 , 34, 1578-85	1.9	18
168	Home blood pressure measurements associated with better blood pressure control: the J-HOME study. <i>Journal of Human Hypertension</i> , 2008 , 22, 197-204	2.6	18
167	The second progress report on the Hypertension Objective treatment based on Measurement by Electrical Devices of Blood Pressure (HOMED-BP) study. <i>Blood Pressure Monitoring</i> , 2004 , 9, 243-7	1.3	18
166	Aldosterone-to-renin ratio and nocturnal blood pressure decline in a general population: the Ohasama study. <i>Journal of Hypertension</i> , 2011 , 29, 1940-7	1.9	17
165	Detection of silent cerebrovascular lesions in individuals with 'masked' and 'white-coat' hypertension by home blood pressure measurement: the Ohasama study. <i>Journal of Hypertension</i> , 2009 , 27, 1049-55	1.9	17
164	The current status of home and office blood pressure control among hypertensive patients with diabetes mellitus: the Japan Home Versus Office Blood Pressure Measurement Evaluation (J-HOME) study. <i>Diabetes Research and Clinical Practice</i> , 2006 , 73, 276-83	7.4	17
163	Incidence of Domestic Violence Against Pregnant Females After the Great East Japan Earthquake in Miyagi Prefecture: The Japan Environment and Children's Study. <i>Disaster Medicine and Public Health Preparedness</i> , 2017 , 11, 216-226	2.8	16
162	New definition and classification of Hypertensive Disorders of Pregnancy (HDP). <i>Hypertension Research in Pregnancy</i> , 2019 , 7, 1-5	0.7	16
161	Eczema and Asthma Symptoms among Schoolchildren in Coastal and Inland Areas after the 2011 Great East Japan Earthquake: The ToMMo Child Health Study. <i>Tohoku Journal of Experimental Medicine</i> , 2015 , 237, 297-305	2.4	16
160	Expression of adrenomedullin 2/intermedin, a possible reno-protective peptide, is decreased in the kidneys of rats with hypertension or renal failure. <i>American Journal of Physiology - Renal Physiology</i> , 2010 , 299, F128-34	4.3	16
159	Uncontrolled hypertension based on morning and evening home blood pressure measurements from the J-HOME study. <i>Hypertension Research</i> , 2009 , 32, 1072-8	4.7	16

158	Aldosterone-to-renin ratio and home blood pressure in subjects with higher and lower sodium intake: the Ohasama study. <i>Hypertension Research</i> , 2011 , 34, 361-6	4.7	16
157	Repeated evening home blood pressure measurement improves prognostic significance for stroke: a 12-year follow-up of the Ohasama study. <i>Blood Pressure Monitoring</i> , 2009 , 14, 93-8	1.3	16
156	Risk Factors for Stroke among Young-Old and Old-Old Community-Dwelling Adults in Japan: The Ohasama Study. <i>Journal of Atherosclerosis and Thrombosis</i> , 2017 , 24, 290-300	4	15
155	Maternal clinic and home blood pressure measurements during pregnancy and infant birth weight: the BOSHI study. <i>Hypertension Research</i> , 2016 , 39, 151-7	4.7	15
154	Pregnancy outcomes of gestational diabetes mellitus according to pre-gestational BMI in a retrospective multi-institutional study in Japan. <i>Endocrine Journal</i> , 2014 , 61, 373-80	2.9	15
153	The velocity of antihypertensive effect of losartan/hydrochlorothiazide and angiotensin II receptor blocker. <i>Journal of Hypertension</i> , 2012 , 30, 1478-86	1.9	15
152	Prognostic significance of morning surge in blood pressure: which definition, which outcome?. <i>Blood Pressure Monitoring</i> , 2008 , 13, 161-2	1.3	15
151	Prevalence and determinants of inadequate use of folic acid supplementation in Japanese pregnant women: the Japan Environment and Children's Study (JECS). <i>Journal of Maternal-Fetal and Neonatal Medicine</i> , 2017 , 30, 588-593	2	14
150	Living situations associated with poor dietary intake among healthy Japanese elderly: the Ohasama Study. <i>Journal of Nutrition, Health and Aging</i> , 2015 , 19, 375-82	5.2	14
149	Parity as a factor affecting the white-coat effect in pregnant women: the BOSHI study. <i>Hypertension Research</i> , 2015 , 38, 770-5	4.7	14
148	Aldosterone-to-renin ratio and nocturnal blood pressure decline assessed by self-measurement of blood pressure at home: the Ohasama Study. <i>Clinical and Experimental Hypertension</i> , 2014 , 36, 108-14	2.2	14
147	Diurnal blood pressure variation and cardiovascular prognosis in a community-based study of Ohasama, Japan. <i>Hypertension Research</i> , 2010 , 33, 652-6	4.7	14
146	Proposal of a risk-stratification system for the Japanese population based on blood pressure levels: the Ohasama study. <i>Hypertension Research</i> , 2008 , 31, 1315-22	4.7	14
145	Impact of COVID-19 pandemic on glycemic control among outpatients with type 2 diabetes in Japan: A hospital-based survey from a country without lockdown. <i>Diabetes Research and Clinical Practice</i> , 2021 , 176, 108840	7.4	14
144	Impaired Higher-Level Functional Capacity as a Predictor of Stroke in Community-Dwelling Older Adults: The Ohasama Study. <i>Stroke</i> , 2016 , 47, 323-8	6.7	13
143	Introversion associated with large differences between screening blood pressure and home blood pressure measurement: The Ohasama study. <i>Journal of Hypertension</i> , 2006 , 24, 2183-9	1.9	13
142	Haplotypes of aldosterone synthase (CYP11B2) gene in the general population of Japan: the Ohasama study. <i>Clinical and Experimental Hypertension</i> , 2001 , 23, 603-10	2.2	13
141	T+31C polymorphism (M235T) of the angiotensinogen gene and home blood pressure in the Japanese general population: the Ohasama Study. <i>Hypertension Research</i> , 2003 , 26, 47-52	4.7	13

140	Beneficial Effects of Human Papillomavirus Vaccine for Prevention of Cervical Abnormalities in Miyagi, Japan. <i>Tohoku Journal of Experimental Medicine</i> , 2016 , 240, 147-151	2.4	13
139	Severity of eczema and mental health problems in Japanese schoolchildren: The ToMMo Child Health Study. <i>Allergology International</i> , 2018 , 67, 481-486	4.4	12
138	New definition and classification of Hypertensive Disorders of Pregnancy (HDP) <i>Hypertension Research in Pregnancy</i> , 2017 , 5, 39-40	0.7	12
137	Relationship of dysregulation of glucose metabolism with white-coat hypertension: the Ohasama study. <i>Hypertension Research</i> , 2010 , 33, 937-43	4.7	12
136	T+31C polymorphism of angiotensinogen gene and nocturnal blood pressure decline: the Ohasama study. <i>American Journal of Hypertension</i> , 2002 , 15, 628-32	2.3	12
135	Association between alcohol consumption during pregnancy and hypertensive disorders of pregnancy in Japan: the Japan Environment and Children's Study. <i>Hypertension Research</i> , 2019 , 42, 85-94	4.7	12
134	The risk of secondary sex ratio imbalance and increased monozygotic twinning after blastocyst transfer: data from the Japan Environment and Children's Study. <i>Reproductive Biology and Endocrinology</i> , 2019 , 17, 27	5	11
133	Protocol and Research Perspectives of the ToMMo Child Health Study after the 2011 Great East Japan Earthquake. <i>Tohoku Journal of Experimental Medicine</i> , 2015 , 236, 123-30	2.4	11
132	Association between N-terminal pro B-type natriuretic peptide and day-to-day blood pressure and heart rate variability in a general population: the Ohasama study. <i>Journal of Hypertension</i> , 2015 , 33, 1536-41	1.9	11
131	Validation of the FM-800 ambulatory blood pressure monitor according to the Association for the Advancement of Medical Instrumentation criteria and the International Protocol. <i>Clinical and Experimental Hypertension</i> , 2010 , 32, 523-7	2.2	11
130	Difference between home and office blood pressures among treated hypertensive patients from the Japan Home versus Office Blood Pressure Measurement Evaluation (J-HOME) study. <i>Hypertension Research</i> , 2008 , 31, 1115-23	4.7	11
129	Development and evaluation of a home nocturnal blood pressure monitoring system using a wrist-cuff device. <i>Blood Pressure Monitoring</i> , 2018 , 23, 318-326	1.3	11
128	Potential identification of vitamin B6 responsiveness in autism spectrum disorder utilizing phenotype variables and machine learning methods. <i>Scientific Reports</i> , 2018 , 8, 14840	4.9	11
127	Mother-offspring aggregation in home versus conventional blood pressure in the Tohoku Study of Child Development (TSCD). <i>Acta Cardiologica</i> , 2012 , 67, 449-56	0.9	10
126	Influence of adrenomedullin 2/intermedin gene polymorphism on blood pressure, renal function and silent cerebrovascular lesions in Japanese: the Ohasama study. <i>Hypertension Research</i> , 2011 , 34, 1327-32	4.7	10
125	Influence of home blood pressure measuring conditions in the evening on the morning-evening home blood pressure difference in treated hypertensive patients: the J-HOME study. <i>Blood Pressure Monitoring</i> , 2009 , 14, 160-5	1.3	10
124	The velocity of antihypertensive effects of seven angiotensin II receptor blockers determined by home blood pressure measurements. <i>Journal of Hypertension</i> , 2016 , 34, 1218-23	1.9	10
123	Menstrual Factors and Stroke Incidence in Japanese Postmenopausal Women: The Ohasama Study. <i>Neuroepidemiology</i> , 2016 , 47, 109-116	5.4	9

122	A retrospective multi-institutional study of treatment for mild gestational diabetes in Japan. <i>Diabetes Research and Clinical Practice</i> , 2014 , 103, 412-8	7.4	9
121	Self-monitoring of ambulatory blood pressure by the Microlife WatchBP O3--an application test. <i>Clinical and Experimental Hypertension</i> , 2011 , 33, 34-40	2.2	9
120	Blood pressure changes during pregnancy. <i>Hypertension Research</i> , 2012 , 35, 563-4; author reply 34	4.7	9
119	Efficacy of Combination Antihypertensive Therapy with Low-Dose Indapamide: Assessment by Blood Pressure Self-Monitoring at Home. <i>Clinical and Experimental Hypertension</i> , 2005 , 27, 331-341	2.2	9
118	Clustering by phenotype and genome-wide association study in autism. <i>Translational Psychiatry</i> , 2020 , 10, 290	8.6	9
117	Nocturnal blood pressure decline based on different time intervals and long-term cardiovascular risk: the Ohasama Study. <i>Clinical and Experimental Hypertension</i> , 2018 , 40, 1-7	2.2	9
116	Association of aldosterone-to-renin ratio with hypertension differs by sodium intake: the Ohasama study. <i>American Journal of Hypertension</i> , 2015 , 28, 208-15	2.3	8
115	Relationships among personality traits, metabolic syndrome, and metabolic syndrome scores: The Kakegawa cohort study. <i>Journal of Psychosomatic Research</i> , 2018 , 107, 20-25	4.1	8
114	Home blood pressure level and decline in renal function among treated hypertensive patients: the J-HOME-Morning Study. <i>Hypertension Research</i> , 2016 , 39, 107-12	4.7	8
113	Correlation Between the Great East Japan Earthquake and Postpartum Depression: A Study in Miyako, Iwate, Japan. <i>Disaster Medicine and Public Health Preparedness</i> , 2015 , 9, 307-312	2.8	8
112	Randomized controlled trial of the effects of consumption of 'Yabukita' or 'Benifuuki' encapsulated tea-powder on low-density lipoprotein cholesterol level and body weight. <i>Food and Nutrition Research</i> , 2017 , 61, 1334484	3.1	8
111	A Brief Review of the 2009 JSSHP Guidelines for the care and treatment of Pregnancy induced Hypertension. <i>Hypertension Research in Pregnancy</i> , 2013 , 1, 5-7	0.7	8
110	Electrocardiographic abnormalities and home blood pressure in treated elderly hypertensive patients: Japan home versus office blood pressure measurement evaluation in the elderly (J-HOME-Elderly) study. <i>Hypertension Research</i> , 2010 , 33, 670-7	4.7	8
109	Pharmacists' awareness and attitude toward blood pressure measurement at home and in the pharmacy in Japan. <i>Clinical and Experimental Hypertension</i> , 2012 , 34, 447-55	2.2	8
108	Prevalence of Masked Hypertension in Subjects Treated with Antihypertensive Drugs as Assessed by Morning versus Evening Home Blood Pressure Measurements: the J-HOME study. <i>Clinical and Experimental Hypertension</i> , 2008 , 30, 277-87	2.2	8
107	Fermented Food Consumption and Psychological Distress in Pregnant Women: A Nationwide Birth Cohort Study of the Japan Environment and Children's Study. <i>Tohoku Journal of Experimental Medicine</i> , 2016 , 240, 309-321	2.4	8
106	Mother-to-infant bonding failure and intimate partner violence during pregnancy as risk factors for father-to-infant bonding failure at 1 month postpartum: an adjunct study of the Japan Environment and Children's Study. <i>Journal of Maternal-Fetal and Neonatal Medicine</i> , 2020 , 33, 2789-2796	2	8
105	Update on the prevalence and determinants of folic acid use in Japan evaluated with 91,538 pregnant women: the Japan Environment and Children's Study. <i>Journal of Maternal-Fetal and Neonatal Medicine</i> , 2020 , 33, 427-436	2	8

104	Association of maternal home blood pressure trajectory during pregnancy with infant birth weight: the BOSHI study. <i>Hypertension Research</i> , 2020 , 43, 550-559	4.7	7
103	Regional Birth Outcomes after the 2011 Great East Japan Earthquake and Tsunami in Miyagi Prefecture. <i>Prehospital and Disaster Medicine</i> , 2018 , 33, 215-219	0.8	7
102	Alcohol intake, circadian blood pressure variation, and stroke. <i>Hypertension</i> , 2009 , 53, 4-5	8.5	7
101	Prognostic significance of home arterial stiffness index derived from self-measurement of blood pressure: the Ohasama Study. <i>American Journal of Hypertension</i> , 2012 , 25, 67-73	2.3	7
100	Progress report on the HOMED-BP Study: hypertension objective treatment based on measurement by electrical devices of blood pressure study. <i>Clinical and Experimental Hypertension</i> , 2004 , 26, 119-27	2.2	7
99	The prevalence and risk factors for postpartum depression symptoms of fathers at one and 6 months postpartum: an adjunct study of the Japan Environment & Children's Study. <i>Journal of Maternal-Fetal and Neonatal Medicine</i> , 2020 , 33, 2797-2804	2	7
98	Reduced sleep efficiency, measured using an objective device, was related to an increased prevalence of home hypertension in Japanese adults. <i>Hypertension Research</i> , 2020 , 43, 23-29	4.7	7
97	Disease prevalence among nursery school children after the Great East Japan earthquake. <i>BMJ Global Health</i> , 2017 , 2, e000127	6.6	6
96	Maternity Log study: a longitudinal lifelog monitoring and multiomics analysis for the early prediction of complicated pregnancy. <i>BMJ Open</i> , 2019 , 9, e025939	3	6
95	N-Terminal Pro-B-Type Natriuretic Peptide Is Not a Significant Predictor of Stroke Incidence After 5 Years - The Ohasama Study. <i>Circulation Journal</i> , 2018 , 82, 2055-2062	2.9	6
94	Preconception folic acid supplementation use and the occurrence of neural tube defects in Japan: A nationwide birth cohort study of the Japan Environment and Children's Study. <i>Congenital Anomalies (discontinued)</i> , 2019 , 59, 110-117	1.1	6
93	Alterations in physique among young children after the Great East Japan Earthquake: Results from a nationwide survey. <i>Journal of Epidemiology</i> , 2017 , 27, 462-468	3.4	6
92	Monitoring and evaluation of out-of-office blood pressure during pregnancy. <i>Hypertension Research</i> , 2017 , 40, 107-109	4.7	6
91	Lower Incidence of Cervical Intraepithelial Neoplasia among Young Women with Human Papillomavirus Vaccination in Miyagi, Japan. <i>Tohoku Journal of Experimental Medicine</i> , 2017 , 243, 329-334	2.4	6
90	Mapping the genetic diversity of HLA haplotypes in the Japanese populations. <i>Scientific Reports</i> , 2015 , 5, 17855	4.9	6
89	Detection of midpregnancy fall in blood pressure by out-of-office monitoring. <i>Hypertension</i> , 2009 , 53, e12-3; author reply e14	8.5	6
88	How many measurements are needed to provide reliable information in terms of the ambulatory arterial stiffness index? The Ohasama study. <i>Hypertension Research</i> , 2011 , 34, 314-8	4.7	6
87	Maternal Baseline Characteristics and Perinatal Outcomes: the Tohoku Medical Megabank Project Birth and Three-Generation Cohort Study. <i>Journal of Epidemiology</i> , 2020 ,	3.4	6

86	Prescription trends in children with pervasive developmental disorders: a claims data-based study in Japan. <i>World Journal of Pediatrics</i> , 2016 , 12, 443-449	4.6	6
85	Antihypertensives prescribed for pregnant women in Japan: Prevalence and timing determined from a database of health insurance claims. <i>Pharmacoepidemiology and Drug Safety</i> , 2018 , 27, 1325-1334 ^{2,6}	2.6	6
84	Difference in the prevalence of gestational diabetes mellitus according to gestational age at 75-g oral glucose tolerance test in Japan: The Japan Assessment of Gestational Diabetes Mellitus Screening trial. <i>Journal of Diabetes Investigation</i> , 2019 , 10, 1576-1585	3.9	5
83	Validation of the Parama-Tech PS-501 device for office blood pressure measurement according to the international protocol. <i>Clinical and Experimental Hypertension</i> , 2012 , 34, 71-3	2.2	5
82	Evaluating home blood pressure in treated hypertensives in comparison with the referential value of casual screening of blood pressure: the Ohasama study. <i>Blood Pressure Monitoring</i> , 2012 , 17, 89-95	1.3	5
81	Parental longevity and offspring's home blood pressure: the Ohasama study. <i>Journal of Hypertension</i> , 2010 , 28, 272-7	1.9	5
80	Progress report on The Hypertension Objective Treatment Based on Measurement by Electrical Devices of Blood Pressure (HOMED-BP) study: status at February 2004. <i>Clinical and Experimental Hypertension</i> , 2007 , 29, 69-81	2.2	5
79	Comparison among research, home, and office blood pressure measurements for pregnant women: The TMM BirThree Cohort Study. <i>Journal of Clinical Hypertension</i> , 2020 , 22, 2004-2013	2.3	5
78	Blood Pressure and Chronic Kidney Disease Stratified by Gender and the Use of Antihypertensive Drugs. <i>Journal of the American Heart Association</i> , 2020 , 9, e015592	6	5
77	Blood pressure changes during twin pregnancies: the Japan Environment and Children's Study. <i>Journal of Hypertension</i> , 2019 , 37, 206-215	1.9	5
76	Stroke risk due to partial white-coat or masked hypertension based on the ACC/AHA guideline's blood pressure threshold: the Ohasama study. <i>Hypertension Research</i> , 2019 , 42, 120-122	4.7	5
75	Selective serotonin reuptake inhibitors and risk of major congenital anomalies for pregnancies in Japan: A nationwide birth cohort study of the Japan Environment and Children's Study. <i>Congenital Anomalies (discontinued)</i> , 2017 , 57, 72-78	1.1	4
74	Pregnant Women's Awareness of Social Capital in the Great East Japan Earthquake-Affected Areas of Miyagi Prefecture: The Japan Environment and Children's Study. <i>Disaster Medicine and Public Health Preparedness</i> , 2017 , 11, 355-364	2.8	4
73	Higher prevalence of hypertensive disorders of pregnancy in women who smoke: the Japan environment and children's study. <i>Hypertension Research</i> , 2019 , 42, 558-566	4.7	4
72	Randomized trial comparing the velocities of the antihypertensive effects on home blood pressure of candesartan and candesartan with hydrochlorothiazide. <i>Hypertension Research</i> , 2015 , 38, 701-7	4.7	4
71	Lacunar Infarcts Rather than White Matter Hyperintensity as a Predictor of Future Higher Level Functional Decline: The Ohasama Study. <i>Journal of Stroke and Cerebrovascular Diseases</i> , 2017 , 26, 376-384 ^{2,8}	2.8	4
70	Present status of clinical care for postpartum patients with hypertensive disorders of pregnancy in Japan: findings from a nationwide questionnaire survey. <i>Hypertension in Pregnancy</i> , 2015 , 34, 209-20	2	4
69	Associations between visit-to-visit variability in blood pressure measured in the office and antihypertensive drugs: the J-HOME-Morning study. <i>Clinical and Experimental Hypertension</i> , 2013 , 35, 285-90	2.2	4

68	Low-dose and very low-dose spironolactone in combination therapy for essential hypertension: evaluation by self-measurement of blood pressure at home. <i>Clinical and Experimental Hypertension</i> , 2011 , 33, 427-36	2.2	4
67	Time-dependent effects of imidapril administration in patients with morning hypertension measured as home blood pressure. <i>Clinical and Experimental Hypertension</i> , 2008 , 30, 243-54	2.2	4
66	Factors affecting heart rate as measured at home among treated hypertensive patients: the Japan home versus office blood pressure measurement evaluation (J-HOME) study. <i>Hypertension Research</i> , 2007 , 30, 1051-7	4.7	4
65	Incorporating self-blood pressure measurements at home in the guideline from the Ohasama study. <i>Blood Pressure Monitoring</i> , 2007 , 12, 407-9	1.3	4
64	Epidemiology of Hypertension Based on Ambulatory Blood Pressure Monitoring and Self-Measurement of Blood Pressure at Home. <i>Journal of Health Science</i> , 2004 , 50, 113-119		4
63	Preconception dysmenorrhea as a risk factor for psychological distress in pregnancy: The Japan Environment and Children's Study. <i>Journal of Affective Disorders</i> , 2019 , 245, 475-483	6.6	4
62	The present situation of home blood pressure measurement among outpatients in Japan. <i>Clinical and Experimental Hypertension</i> , 2020 , 42, 67-74	2.2	4
61	Multiple measurements of the urinary sodium-to-potassium ratio strongly related home hypertension: TMM Cohort Study. <i>Hypertension Research</i> , 2020 , 43, 62-71	4.7	4
60	Relation between disaster exposure, maternal characteristics, and obstetric outcomes: the Tohoku Medical Megabank Project Birth and Three-Generation Cohort Study. <i>Journal of Epidemiology</i> , 2021 ,	3.4	4
59	Strategic Methods for Recruiting Grandparents: The Tohoku Medical Megabank Birth and Three-Generation Cohort Study. <i>Tohoku Journal of Experimental Medicine</i> , 2018 , 246, 97-105	2.4	4
58	Predictive power of home blood pressure indices at baseline and during follow-up in hypertensive patients: HOMED-BP study. <i>Hypertension Research</i> , 2018 , 41, 622-628	4.7	4
57	Interannual Changes in the Prevalence of Intimate Partner Violence Against Pregnant Women in Miyagi Prefecture After the Great East Japan Earthquake: The Japan Environment and Children's Study. <i>Journal of Interpersonal Violence</i> , 2021 , 36, 10013-10028	2.2	3
56	Salt intake and the validity of a salt intake assessment system based on a 24-h dietary recall method in pregnant Japanese women. <i>Clinical and Experimental Hypertension</i> , 2015 , 37, 459-62	2.2	3
55	Differential effects of angiotensin II receptor blocker and losartan/hydrochlorothiazide combination on central blood pressure and augmentation index. <i>Clinical and Experimental Hypertension</i> , 2015 , 37, 294-302	2.2	3
54	Hypertensive disorders of pregnancy, obesity, and hypertension in later life by age group: a cross-sectional analysis. <i>Hypertension Research</i> , 2020 , 43, 1277-1283	4.7	3
53	Comparison of nocturnal blood pressure based on home versus ambulatory blood pressure measurement: The Ohasama Study. <i>Clinical and Experimental Hypertension</i> , 2020 , 42, 685-691	2.2	3
52	Age-Related Trends in Home Blood Pressure, Home Pulse Rate, and Day-to-Day Blood Pressure and Pulse Rate Variability Based on Longitudinal Cohort Data: The Ohasama Study. <i>Journal of the American Heart Association</i> , 2019 , 8, e012121	6	3
51	Maternal body mass index is a better indicator of large-for-gestational-age infants compared with a 75-g oral glucose tolerance test in early pregnancy: The JAGS trial. <i>Diabetes Research and Clinical Practice</i> , 2017 , 132, 10-18	7.4	3

50	Home blood pressure predicts stroke incidence among older adults with impaired physical function: the Ohasama study. <i>Journal of Hypertension</i> , 2017 , 35, 2395-2401	1.9	3
49	Differences between clinic and home blood pressure measurements during pregnancy. <i>Journal of Hypertension</i> , 2015 , 33, 1492-3	1.9	3
48	Blood pressure-lowering effect and duration of action of bedtime administration of doxazosin determined by home blood pressure measurement. <i>Clinical and Experimental Hypertension</i> , 2010 , 32, 311-7	2.2	3
47	Associated factors of home versus ambulatory heart rate variability in the general population: the Ohasama study. <i>Clinical and Experimental Hypertension</i> , 2011 , 33, 404-10	2.2	3
46	Control of home heart rate and home blood pressure levels in treated patients with hypertension: the J-HOME study. <i>Blood Pressure Monitoring</i> , 2007 , 12, 289-95	1.3	3
45	Efficacy of Combination Antihypertensive Therapy with Low-Dose Indapamide: Assessment by Blood Pressure Self-Monitoring at Home. <i>Clinical and Experimental Hypertension</i> , 2005 , 27, 331-341	2.2	3
44	Associations between glycosylated hemoglobin level at less than 24 weeks of gestation and adverse pregnancy outcomes in Japan: The Japan Environment and Children's Study (JECS). <i>Diabetes Research and Clinical Practice</i> , 2020 , 169, 108377	7.4	3
43	Knowledge, Attitude, and Practices Toward Blood Pressure Measurement at Home Among Japanese Nurses. <i>Home Healthcare Now</i> , 2016 , 34, 210-7	0.5	3
42	Public Attitudes toward an Epidemiological Study with Genomic Analysis in the Great East Japan Earthquake Disaster Area. <i>Prehospital and Disaster Medicine</i> , 2016 , 31, 330-4	0.8	2
41	Personality traits as predictors of decline in higher-level functional capacity over a 7-year follow-up in older adults: the Ohasama study. <i>Tohoku Journal of Experimental Medicine</i> , 2014 , 234, 197-207	2.4	2
40	Change of the management of treated hypertensive patients with or without diabetes in Japan. <i>Clinical and Experimental Hypertension</i> , 2013 , 35, 79-86	2.2	2
39	Out-of-office blood pressure control among treated subjects. <i>Hypertension</i> , 2007 , 49, e40-1; author reply e42	8.5	2
38	Definition of masked hypertension. <i>Journal of Hypertension</i> , 2007 , 25, 1511-2; author reply 1512-3	1.9	2
37	N-Terminal Pro-B-Type Natriuretic Peptide Is a Predictor of Chronic Kidney Disease in an Asian General Population - The Ohasama Study. <i>Circulation Reports</i> , 2019 , 2, 24-32	0.7	2
36	The Significance of Observing the Risk of Non-communicable Diseases after Large-scale Disasters and Communicable Disease Epidemics. <i>JMA Journal</i> , 2021 , 4, 305-310	1	2
35	Committee report on cerebrovascular disorders, including eclampsia and emergency medical services, of the Japan Society for the Study of Hypertension in Pregnancy. <i>Hypertension Research in Pregnancy</i> , 2013 , 1, 8-12	0.7	2
34	Low birth weight and abnormal pre-pregnancy body mass index were at higher risk for hypertensive disorders of pregnancy. <i>Pregnancy Hypertension</i> , 2020 , 22, 119-125	2.6	2
33	Epidemiological studies regarding hypertensive disorders of pregnancy: A review. <i>Journal of Obstetrics and Gynaecology Research</i> , 2020 , 46, 1672-1677	1.9	2

32	Lifetime risk of stroke stratified by chronic kidney disease and hypertension in the general Asian population: the Ohasama study. <i>Hypertension Research</i> , 2021 , 44, 866-873	4.7	2
31	The prevalence of psychological distress during pregnancy in Miyagi Prefecture for 3 years after the Great East Japan Earthquake. <i>Environmental Health and Preventive Medicine</i> , 2021 , 26, 27	4.2	2
30	Genome-wide association study for white coat effect in Japanese middle-aged to elderly people: The HOMED-BP study. <i>Clinical and Experimental Hypertension</i> , 2018 , 40, 363-369	2.2	2
29	Detailed association between serum uric acid levels and the incidence of chronic kidney disease stratified by sex in middle-aged adults. <i>Atherosclerosis</i> , 2021 , 330, 107-113	3.1	2
28	Correlation Between the Great East Japan Earthquake and Postpartum Depression: A Study in Miyako, Iwate, Japan. <i>Disaster Medicine and Public Health Preparedness</i> , 2015 , 9, 307-12	2.8	2
27	Accumulation of evidence regarding home blood pressure during pregnancy is necessary. <i>Hypertension Research</i> , 2017 , 40, 635-636	4.7	1
26	Home blood pressure measurement (HBPM) for the early detection of hypertensive disorders of pregnancy (HDP) in Japanese women: a multicenter prospective study. <i>Hypertension Research in Pregnancy</i> , 2017 , 5, 36-38	0.7	1
25	Individual assessment of inherent arterial stiffness using nomogram and pulse wave velocity index: the Ohasama study. <i>Clinical and Experimental Hypertension</i> , 2011 , 33, 147-52	2.2	1
24	Genetic polymorphisms in the beta-subunit of the epithelial sodium channel (β ENaC) gene in the Japanese population. <i>Clinical and Experimental Nephrology</i> , 2002 , 6, 130-4	2.5	1
23	Pregnancy outcomes in Japanese pregnant women with either chronic hypertension or white coat hypertension (JP-WCH): protocol for a prospective multicenter cohort study. <i>Hypertension Research in Pregnancy</i> , 2018 , 6, 7-10	0.7	1
22	Design and Progress of Oral Health Examinations in the Tohoku Medical Megabank Project. <i>Tohoku Journal of Experimental Medicine</i> , 2020 , 251, 97-115	2.4	1
21	Clustering by phenotype and genome-wide association study in autism		1
20	Prefabricated Temporary Housing and Eczema or Respiratory Symptoms in Schoolchildren after the Great East Japan Earthquake: The ToMMo Child Health Study. <i>Disaster Medicine and Public Health Preparedness</i> , 2019 , 13, 905-911	2.8	1
19	Machine learning approaches to predict gestational age in normal and complicated pregnancies via urinary metabolomics analysis. <i>Scientific Reports</i> , 2021 , 11, 17777	4.9	1
18	Spousal similarities in cardiometabolic risk factors: A cross-sectional comparison between Dutch and Japanese data from two large biobank studies. <i>Atherosclerosis</i> , 2021 , 334, 85-92	3.1	1
17	dbTMM: an integrated database of large-scale cohort, genome and clinical data for the Tohoku Medical Megabank Project. <i>Human Genome Variation</i> , 2021 , 8, 44	1.8	1
16	Maternal personality and postpartum mental disorders in Japan: the Tohoku Medical Megabank Project Birth and Three-Generation Cohort Study.. <i>Scientific Reports</i> , 2022 , 12, 6400	4.9	1
15	Evaluating folic acid supplementation among Japanese pregnant women with dietary intake of folic acid lower than 480 μ g per day: results from TMM Birth Three Cohort Study. <i>Journal of Maternal-Fetal and Neonatal Medicine</i> , 2020 , 1-6	2	0

14	Prenatal folic acid supplementation and autism spectrum disorder in 3-year-old offspring: the Japan environment and children's study. <i>Journal of Maternal-Fetal and Neonatal Medicine</i> , 2021 , 1-10	2	○
13	The A118G single-nucleotide polymorphism in OPRM1 is a risk factor for asthma severity. <i>Allergology International</i> , 2021 , 71, 55-55	4.4	○
12	Preoperative Risks of Cerebral Infarction in Pediatric Moyamoya Disease. <i>Stroke</i> , 2021 , 52, 2302-2310	6.7	○
11	Impacts of the urinary sodium-to-potassium ratio, sleep efficiency, and conventional risk factors on home hypertension in a general Japanese population. <i>Hypertension Research</i> , 2021 , 44, 858-865	4.7	○
10	Associations of education and income with hazardous drinking among postpartum women in Japan: results from the TMM BirThree Cohort Study. <i>Environmental Health and Preventive Medicine</i> , 2021 , 26, 70	4.2	○
9	The association of disproportionately enlarged subarachnoid space hydrocephalus with cognitive deficit in a general population: the Ohasama study. <i>Scientific Reports</i> , 2021 , 11, 17061	4.9	○
8	Maternal personality and alcohol use during pregnancy in Japan: The Tohoku Medical Megabank Project Birth and Three-Generation Cohort Study. <i>Addictive Behaviors</i> , 2021 , 122, 107020	4.2	○
7	Families' Health after the Great East Japan Earthquake: Findings from the Tohoku Medical Megabank Project Birth and Three-Generation Cohort Study.. <i>Tohoku Journal of Experimental Medicine</i> , 2022 , 256, 93-101	2.4	○
6	Cumulative exposure to maternal psychological distress in the prenatal and postnatal periods and atopic dermatitis in children: findings from the TMM BirThree Cohort Study.. <i>BMC Pregnancy and Childbirth</i> , 2022 , 22, 242	3.2	○
5	Ambulatory Blood Pressure Measurement and Home Blood Pressure Measurement. <i>Comprehensive Gynecology and Obstetrics</i> , 2018 , 199-208	○	
4	Hypotensive and heart rate-lowering effects of low-dose bisoprolol determined based on self-measured blood pressure at home. <i>Clinical and Experimental Hypertension</i> , 2012 , 34, 284-9	2.2	
3	Diurnal variability of hazards. <i>Journal of Hypertension</i> , 2007 , 25, 479-480	1.9	
2	Blood Pressure Phenotypes Defined by Ambulatory Blood Pressure Monitoring and Carotid Artery Changes in Community-Dwelling Older Japanese Adults: The Ohasama Study. <i>Tohoku Journal of Experimental Medicine</i> , 2020 , 252, 269-279	2.4	
1	Association of subjective health and abnormal cervical cytology in Japanese pregnant women: An adjunct study of the Japan Environment and Children's Study. <i>Preventive Medicine Reports</i> , 2021 , 24, 101525	2.6	