

# Yang Shin Park

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3314101/publications.pdf>

Version: 2024-02-01

26  
papers

385  
citations

759233

12  
h-index

794594

19  
g-index

26  
all docs

26  
docs citations

26  
times ranked

771  
citing authors

#	ARTICLE	IF	CITATIONS
1	Usefulness of Controlled Aliasing in Parallel Imaging Results in Higher Acceleration in Gadoteric Acid-Enhanced Liver Magnetic Resonance Imaging to Clarify the Hepatic Arterial Phase. <i>Investigative Radiology</i> , 2014, 49, 183-188.	6.2	45
2	Using intravoxel incoherent motion (IVIM) MR imaging to predict lipiodol uptake in patients with hepatocellular carcinoma following transcatheter arterial chemoembolization: A preliminary result. <i>Magnetic Resonance Imaging</i> , 2014, 32, 638-646.	1.8	39
3	Using Gd-EOB-DTPA-enhanced 3-T MRI for the differentiation of infiltrative hepatocellular carcinoma and focal confluent fibrosis in liver cirrhosis. <i>Magnetic Resonance Imaging</i> , 2013, 31, 1137-1142.	1.8	32
4	Use of Imaging to Predict Complete Response of Colorectal Liver Metastases after Chemotherapy: MR Imaging versus CT Imaging. <i>Radiology</i> , 2017, 284, 423-431.	7.3	31
5	Comparison of biannual ultrasonography and annual non-contrast liver magnetic resonance imaging as surveillance tools for hepatocellular carcinoma in patients with liver cirrhosis (MAGNUS-HCC): a study protocol. <i>BMC Cancer</i> , 2017, 17, 877.	2.6	30
6	Differentiation of hepatocellular carcinoma from its various mimickers in liver magnetic resonance imaging: What are the tips when using hepatocyte-specific agents?. <i>World Journal of Gastroenterology</i> , 2016, 22, 284.	3.3	28
7	Multiparametric MR Index for the Diagnosis of Non-Alcoholic Steatohepatitis in Patients with Non-Alcoholic Fatty Liver Disease. <i>Scientific Reports</i> , 2020, 10, 2671.	3.3	28
8	Hepatic Arterial Phase in Gadoteric Acid-Enhanced Liver Magnetic Resonance Imaging. <i>Investigative Radiology</i> , 2016, 51, 127-133.	6.2	21
9	Radiologic Evaluation and Structured Reporting Form for Extrahepatic Bile Duct Cancer: 2019 Consensus Recommendations from the Korean Society of Abdominal Radiology. <i>Korean Journal of Radiology</i> , 2021, 22, 41.	3.4	19
10	Application of High-Speed T1 Sequences for High-Quality Hepatic Arterial Phase Magnetic Resonance Imaging. <i>Investigative Radiology</i> , 2017, 52, 605-611.	6.2	18
11	Washout appearance in Gd-EOB-DTPA-enhanced MR imaging: A differentiating feature between hepatocellular carcinoma with paradoxical uptake on the hepatobiliary phase and focal nodular hyperplasia-like nodules. <i>Journal of Magnetic Resonance Imaging</i> , 2017, 45, 1599-1608.	3.4	18
12	Diagnosis and Surveillance of Incidental Pancreatic Cystic Lesions: 2017 Consensus Recommendations of the Korean Society of Abdominal Radiology. <i>Korean Journal of Radiology</i> , 2019, 20, 542.	3.4	18
13	Effect of Gd-EOB-DTPA on hepatic fat quantification using high-speed T2-corrected multi-echo acquisition in 1H MR spectroscopy. <i>Magnetic Resonance Imaging</i> , 2014, 32, 886-890.	1.8	10
14	Intravoxel incoherent motion magnetic resonance imaging to predict vesicoureteral reflux in children with urinary tract infection. <i>European Radiology</i> , 2016, 26, 1670-1677.	4.5	10
15	Multiparametric magnetic resonance imaging of endometrial polypoid lesions. <i>Abdominal Radiology</i> , 2020, 45, 3869-3881.	2.1	9
16	Use of hepatobiliary phase images in Gd-EOB-DTPA-enhanced MRI of breast cancer hepatic metastasis to predict response to chemotherapy. <i>Clinical Imaging</i> , 2017, 43, 127-131.	1.5	6
17	Second shot arterial phase to overcome degraded hepatic arterial phase in liver MR imaging. <i>European Radiology</i> , 2019, 29, 2821-2829.	4.5	5
18	Intraoperative Contrast-Enhanced Sonographic Portography Combined With Indigo Carmine Dye Injection for Anatomic Liver Resection in Hepatocellular Carcinoma. <i>Journal of Ultrasound in Medicine</i> , 2014, 33, 1287-1291.	1.7	4

#	ARTICLE	IF	CITATIONS
19	Abbreviated Gadoteric Acid-enhanced MRI with Second-Shot Arterial Phase Imaging for Liver Metastasis Evaluation. <i>Radiology Imaging Cancer</i> , 2019, 1, e190006.	1.6	4
20	The value of paradoxical uptake of hepatocellular carcinoma on the hepatobiliary phase of gadoteric acid-enhanced liver magnetic resonance imaging for the prediction of lipiodol uptake after transcatheter arterial chemoembolization. <i>European Journal of Radiology</i> , 2017, 89, 169-176.	2.6	3
21	Ultrasonographic index for the diagnosis of non-alcoholic steatohepatitis in patients with non-alcoholic fatty liver disease. <i>Quantitative Imaging in Medicine and Surgery</i> , 2022, 12, 1815-1829.	2.0	3
22	Ultrasonographic atlas of splenic lesions. <i>Ultrasonography</i> , 2021, , .	2.3	2
23	Fluoroscopy Guided Transurethral Placement of Ureteral Metallic Stents. <i>Iranian Journal of Radiology</i> , 2015, 12, e20269.	0.2	1
24	Ectopic lesions in the abdomen and pelvis: a multimodality pictorial review. <i>Abdominal Radiology</i> , 2022, 47, 2254-2276.	2.1	1
25	Re-use study of low-tube-voltage CT arterial phase in the spoiled gadoteric-acid liver MRI. <i>Clinical Imaging</i> , 2015, 39, 820-826.	1.5	0
26	Low-Tube Voltage Computed Tomography During Hepatic Arterial Phase. <i>Journal of Computer Assisted Tomography</i> , 2017, 41, 401-406.	0.9	0