Yang Shin Park

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/3314101/publications.pdf

Version: 2024-02-01

759233 794594 26 385 12 19 citations h-index g-index papers 26 26 26 771 docs citations times ranked citing authors all docs

#	Article	IF	CITATIONS
1	Ultrasonographic index for the diagnosis of non-alcoholic steatohepatitis in patients with non-alcoholic fatty liver disease. Quantitative Imaging in Medicine and Surgery, 2022, 12, 1815-1829.	2.0	3
2	Ectopic lesions in the abdomen and pelvis: a multimodality pictorial review. Abdominal Radiology, 2022, 47, 2254-2276.	2.1	1
3	Radiologic Evaluation and Structured Reporting Form for Extrahepatic Bile Duct Cancer: 2019 Consensus Recommendations from the Korean Society of Abdominal Radiology. Korean Journal of Radiology, 2021, 22, 41.	3.4	19
4	Ultrasonographic atlas of splenic lesions. Ultrasonography, 2021, , .	2.3	2
5	Multiparametric magnetic resonance imaging of endometrial polypoid lesions. Abdominal Radiology, 2020, 45, 3869-3881.	2.1	9
6	Multiparametric MR Index for the Diagnosis of Non-Alcoholic Steatohepatitis in Patients with Non-Alcoholic Fatty Liver Disease. Scientific Reports, 2020, 10, 2671.	3.3	28
7	Abbreviated Gadoxetic Acid–enhanced MRI with Second-Shot Arterial Phase Imaging for Liver Metastasis Evaluation. Radiology Imaging Cancer, 2019, 1, e190006.	1.6	4
8	Diagnosis and Surveillance of Incidental Pancreatic Cystic Lesions: 2017 Consensus Recommendations of the Korean Society of Abdominal Radiology. Korean Journal of Radiology, 2019, 20, 542.	3.4	18
9	Second shot arterial phase to overcome degraded hepatic arterial phase in liver MR imaging. European Radiology, 2019, 29, 2821-2829.	4.5	5
10	Low–Tube Voltage Computed Tomography During Hepatic Arterial Phase. Journal of Computer Assisted Tomography, 2017, 41, 401-406.	0.9	0
11	Use of Imaging to Predict Complete Response of Colorectal Liver Metastases after Chemotherapy: MR Imaging versus CT Imaging. Radiology, 2017, 284, 423-431.	7.3	31
12	The value of paradoxical uptake of hepatocellular carcinoma on the hepatobiliary phase of gadoxetic acid-enhanced liver magnetic resonance imaging for the prediction of lipiodol uptake after transcatheter arterial chemoembolization. European Journal of Radiology, 2017, 89, 169-176.	2.6	3
13	Use of hepatobiliary phase images in Gd-EOB-DTPA-enhanced MRI of breast cancer hepatic metastasis to predict response to chemotherapy. Clinical Imaging, 2017, 43, 127-131.	1.5	6
14	Application of High-Speed T1 Sequences for High-Quality Hepatic Arterial Phase Magnetic Resonance Imaging. Investigative Radiology, 2017, 52, 605-611.	6.2	18
15	Washout appearance in Gdâ€EOBâ€DTPAâ€enhanced MR imaging: A differentiating feature between hepatocellular carcinoma with paradoxical uptake on the hepatobiliary phase and focal nodular hyperplasiaâ€like nodules. Journal of Magnetic Resonance Imaging, 2017, 45, 1599-1608.	3.4	18
16	Comparison of biannual ultrasonography and annual non-contrast liver magnetic resonance imaging as surveillance tools for hepatocellular carcinoma in patients with liver cirrhosis (MAGNUS-HCC): a study protocol. BMC Cancer, 2017, 17, 877.	2.6	30
17	Differentiation of hepatocellular carcinoma from its various mimickers in liver magnetic resonance imaging: What are the tips when using hepatocyte-specific agents?. World Journal of Gastroenterology, 2016, 22, 284.	3.3	28
18	Hepatic Arterial Phase in Gadoxetic Acid–Enhanced Liver Magnetic Resonance Imaging. Investigative Radiology, 2016, 51, 127-133.	6.2	21

#	Article	IF	CITATION
19	Intravoxel incoherent motion magnetic resonance imaging to predict vesicoureteral reflux in children with urinary tract infection. European Radiology, 2016, 26, 1670-1677.	4.5	10
20	"Reuse―study of low-tube-voltage CT arterial phase in the spoiled gadoxetic-acid liver MRI. Clinical Imaging, 2015, 39, 820-826.	1.5	0
21	Fluoroscopy Guided Transurethral Placement of Ureteral Metallic Stents. Iranian Journal of Radiology, 2015, 12, e20269.	0.2	1
22	Usefulness of Controlled Aliasing in Parallel Imaging Results in Higher Acceleration in Gadoxetic Acid–Enhanced Liver Magnetic Resonance Imaging to Clarify the Hepatic Arterial Phase. Investigative Radiology, 2014, 49, 183-188.	6.2	45
23	Intraoperative Contrastâ€Enhanced Sonographic Portography Combined With Indigo Carmine Dye Injection for Anatomic Liver Resection in Hepatocellular Carcinoma. Journal of Ultrasound in Medicine, 2014, 33, 1287-1291.	1.7	4
24	Effect of Gd-EOB-DTPA on hepatic fat quantification using high-speed T2-corrected multi-echo acquisition in 1H MR spectroscopy. Magnetic Resonance Imaging, 2014, 32, 886-890.	1.8	10
25	Using intravoxel incoherent motion (IVIM) MR imaging to predict lipiodol uptake in patients with hepatocellular carcinoma following transcatheter arterial chemoembolization: A preliminary result. Magnetic Resonance Imaging, 2014, 32, 638-646.	1.8	39
26	Using Gd-EOB-DTPA-enhanced 3-T MRI for the differentiation of infiltrative hepatocellular carcinoma and focal confluent fibrosis in liver cirrhosis. Magnetic Resonance Imaging, 2013, 31, 1137-1142.	1.8	32