

Dimitrios E Kouzoukas

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3316253/publications.pdf>

Version: 2024-02-01

18
papers

222
citations

1162889

8
h-index

1058333

14
g-index

19
all docs

19
docs citations

19
times ranked

401
citing authors

#	ARTICLE	IF	CITATIONS
1	Precision Mapping of COVID-19 Vulnerable Locales by Epidemiological and Socioeconomic Risk Factors, Developed Using South Korean Data. <i>International Journal of Environmental Research and Public Health</i> , 2021, 18, 604.	1.2	14
2	Moderate blood alcohol and brain neurovulnerability: Selective depletion of calcium-independent PLA2, omega-3 docosahexaenoic acid and its synaptamide derivative as a potential harbinger of deficits in anti-inflammatory reserve. <i>Alcoholism: Clinical and Experimental Research</i> , 2021, .	1.4	1
3	PARP inhibition in vivo blocks alcohol-induced brain neurodegeneration and neuroinflammatory cytosolic phospholipase A2 elevations. <i>Neurochemistry International</i> , 2019, 129, 104497.	1.9	7
4	MIF mediates bladder pain, not inflammation, in cyclophosphamide cystitis. <i>Cytokine: X</i> , 2019, 1, 100003.	0.5	6
5	Subneurodegenerative Levels of Binge Alcohol Reduce Calcium-independent Phospholipase A2 (iPLA2) Levels in Rat Brain: An Unrecognized Initial Sign of Neuroinflammation. <i>FASEB Journal</i> , 2019, 33, 499.5.	0.2	0
6	Clonidine as a preoperative sedative. <i>Special Care in Dentistry</i> , 2018, 38, 80-88.	0.4	1
7	Elevated Urine Levels of Macrophage Migration Inhibitory Factor in Inflammatory Bladder Conditions: A Potential Biomarker for a Subgroup of Interstitial Cystitis/Bladder Pain Syndrome Patients. <i>Urology</i> , 2018, 116, 55-62.	0.5	26
8	Activation of cyclic GMP-dependent protein kinase blocks alcohol-mediated cell death and calcium disruption in cerebellar granule neurons. <i>Neuroscience Letters</i> , 2018, 676, 108-112.	1.0	4
9	Preclinical study of a Kv11.1 potassium channel activator as antineoplastic approach for breast cancer. <i>Oncotarget</i> , 2018, 9, 3321-3337.	0.8	41
10	Disulfide high mobility group box-1 causes bladder pain through bladder Toll-like receptor 4. <i>BMC Physiology</i> , 2017, 17, 6.	3.6	22
11	Macrophage migration inhibitory factor mediates protease-activated receptor 4-induced bladder pain through urothelial high mobility group box 1. <i>Physiological Reports</i> , 2017, 5, e13549.	0.7	12
12	MP28-14 PROTEASE-ACTIVATED RECEPTOR 4 INDUCES BLADDER PAIN THROUGH DISULFIDE HIGH MOBILITY GROUP BOX-1 ACTING ON RECEPTORS FOR ADVANCED GLYCATION ENDPRODUCTS. <i>Journal of Urology</i> , 2016, 195, .	0.2	0
13	MP72-01 URINE LEVELS OF MACROPHAGE MIGRATION INHIBITORY FACTOR ARE INCREASED IN PATIENTS WITH BLADDER INFLAMMATION. <i>Journal of Urology</i> , 2016, 195, .	0.2	0
14	Protease-Activated Receptor 4 Induces Bladder Pain through High Mobility Group Box-1. <i>PLoS ONE</i> , 2016, 11, e0152055.	1.1	23
15	MP21-17 MACROPHAGE MIGRATION INHIBITORY FACTOR MEDIATES PROTEASE ACTIVATED RECEPTOR-INDUCED BLADDER PAIN. <i>Journal of Urology</i> , 2015, 193, .	0.2	0
16	Macrophage Migration Inhibitory Factor Mediates PAR-Induced Bladder Pain. <i>PLoS ONE</i> , 2015, 10, e0127628.	1.1	24
17	Intracellular calcium plays a critical role in the alcohol-mediated death of cerebellar granule neurons. <i>Journal of Neurochemistry</i> , 2013, 124, 323-335.	2.1	34
18	Neuropsychological Correlates of Normal Variation in Emotional Response to Visual Stimuli. <i>Journal of Nervous and Mental Disease</i> , 2007, 195, 112-118.	0.5	6