Anna Bednarek-Skublewska

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/3340437/publications.pdf

Version: 2024-02-01

36 738 13 27 g-index

37 37 37 37 1094

times ranked

citing authors

docs citations

all docs

#	Article	IF	Citations
1	Prevalence and Correlates of Self-Reported Sexual Dysfunction in CKD: A Meta-analysis of Observational Studies. American Journal of Kidney Diseases, 2010, 56, 670-685.	2.1	155
2	Fruit and Vegetable Intake and Mortality in Adults undergoing Maintenance Hemodialysis. Clinical Journal of the American Society of Nephrology: CJASN, 2019, 14, 250-260.	2.2	85
3	Sexual Dysfunction in Women with ESRD Requiring Hemodialysis. Clinical Journal of the American Society of Nephrology: CJASN, 2012, 7, 974-981.	2.2	82
4	Dental Health and Mortality in People With End-Stage Kidney Disease Treated With Hemodialysis: A Multinational Cohort Study. American Journal of Kidney Diseases, 2015, 66, 666-676.	2.1	55
5	Prevalence and correlates of erectile dysfunction in men on chronic haemodialysis: a multinational cross-sectional study. Nephrology Dialysis Transplantation, 2012, 27, 2479-2488.	0.4	44
6	The Association of Mediterranean and DASH Diets with Mortality in Adults on Hemodialysis: The DIET-HD Multinational Cohort Study. Journal of the American Society of Nephrology: JASN, 2018, 29, 1741-1751.	3.0	33
7	Patient satisfaction with in-centre haemodialysis care: an international survey. BMJ Open, 2014, 4, e005020.	0.8	29
8	The prevalence and correlates of low sexual functioning in women on hemodialysis: A multinational, cross-sectional study. PLoS ONE, 2017, 12, e0179511.	1.1	26
9	Nutrition and dietary intake and their association with mortality and hospitalisation in adults with chronic kidney disease treated with haemodialysis: protocol for DIET-HD, a prospective multinational cohort study. BMJ Open, 2015, 5, e006897-e006897.	0.8	24
10	Monocyte chemoattractant protein-1 (MCP-1) gene polymorphism as a potential risk factor for cardiovascular disease in hemodialyzed patients. Cytokine, 2008, 44, 361-365.	1.4	20
11	Oral disease in adults treated with hemodialysis: prevalence, predictors, and association with mortality and adverse cardiovascular events: the rationale and design of the ORAL Diseases in hemodialysis (ORAL-D) study, a prospective, multinational, longitudinal, observational, cohort study. BMC Nephrology, 2013, 14, 90.	0.8	19
12	Depression and all-cause and cardiovascular mortality in patients on haemodialysis: a multinational cohort study. Nephrology Dialysis Transplantation, 2016, 32, gfw016.	0.4	17
13	Dietary n-3 polyunsaturated fatty acid intake and all-cause and cardiovascular mortality in adults on hemodialysis: The DIET-HD multinational cohort study. Clinical Nutrition, 2019, 38, 429-437.	2.3	16
14	The prognostic value of positive T-wave in lead aVR in hemodialysis patients. Clinical and Experimental Nephrology, 2015, 19, 1157-1164.	0.7	14
15	Patterns of oral disease in adults with chronic kidney disease treated with hemodialysis. Nephrology Dialysis Transplantation, 2016, 31, 1647-1653.	0.4	14
16	Periodontitis and early mortality among adults treated with hemodialysis: a multinational propensity-matched cohort study. BMC Nephrology, 2017, 18, 166.	0.8	14
17	Dietary Patterns and Mortality in a Multinational Cohort of Adults Receiving Hemodialysis. American Journal of Kidney Diseases, 2020, 75, 361-372.	2.1	12
18	The effect of a single dialysis session on spatial QRS-T angle in haemodialysis patients. Nephrology Dialysis Transplantation, 2010, 25, 3723-3729.	0.4	11

#	Article	IF	Citations
19	Effects of vitamin D3 on selected biochemical parameters of nutritional status, inflammation, and cardiovascular disease in patients undergoing long-term hemodialysis., 2010, 120, 167-74.		10
20	Complement receptor 1 gene polymorphism and cardiovascular disease in dialyzed end-stage renal disease patients. Human Immunology, 2010, 71, 878-882.	1.2	9
21	lodine-123 metaiodobenzylguanidine myocardial imaging in haemodialysed patients asymptomatic for coronary artery disease. Nuclear Medicine Communications, 2011, 32, 515-521.	0.5	9
22	Relationship between renalase and N-terminal pro-B-type natriuretic peptide (NT pro-BNP) in haemodialysis patients. Annals of Agricultural and Environmental Medicine, 2014, 21, 132-5.	0.5	8
23	Relationship between microRNA-146a expression and plasma renalase levels in hemodialyzed patients. PLoS ONE, 2017, 12, e0179218.	1.1	7
24	Plasma and erythrocyte relationship of catecholamines in hemodialysis patients. Annals of Agricultural and Environmental Medicine, 2014, 21, 562-566.	0.5	6
25	Relationship of haemodialysis therapy duration and cardiac adrenergic system function assessed by iodine-123 metaiodobenzylguanidine imaging in haemodialysed nondiabetic patients. Nuclear Medicine Communications, 2012, 33, 155-163.	0.5	5
26	Levels of renalase and advanced oxidation protein products with regard to catecholamines in haemodialysed patients. Annals of Agricultural and Environmental Medicine, 2017, 24, 453-458.	0.5	3
27	Prevalence of Legionella antibodies in immunocompromised patients. Open Medicine (Poland), 2013, 8, 208-212.	0.6	2
28	Assessment of Cytokine Release Afterin VitroStimulation of Whole Blood withLegionella Pneumophilain Immunocompromised Patients. Immunological Investigations, 2013, 42, 1-17.	1.0	2
29	Oral mucosal lesions and risk of all-cause and cardiovascular mortality in people treated with long-term haemodialysis: The ORAL-D multinational cohort study. PLoS ONE, 2019, 14, e0218684.	1.1	2
30	Plasma levels of catecholamines and asymmetric dimethylarginine levels as predictive values of mortality among hemodialysis patients. Current Issues in Pharmacy and Medical Sciences, 2014, 27, 37-40.	0.1	2
31	Influence of Homocysteine on the Function of Native Arteriovenous Fistula in Hemodialysis Patients. Dialysis and Transplantation, 2006, 35, 604-609.	0.2	1
32	FP630ASSOCIATION OF FRUIT AND VEGETABLE INTAKE WITH ALL-CAUSE MORTALITY IN HEMODIALYSIS PATIENTS (DIET-HD): A PROSPECTIVE COHORT STUDY. Nephrology Dialysis Transplantation, 2018, 33, i255-i25'6.	0.4	1
33	Concentrations of branched amino acids: isoleucine, valine, leucine in serum of hemodialysis patients during one-year observation. Annales Universitatis Mariae Curie-Sklodowska Sectio D: Medicina, 2002, 57, 322-9.	0.0	1
34	FP631PREVALENCE OF ORAL MUCOSAL LESIONS IN HEMODIALYSIS PATIENTS AND ASSOCIATION WITH MORTALITY: A PROSPECTIVE COHORT STUDY. Nephrology Dialysis Transplantation, 2018, 33, i256-i256.	0.4	0
35	Retrospective single-center analysis of the reasons and serious complications of peritoneal-related peritonitis in patients with end-stage renal disease undergoing peritoneal dialysis treatment. Zdrowie Publiczne, 2018, 128, 103-106.	0.2	0
36	Sialic acid content in erythrocyte membranes of patients on chronic hemodialysis., 2009, 119, 194-9.		0