

Marit-Solveig Seidenkrantz

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/3340623/marit-solveig-seidenkrantz-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

121
papers

3,776
citations

36
h-index

57
g-index

152
ext. papers

4,270
ext. citations

3.9
avg, IF

5.38
L-index

#	Paper	IF	Citations
121	Holocene palaeoceanography of the Northeast Greenland shelf. <i>Climate of the Past</i> , 2022 , 18, 103-127	3.9	2
120	Characterization of organic matter in marine sediments to estimate age offset of bulk radiocarbon dating. <i>Quaternary Geochronology</i> , 2022 , 67, 101242	2.7	0
119	Linkages between ocean circulation and the Northeast Greenland Ice Stream in the Early Holocene. <i>Quaternary Science Reviews</i> , 2022 , 286, 107530	3.9	0
118	Seasonal climate variations in the Baltic Sea during the Last Interglacial based on foraminiferal geochemistry. <i>Quaternary Science Reviews</i> , 2021 , 272, 107220	3.9	0
117	Evidence for influx of Atlantic water masses to the Labrador Sea during the Last Glacial Maximum. <i>Scientific Reports</i> , 2021 , 11, 6788	4.9	0
116	Comparative analysis of six common foraminiferal species of the genera <i>Cassidulina</i> , <i>Paracassidulina</i> , and <i>Islandiella</i> from the Arctic-North Atlantic domain. <i>Journal of Micropalaeontology</i> , 2021 , 40, 37-60	2	11
115	When were the straits between the Baltic Sea and the Kattegat inundated by the sea during the Holocene?. <i>Boreas</i> , 2021 , 50, 1079	2.4	2
114	Mid to late-Holocene sea-surface temperature variability off north-eastern Newfoundland and its linkage to the North Atlantic Oscillation. <i>Holocene</i> , 2021 , 31, 3-15	2.6	5
113	Spatio-temporal changes in ocean conditions and primary production in Baffin Bay and the Labrador Sea. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2021 , 563, 110175	2.9	2
112	Insolation vs. meltwater control of productivity and sea surface conditions off SW Greenland during the Holocene. <i>Boreas</i> , 2021 , 50, 631-651	2.4	1
111	Episodic Atlantic Water Inflow Into the Independence Fjord System (Eastern North Greenland) During the Holocene and Last Glacial Period. <i>Frontiers in Earth Science</i> , 2020 , 8,	3.5	1
110	Macrofaunal control of microbial community structure in continental margin sediments. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2020 , 117, 15911-15922	11.5	11
109	Holocene Hydrographic Variations From the Baltic-North Sea Transitional Area (IODP Site M0059). <i>Paleoceanography and Paleoclimatology</i> , 2020 , 35, e2019PA003722	3.3	6
108	Reconstruction of Holocene oceanographic conditions in eastern Baffin Bay. <i>Climate of the Past</i> , 2020 , 16, 1075-1095	3.9	11
107	Distribution of dinocyst assemblages in surface sediment samples from the West Greenland margin. <i>Marine Micropaleontology</i> , 2020 , 159, 101818	1.7	3
106	Early diagenesis of foraminiferal calcite under anoxic conditions: A case study from the Landsort Deep, Baltic Sea (IODP Site M0063). <i>Chemical Geology</i> , 2020 , 558, 119871	4.2	1
105	Glomulina oculus, New Calcareous Foraminiferal Species from the High Arctic: A Potential Indicator of a Nearby Marine-Terminating Glacier. <i>Journal of Foraminiferal Research</i> , 2020 , 50, 219-234	1.1	3

104	Learning from the past: Impact of the Arctic Oscillation on sea ice and marine productivity off northwest Greenland over the last 9,000 years. <i>Global Change Biology</i> , 2020 , 26, 6767-6786	11.4	9
103	Middle to Late Holocene Variations in Salinity and Primary Productivity in the Central Baltic Sea: A Multiproxy Study From the Landsort Deep. <i>Frontiers in Marine Science</i> , 2019 , 6,	4.5	7
102	Glacial history and palaeo-environmental change of southern Taimyr Peninsula, Arctic Russia, during the Middle and Late Pleistocene. <i>Earth-Science Reviews</i> , 2019 , 196, 102832	10.2	5
101	Holocene water mass changes in the Labrador Current. <i>Holocene</i> , 2019 , 29, 676-690	2.6	11
100	Data set on sedimentology, palaeoecology and chronology of middle to late pleistocene deposits on the Taimyr Peninsula, Arctic Russia. <i>Data in Brief</i> , 2019 , 25, 104267	1.2	4
99	A reconstruction of warm-water inflow to Upernavik Isstrøm since 1925 CE and its relation to glacier retreat. <i>Climate of the Past</i> , 2019 , 15, 1171-1186	3.9	7
98	Southwest Greenland shelf glaciation during MIS 4 more extensive than during the Last Glacial Maximum. <i>Scientific Reports</i> , 2019 , 9, 15617	4.9	9
97	Development of the western Limfjord, Denmark, after the last deglaciation: a review with new data. <i>Bulletin of the Geological Society of Denmark</i> , 2019 , 67, 53-73	1	4
96	Glacial Runoff Promotes Deep Burial of Sulfur Cycling-Associated Microorganisms in Marine Sediments. <i>Frontiers in Microbiology</i> , 2019 , 10, 2558	5.7	10
95	Late Holocene Sea Surface Instabilities in the Disko Bugt Area, West Greenland, in Phase With 180 Oscillations at Camp Century. <i>Paleoceanography and Paleoclimatology</i> , 2018 , 33, 227-243	3.3	15
94	East Asian Winter Monsoon Variations and Their Links to Arctic Sea Ice During the Last Millennium, Inferred From Sea Surface Temperatures in the Okinawa Trough. <i>Paleoceanography and Paleoclimatology</i> , 2018 , 33, 61-75	3.3	5
93	Linking the Modern Distribution of Biogenic Proxies in High Arctic Greenland Shelf Sediments to Sea Ice, Primary Production, and Arctic-Atlantic Inflow. <i>Journal of Geophysical Research G: Biogeosciences</i> , 2018 , 123, 760-786	3.7	23
92	Subpolar North Atlantic sea surface temperature since 6 ka BP: Indications of anomalous ocean-atmosphere interactions at 4-2 ka BP. <i>Quaternary Science Reviews</i> , 2018 , 194, 128-142	3.9	18
91	Meltwater and seasonality influence on Subpolar Gyre circulation during the Holocene. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2018 , 502, 104-118	2.9	11
90	Benthic foraminiferal assemblages and test accumulation in coastal microhabitats on San Salvador, Bahamas. <i>Journal of Micropalaeontology</i> , 2018 , 37, 499-518	2	0
89	Relative Sea-Level Changes and Ice Sheet History in Funderup Land, North Greenland. <i>Frontiers in Earth Science</i> , 2018 , 6,	3.5	11
88	The Holocene Great Belt connection to the southern Kattegat, Scandinavia: Ancylus Lake drainage and Early Littorina Sea transgression. <i>Boreas</i> , 2017 , 46, 53-68	2.4	22
87	Upper water mass variability in the Anegada-Jungfern Passage, NE Caribbean, during the last 11,100 cal. yr. <i>Holocene</i> , 2017 , 27, 1291-1307	2.6	1

86	A record of Holocene sea-ice variability off West Greenland and its potential forcing factors. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2017 , 475, 115-124	2.9	19
85	Minimal Holocene retreat of large tidewater glaciers in Kĕge Bugt, southeast Greenland. <i>Scientific Reports</i> , 2017 , 7, 12330	4.9	8
84	Atlantic Water advection vs. glacier dynamics in northern Spitsbergen since early deglaciation. <i>Climate of the Past</i> , 2017 , 13, 1717-1749	3.9	22
83	Reconstructing Holocene temperature and salinity variations in the western Baltic Sea region: a multi-proxy comparison from the Little Belt (IODP Expedition B47, Site M0059). <i>Biogeosciences</i> , 2017 , 14, 5607-5632	4.6	16
82	Microbial turnover times in the deep seabed studied by amino acid racemization modelling. <i>Scientific Reports</i> , 2017 , 7, 5680	4.9	37
81	Sea ice and primary production proxies in surface sediments from a High Arctic Greenland fjord: Spatial distribution and implications for palaeoenvironmental studies. <i>Ambio</i> , 2017 , 46, 106-118	6.5	27
80	Early onset of industrial-era warming across the oceans and continents. <i>Nature</i> , 2016 , 536, 411-8	50.4	167
79	Water balance in the complex mountainous terrain of Bhutan and linkages to land use. <i>Journal of Hydrology: Regional Studies</i> , 2016 , 7, 55-68	3.6	11
78	The role of sea ice for vascular plant dispersal in the Arctic. <i>Biology Letters</i> , 2016 , 12,	3.6	14
77	Ice stream retreat following the LGM and onset of the west Greenland current in Uummannaq Trough, west Greenland. <i>Quaternary Science Reviews</i> , 2016 , 147, 27-46	3.9	36
76	Solar forcing as an important trigger for West Greenland sea-ice variability over the last millennium. <i>Quaternary Science Reviews</i> , 2016 , 131, 148-156	3.9	26
75	Holocene oceanographic changes in SW Labrador Sea, off Newfoundland. <i>Holocene</i> , 2016 , 26, 274-289	2.6	14
74	Spatial Variation of Temperature and Precipitation in Bhutan and Links to Vegetation and Land Cover. <i>Mountain Research and Development</i> , 2016 , 36, 66	1.4	21
73	Solar forcing of Holocene summer sea-surface temperatures in the northern North Atlantic. <i>Geology</i> , 2015 , 43, 203-206	5	67
72	Robust global ocean cooling trend for the pre-industrial Common Era. <i>Nature Geoscience</i> , 2015 , 8, 671-678	7.3	134
71	Heinrich 0 on the east Canadian margin: Source, distribution, and timing. <i>Paleoceanography</i> , 2015 , 30, 1613-1624		11
70	Sequence of events from the onset to the demise of the Last Interglacial: Evaluating strengths and limitations of chronologies used in climatic archives. <i>Quaternary Science Reviews</i> , 2015 , 129, 1-36	3.9	104
69	Formate, acetate, and propionate as substrates for sulfate reduction in sub-arctic sediments of Southwest Greenland. <i>Frontiers in Microbiology</i> , 2015 , 6, 846	5.7	49

68	Variable influx of West Greenland Current water into the Labrador Current through the last 7200 years: a multiproxy record from Trinity Bay (NE Newfoundland). <i>Arktos</i> , 2015 , 1, 1	0.9	7
67	Evidence for external forcing of the Atlantic Multidecadal Oscillation since termination of the Little Ice Age. <i>Nature Communications</i> , 2014 , 5, 3323	17.4	85
66	The Holocene marine diatom flora of Eastern Newfoundland bays. <i>Diatom Research</i> , 2014 , 29, 441-454	0.9	10
65	A diatom-based sea-ice reconstruction for the Vaigat Strait (Disko Bugt, West Greenland) over the last 5000yr. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2014 , 403, 66-79	2.9	33
64	A 100-year record of changes in water renewal rate in Sermilik fjord and its influence on calving of Helheim glacier, southeast Greenland. <i>Continental Shelf Research</i> , 2014 , 85, 21-29	2.4	8
63	Labrador current variability over the last 2000 years. <i>Earth and Planetary Science Letters</i> , 2014 , 400, 26-33	3.3	42
62	Impact of Medieval Fjord Hydrography and Climate on the Western and Eastern Settlements in Norse Greenland. <i>Journal of the North Atlantic</i> , 2014 , 601, 1-13	0.4	5
61	A multi-proxy reconstruction of oceanographic conditions around the Younger Dryas-Holocene transition in Placentia Bay, Newfoundland. <i>Marine Micropaleontology</i> , 2014 , 112, 39-49	1.7	18
60	Diatom-based reconstruction of summer sea-surface salinity in the South China Sea over the last 15 000 years. <i>Boreas</i> , 2014 , 43, 208-219	2.4	6
59	Atmospheric and oceanic influence on mass balance of northern North Atlantic region land-terminating glaciers. <i>Geografiska Annaler, Series A: Physical Geography</i> , 2014 , 96, n/a-n/a	1.1	5
58	Early Holocene large-scale meltwater discharge from Greenland documented by foraminifera and sediment parameters. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2013 , 391, 71-81	2.9	31
57	Ocean lead at the termination of the Younger Dryas cold spell. <i>Nature Communications</i> , 2013 , 4, 1664	17.4	51
56	Sea ice in the paleoclimate system: the challenge of reconstructing sea ice from proxies and an introduction. <i>Quaternary Science Reviews</i> , 2013 , 79, 1-8	3.9	67
55	Evaluation of the sea ice proxy IP25 against observational and diatom proxy data in the SW Labrador Sea. <i>Quaternary Science Reviews</i> , 2013 , 79, 53-62	3.9	36
54	Benthic foraminifera as palaeo sea-ice indicators in the subarctic realm: examples from the Labrador Sea-Baffin Bay region. <i>Quaternary Science Reviews</i> , 2013 , 79, 135-144	3.9	61
53	Mid- to late-Holocene oceanographic variability on the Southeast Greenland shelf. <i>Holocene</i> , 2013 , 23, 167-178	2.6	32
52	Environmental reconstructions of Eemian Stage interglacial marine records in the Lower Vistula area, southern Baltic Sea. <i>Boreas</i> , 2012 , 41, 209-234	2.4	9
51	A diatom-based reconstruction of summer sea-surface salinity in the Southern Okinawa Trough, East China Sea, over the last millennium. <i>Journal of Quaternary Science</i> , 2012 , 27, 771-779	2.3	7

50	Greenland climate change: from the past to the future. <i>Wiley Interdisciplinary Reviews: Climate Change</i> , 2012 , 3, 427-449	8.4	22
49	Climate-driven fluctuations in freshwater flux to Sermilik Fjord, East Greenland, during the last 4000 years. <i>Holocene</i> , 2012 , 22, 155-164	2.6	15
48	Evidence of Suess solar-cycle bursts in subtropical Holocene speleothem $\delta^{18}O$ records. <i>Holocene</i> , 2012 , 22, 597-602	2.6	17
47	Exotic pollen as an indicator of variable atmospheric circulation over the Labrador Sea region during the mid to late Holocene. <i>Journal of Quaternary Science</i> , 2011 , 26, 286-296	2.3	34
46	Tracking the Atlantic Multidecadal Oscillation through the last 8,000 years. <i>Nature Communications</i> , 2011 , 2, 178	17.4	238
45	Lacustrine evidence of Holocene environmental change from three Faroese lakes: a multiproxy XRF and stable isotope study. <i>Quaternary Science Reviews</i> , 2010 , 29, 2764-2780	3.9	27
44	A diatom-based reconstruction of Early Holocene hydrographic and climatic change in a southwest Greenland fjord. <i>Marine Micropaleontology</i> , 2009 , 70, 166-176	1.7	40
43	Taking the pulse of the Sun during the Holocene by joint analysis of ^{14}C and ^{10}Be . <i>Geophysical Research Letters</i> , 2009 , 36,	4.9	51
42	The 7 Aarhus Statements on Climate Change. <i>IOP Conference Series: Earth and Environmental Science</i> , 2009 , 8, 011002	0.3	
41	Comparing past and present climate $\delta^{18}O$ tool to distinguish between natural and human-induced climate change. <i>IOP Conference Series: Earth and Environmental Science</i> , 2009 , 8, 012012	0.3	3
40	Termination of the Medieval Warm Period: Linking sub-polar and tropical N Atlantic circulation changes to ENSO. <i>PAGES News</i> , 2009 , 17, 76-77		8
39	Glacial and palaeoenvironmental history of the Cape Chelyuskin area, Arctic Russia. <i>Polar Research</i> , 2008 , 27, 222-248	2	18
38	Deglacial and Holocene conditions in northernmost Baffin Bay: sediments, foraminifera, diatoms and stable isotopes. <i>Boreas</i> , 2008 , 37, 346-376	2.4	87
37	Variable North Atlantic climate seesaw patterns documented by a late Holocene marine record from Disko Bugt, West Greenland. <i>Marine Micropaleontology</i> , 2008 , 68, 66-83	1.7	77
36	Foraminifera from the Quaternary sequence in the Anholt boring, Denmark. <i>Boreas</i> , 2008 , 22, 283-290	2.4	20
35	Quaternary sequences and their relations to the pre-Quaternary in the vicinity of Anholt, Kattegat, Scandinavia. <i>Boreas</i> , 2008 , 22, 291-298	2.4	19
34	Middle Weichselian to Holocene palaeoecology in the eastern Kattegat, Scandinavia: foraminifera, ostracods and ^{14}C measurements. <i>Boreas</i> , 2008 , 22, 299-310	2.4	36
33	Subrecent changes in the foraminiferal distribution in the Kattegat and the Skagerrak, Scandinavia: anthropogenic influence and natural causes. <i>Boreas</i> , 2008 , 22, 383-395	2.4	23

32	Palaeoenvironments in the Skagerrak-Kattegat basin in the eastern North Sea during the last deglaciation. <i>Boreas</i> , 2008 , 25, 65-78	2.4	32
31	A late Holocene palaeoenvironmental record from Altona Bay, St. Croix, US Virgin Islands. <i>Geografisk Tidsskrift</i> , 2008 , 108, 59-70	1.5	6
30	Hydrography and climate of the last 4400 years in a SW Greenland fjord: implications for Labrador Sea palaeoceanography. <i>Holocene</i> , 2007 , 17, 387-401	2.6	97
29	Late-Holocene environment and climatic changes in Ameralik Fjord, southwest Greenland: evidence from the sedimentary record. <i>Holocene</i> , 2006 , 16, 685-695	2.6	45
28	Severnaya Zemlya, Arctic Russia: a nucleation area for Kara Sea ice sheets during the Middle to Late Quaternary. <i>Quaternary Science Reviews</i> , 2006 , 25, 2894-2936	3.9	30
27	Last Interglacial marine environments in the White Sea region, northwestern Russia. <i>Boreas</i> , 2006 , 35, 493-520	2.4	37
26	Last Interglacial (Eemian) hydrographic conditions in the southeastern Baltic Sea, NE Europe, based on dinoflagellate cysts. <i>Quaternary International</i> , 2005 , 130, 3-30	2	48
25	Evidence for solar forcing of sea-surface temperature on the North Icelandic Shelf during the late Holocene. <i>Geology</i> , 2005 , 33, 73	5	130
24	Diatoms from the surface sediments of the South China Sea and their relationships to modern hydrography. <i>Marine Micropaleontology</i> , 2004 , 53, 279-292	1.7	42
23	Environmental changes off North Iceland during the deglaciation and the Holocene: foraminifera, diatoms and stable isotopes. <i>Marine Micropaleontology</i> , 2004 , 50, 273-305	1.7	71
22	New foraminiferal taxa and revised biostratigraphy of Jurassic marginal marine deposits on Anholt, Denmark. <i>Micropaleontology</i> , 2003 , 49, 27	2	6
21	Last Interglacial and Early Glacial Circulation in the Northern North Atlantic Ocean. <i>Quaternary Research</i> , 2002 , 58, 22-26	1.9	25
20	Late-Holocene summer sea-surface temperatures based on a diatom record from the north Icelandic shelf. <i>Holocene</i> , 2002 , 12, 137-147	2.6	3
19	Late-Holocene summer sea-surface temperatures based on a diatom record from the north Icelandic shelf. <i>Holocene</i> , 2002 , 12, 137-147	2.6	85
18	MODERN DISTRIBUTION OF BENTHIC FORAMINIFERA ON THE NORTH ICELANDIC SHELF AND SLOPE. <i>Journal of Foraminiferal Research</i> , 2002 , 32, 217-244	1.1	96
17	Diatom surface sediment assemblages around Iceland and their relationships to oceanic environmental variables. <i>Marine Micropaleontology</i> , 2001 , 41, 73-96	1.7	88
16	Marine late Saalian to Eemian environments and climatic variability in the Danish shelf area. <i>Geologie En Mijnbouw/Netherlands Journal of Geosciences</i> , 2000 , 79, 335-343	1.1	25
15	Benthic foraminifera as indicators of changing Mediterranean-Atlantic water exchange in the late Miocene. <i>Marine Geology</i> , 2000 , 163, 387-407	3.3	80

14	Santorini (Greece) before the Minoan eruption: a reconstruction of the ring-island, natural resources and clay deposits from the Akrotiri excavation. <i>Geological Society Special Publication</i> , 2000 , 171, 71-80	1.7	3
13	Deep-water changes: the near-synchronous disappearance of a group of benthic foraminifera from the Late Miocene Mediterranean. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 1999 , 152, 259-281 ⁹		94
12	Eemian Climatic and Hydrographical Instability on a Marine Shelf in Northern Denmark. <i>Quaternary Research</i> , 1997 , 47, 218-234	1.9	45
11	Two-step deglaciation at the oxygen isotope stage 6/5E transition: The Zeifen-Kattegat climate oscillation. <i>Quaternary Science Reviews</i> , 1996 , 15, 63-75	3.9	83
10	Quaternary palaeoceanography and palaeogeography in northern Denmark: a review of results from the Skagen cores. <i>Bulletin of the Geological Society of Denmark</i> , 1996 , 43, 22-31	1	24
9	&i>Cassidulina teretis&i>; Tappan and &i>Cassidulina neoteretis&i>; new species (Foraminifera): stratigraphic markers for deep sea and outer shelf areas. <i>Journal of Micropalaeontology</i> , 1995 , 14, 145-157	2	63
8	Marine evidence for climatic instability during the last interglacial in shelf records from northwest Europe. <i>Journal of Quaternary Science</i> , 1995 , 10, 77-82	2.3	68
7	Marine High Resolution Records of the Last Interglacial in Northwest Europe: A Review. <i>Géographie Physique Et Quaternaire</i> , 1994 , 48, 157-168		19
6	Biostratigraphy and palaeoenvironmental analysis of a Lower to Middle Jurassic succession on Anholt, Denmark. <i>Journal of Micropalaeontology</i> , 1993 , 12, 201-218	2	10
5	Benthic foraminiferal and stable isotope evidence for a Younger Dryas-style cold spell at the Saalian-Eemian transition, Denmark. <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 1993 , 102, 103-120 ^{2,9}		41
4	Plio-Pleistocene foraminiferal paleoecology and stratigraphy in the northernmost North Sea. <i>Journal of Foraminiferal Research</i> , 1992 , 22, 363-378	1.1	17
3	Investigations of past climate and sea-ice variability in the fjord area by Station Nord, eastern North Greenland. <i>Geological Survey of Denmark and Greenland Bulletin</i> , 35 , 67-70		2
2	The 8.2 ka cooling event related to extensive melting of the Greenland Ice Sheet		3
1	Early Holocene palaeoceanographic and glaciological changes in southeast Greenland. <i>Holocene</i> , 095968362210807		