

# Irena A Pidek

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3343439/publications.pdf>

Version: 2024-02-01

33  
papers

607  
citations

759233

12  
h-index

610901

24  
g-index

37  
all docs

37  
docs citations

37  
times ranked

940  
citing authors

| #  | ARTICLE                                                                                                                                                                                                                                            | IF  | CITATIONS |
|----|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|-----------|
| 1  | Two pollen-based methods of Eemian climate reconstruction employed in the study of the Å»abieniec-Jagodne palaeolakes in central Poland. <i>Quaternary International</i> , 2022, 632, 21-35.                                                       | 1.5 | 3         |
| 2  | Palaeoecological record of long Eemian series from KozÅ,Å³w (Central Poland) with reference to palaeoclimatic and palaeohydrological interpretation. <i>Quaternary International</i> , 2022, 632, 36-50.                                           | 1.5 | 4         |
| 3  | A high-resolution pollen and diatom record of mid-to late-Eemian at KozÅ,Å³w (Central Poland) reveals no drastic climate changes in the hornbeam phase of this interglacial. <i>Quaternary International</i> , 2021, 583, 14-30.                   | 1.5 | 5         |
| 4  | Environmental and climate change during the Late Saalianâ€Eemian Interglacial at the Struga site (Central Poland): A diatom record against the background of palynostratigraphy. <i>Review of Palaeobotany and Palynology</i> , 2021, 288, 104386. | 1.5 | 2         |
| 5  | Patterns in recent and Holocene pollen accumulation rates across Europe â€the Pollen Monitoring Programme Database as a tool for vegetation reconstruction. <i>Biogeosciences</i> , 2021, 18, 4511-4534.                                           | 3.3 | 5         |
| 6  | Petrographic features of tills as a tool in solving stratigraphical and palaeogeographical problems â€A case study from Central-Eastern Poland. <i>Quaternary International</i> , 2019, 501, 45-58.                                                | 1.5 | 10        |
| 7  | Environmental Conditions of Settlement of the Danubian Communities in the Northern Foreland of the Sandomierz Upland. <i>Archaeologia Polona</i> , 2019, 57, 213-231.                                                                              | 0.2 | 1         |
| 8  | The east-west migration of trees during the Eemian Interglacial registered on isopollen maps of Poland. <i>Quaternary International</i> , 2018, 467, 178-191.                                                                                      | 1.5 | 19        |
| 9  | Palaeoecological evolution of a spring-fed fen in PawÅ,Å³w (eastern Poland). <i>Grana</i> , 2018, 57, 345-363.                                                                                                                                     | 0.8 | 7         |
| 10 | Late Saalian and Eemian Interglacial at the Struga site (Garwolin Plain, central Poland). <i>Acta Palaeobotanica</i> , 2018, 58, 219-229.                                                                                                          | 0.7 | 4         |
| 11 | New palaeoclimate reconstructions based on multidisciplinary investigation in the FerdynandÅ³w 2011 stratotype site (eastern Poland). <i>Geological Quarterly</i> , 2017, , .                                                                      | 0.2 | 1         |
| 12 | Instability of the environment at the end of the Eemian Interglacial as illustrated by isopollen maps of Poland. <i>Geological Quarterly</i> , 2016, , .                                                                                           | 0.2 | 2         |
| 13 | SELECTED GEOCHEMICAL CRITERIA IN MIRE PROFILES OF PORÅ-BY WOJSÅAWSKIE (SANDOMIERZ BASIN,) Tj ETQq1 1 0.784314 rgBT 0,5 0                                                                                                                           | 0.5 | 0         |
| 14 | Palynostratigraphy and vegetation changes during the early Middle Pleistocene, based on new studies of deposits from FerdynandÅ³w (central eastern Poland). <i>Acta Palaeobotanica</i> , 2015, 55, 54-67.                                          | 0.7 | 2         |
| 15 | Taxon-specific pollen deposition dynamics in a temperate forest zone, SE Poland: the impact of physiological rhythmicity and weather controls. <i>Aerobiologia</i> , 2015, 31, 219-238.                                                            | 1.7 | 3         |
| 16 | Origin and evolution of the Bezedna lakeâ€mire complex in the Lublin area (East Poland): a case study for permafrost lakes in karstic regions. <i>Journal of Paleolimnology</i> , 2015, 53, 191-213.                                               | 1.6 | 10        |
| 17 | What was an interglacial river like? Sedimentological investigation of Holsteinian fluvial deposits in eastern Poland. <i>Geological Quarterly</i> , 2015, , .                                                                                     | 0.2 | 1         |
| 18 | How to resolve Pleistocene stratigraphic problems by different methods? A case study from eastern Poland. <i>Geological Quarterly</i> , 2014, 58, .                                                                                                | 0.2 | 3         |

| #  | ARTICLE                                                                                                                                                                                                                                                         | IF  | CITATIONS |
|----|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----|-----------|
| 19 | The European Modern Pollen Database (EMPD) project. <i>Vegetation History and Archaeobotany</i> , 2013, 22, 521-530.                                                                                                                                            | 2.1 | 101       |
| 20 | Pollen based quantitative climate reconstructions from the Middle Pleistocene sequences in $\text{Åuk\textsuperscript{3}w}$ and Zdany (E Poland): Species and modern analogues based approach. <i>Review of Palaeobotany and Palynology</i> , 2013, 192, 65-78. | 1.5 | 16        |
| 21 | Pollen percentage thresholds of <i>Abies alba</i> based on 13-year annual records of pollen deposition in modified Tauber traps: perspectives of application to fossil situations. <i>Review of Palaeobotany and Palynology</i> , 2013, 195, 26-36.             | 1.5 | 27        |
| 22 | Pollen-based vegetation and climate reconstruction of the Ferdynandovian sequence from $\text{Åuk\textsuperscript{3}w}$ (eastern Poland). <i>Acta Palaeobotanica</i> , 2013, 53, 115-138.                                                                       | 0.7 | 11        |
| 23 | The Holocene palaeoenvironmental changes reflected in the multi-proxy studies of Lake $\text{S\textsuperscript{A}one}$ sediments (SE Poland). <i>Palaeogeography, Palaeoclimatology, Palaeoecology</i> , 2012, 363-364, 79-98.                                  | 2.3 | 23        |
| 24 | Nine-year record of <i>Alnus</i> pollen deposition in the Roztocze region (SE Poland) with relation to vegetation data. <i>Acta Agrobotanica</i> , 2012, 60, 57-64.                                                                                             | 1.0 | 14        |
| 25 | Chronology and directions of Late Glacial paleoenvironmental changes: A multi-proxy study on sediments of Lake $\text{S\textsuperscript{A}one}$ (SE Poland). <i>Quaternary International</i> , 2011, 238, 89-106.                                               | 1.5 | 24        |
| 26 | Pollen dispersal and deposition characteristics of <i>Abies alba</i> , <i>Fagus sylvatica</i> and <i>Pinus sylvestris</i> , Roztocze region (SE Poland). <i>Vegetation History and Archaeobotany</i> , 2010, 19, 91-101.                                        | 2.1 | 68        |
| 27 | Variation in annual pollen accumulation rates of <i>Fagus</i> along a N-S transect in Europe based on pollen traps. <i>Vegetation History and Archaeobotany</i> , 2010, 19, 259-270.                                                                            | 2.1 | 41        |
| 28 | Annual pollen traps reveal the complexity of climatic control on pollen productivity in Europe and the Caucasus. <i>Vegetation History and Archaeobotany</i> , 2010, 19, 285-307.                                                                               | 2.1 | 51        |
| 29 | Comparing pollen spectra from modified Tauber traps and moss samples: examples from a selection of woodlands across Europe. <i>Vegetation History and Archaeobotany</i> , 2010, 19, 271-283.                                                                    | 2.1 | 65        |
| 30 | From early pollen trapping experiments to the Pollen Monitoring Programme. <i>Vegetation History and Archaeobotany</i> , 2010, 19, 247-258.                                                                                                                     | 2.1 | 61        |
| 31 | Pollen-vegetation relationships for pine and spruce in southeast Poland on the basis of volumetric and Tauber trap records. <i>Grana</i> , 2010, 49, 215-226.                                                                                                   | 0.8 | 17        |
| 32 | Older and Younger Holsteinian climate oscillations in the palaeobotanical record of the Brus profile (SE Poland). <i>Geological Quarterly</i> , 0, , .                                                                                                          | 0.2 | 0         |
| 33 | <i>Carpinus betulus</i> pollen accumulation rates in Roztocze (SE Poland) in relation to presence of <i>Carpinus</i> in Ferdynandovian pollen diagrams. <i>Ecological Questions</i> , 0, 26, 95.                                                                | 0.3 | 0         |