

Niels Wiinberg

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3345432/publications.pdf>

Version: 2024-02-01

53
papers

1,551
citations

304701

22
h-index

302107

39
g-index

53
all docs

53
docs citations

53
times ranked

2423
citing authors

#	ARTICLE	IF	CITATIONS
1	24-h ambulatory blood pressure in 352 normal Danish subjects, related to age and gender*. American Journal of Hypertension, 1995, 8, 978-986.	2.0	341
2	Smoking Related to 24-h Ambulatory Blood Pressure and Heart Rate A Study in 352 Normotensive Danish Subjects. American Journal of Hypertension, 1997, 10, 483-491.	2.0	88
3	Haemodynamic changes following treatment of subclinical and overt hyperthyroidism. European Journal of Endocrinology, 2001, 145, 391-396.	3.7	87
4	N-terminal-pro-B-type natriuretic peptide (NT-pro-BNP) in different thyroid function states. Clinical Endocrinology, 2004, 60, 54-59.	2.4	82
5	Noninvasive 24-hour ambulatory arterial blood pressure monitoring in cirrhosis. Hepatology, 1995, 22, 88-95.	7.3	76
6	Soya isoflavone-enriched cereal bars affect markers of endothelial function in postmenopausal women. British Journal of Nutrition, 2006, 95, 1120-1126.	2.3	55
7	Vitamin D Levels and Asymptomatic Coronary Artery Disease in Type 2 Diabetic Patients With Elevated Urinary Albumin Excretion Rate. Diabetes Care, 2012, 35, 168-172.	8.6	46
8	Diets Rich in Conjugated Linoleic Acid and Vaccenic Acid Have No Effect on Blood Pressure and Isoobaric Arterial Elasticity in Healthy Young Men. Journal of Nutrition, 2006, 136, 992-997.	2.9	43
9	Autonomic dysfunction in HIV patients on antiretroviral therapy: studies of heart rate variability. Clinical Physiology and Functional Imaging, 2007, 27, 363-367.	1.2	40
10	Effects of a Ginkgo biloba extract on forearm haemodynamics in healthy volunteers. Clinical Physiology and Functional Imaging, 2002, 22, 375-378.	1.2	38
11	Increased prevalence of coronary artery disease risk markers in patients with chronic hepatitis C – a cross-sectional study. Vascular Health and Risk Management, 2014, 10, 55.	2.3	36
12	Additive prognostic value of plasma N-terminal pro-brain natriuretic peptide and coronary artery calcification for cardiovascular events and mortality in asymptomatic patients with type 2 diabetes. Cardiovascular Diabetology, 2015, 14, 59.	6.8	35
13	Markers of collagen synthesis is related to blood pressure and vascular hypertrophy: a LIFE substudy. Journal of Human Hypertension, 2005, 19, 301-307.	2.2	33
14	Carotid intima-media thickness in individuals with and without type 2 diabetes: a reproducibility study. Cardiovascular Diabetology, 2010, 9, 40.	6.8	33
15	Daily Consumption for Six Weeks of a Lignan Complex Isolated from Flaxseed Does Not Affect Endothelial Function in Healthy Postmenopausal Women. Journal of Nutrition, 2006, 136, 2314-2318.	2.9	30
16	Metformin versus placebo in combination with insulin analogues in patients with type 2 diabetes mellitus”the randomised, blinded Copenhagen Insulin and Metformin Therapy (CIMT) trial. BMJ Open, 2016, 6, e008376.	1.9	30
17	Silent Ischemic Heart Disease and Pericardial Fat Volume in HIV-Infected Patients: A Case-Control Myocardial Perfusion Scintigraphy Study. PLoS ONE, 2013, 8, e72066.	2.5	30
18	Study rationale and design of the CIMT trial: The Copenhagen Insulin and Metformin Therapy Trial. Diabetes, Obesity and Metabolism, 2009, 11, 315-322.	4.4	26

#	ARTICLE	IF	CITATIONS
19	NT-proBNP levels, atherosclerosis and vascular function in asymptomatic type 2 diabetic patients with microalbuminuria: peripheral reactive hyperaemia index but not NT-proBNP is an independent predictor of coronary atherosclerosis. <i>Cardiovascular Diabetology</i> , 2011, 10, 71.	6.8	26
20	Interarm Difference in Blood Pressure: Reproducibility and Association with Peripheral Vascular Disease. <i>International Journal of Vascular Medicine</i> , 2014, 2014, 1-4.	1.0	26
21	Coronary and peripheral endothelial function in HIV patients studied with positron emission tomography and flow-mediated dilation: relation to hypercholesterolemia. <i>European Journal of Nuclear Medicine and Molecular Imaging</i> , 2008, 35, 2049-2058.	6.4	24
22	Elevated NT-proBNP and coronary calcium score in relation to coronary artery disease in asymptomatic type 2 diabetic patients with elevated urinary albumin excretion rate. <i>Nephrology Dialysis Transplantation</i> , 2011, 26, 3242-3249.	0.7	24
23	Effect of Intensive Multifactorial Treatment Compared With Routine Care on Aortic Stiffness and Central Blood Pressure Among Individuals With Screen-Detected Type 2 Diabetes. <i>Diabetes Care</i> , 2012, 35, 2207-2214.	8.6	24
24	Carotid intima-media thickness is reduced 12months after gastric bypass surgery in obese patients with type 2 diabetes or impaired glucose tolerance. <i>Journal of Diabetes and Its Complications</i> , 2014, 28, 517-522.	2.3	23
25	Carotid intima-media thickness in HIV patients treated with antiretroviral therapy. <i>Clinical Physiology and Functional Imaging</i> , 2007, 27, 173-179.	1.2	22
26	Oscillometric blood pressure measurement: a simple method in screening for peripheral arterial disease. <i>Clinical Physiology and Functional Imaging</i> , 2008, 28, 426-429.	1.2	19
27	Higher Physical Activity Is Associated With Lower Aortic Stiffness but Not With Central Blood Pressure. <i>Medicine (United States)</i> , 2015, 94, e485.	1.0	19
28	Reduction in coronary and peripheral vasomotor function in patients with HIV after initiation of antiretroviral therapy: a longitudinal study with positron emission tomography and flow-mediated dilation. <i>Nuclear Medicine Communications</i> , 2010, 31, 874-880.	1.1	18
29	24-h Ambulatory blood pressure in patients with ECG-determined left ventricular hypertrophy: left ventricular geometry and urinary albumin excretion—a LIFE substudy. <i>Journal of Human Hypertension</i> , 2004, 18, 391-396.	2.2	17
30	High prevalence of peripheral arterial disease in patients with previous cerebrovascular or coronary event. <i>Blood Pressure</i> , 2010, 19, 308-312.	1.5	14
31	Osteoprotegerin and coronary artery disease in type 2 diabetic patients with microalbuminuria. <i>Cardiovascular Diabetology</i> , 2011, 10, 70.	6.8	14
32	Diagnostic Value of the Nycocard®, Nycomed® D-Dimer Assay for the Diagnosis of Deep Venous Thrombosis and Pulmonary Embolism. <i>Thrombosis Research</i> , 2000, 100, 287-292.	1.7	13
33	Measurement of Arm Blood Pressure Using Different Oscillometry Manometers Compared to Auscultatory Readings. <i>Blood Pressure</i> , 2003, 12, 155-159.	1.5	13
34	Increased systolic ambulatory blood pressure and microalbuminuria in treated and non-treated hypertensive smokers. <i>Blood Pressure</i> , 2004, 13, 362-368.	1.5	13
35	Vitamin D, carotid intima-media thickness and bone structure in patients with type 2 diabetes. <i>Endocrine Connections</i> , 2015, 4, 128-135.	1.9	13
36	Losartan versus atenolol on 24-hour ambulatory blood pressure. A LIFE substudy. <i>Blood Pressure</i> , 2007, 16, 392-397.	1.5	12

#	ARTICLE	IF	CITATIONS
37	NT-proBNP, echocardiographic abnormalities and subclinical coronary artery disease in high risk type 2 diabetic patients. <i>Cardiovascular Diabetology</i> , 2012, 11, 19.	6.8	12
38	Effects of biphasic, basal-bolus or basal insulin analogue treatments on carotid intima-media thickness in patients with type 2 diabetes mellitus: the randomised Copenhagen Insulin and Metformin Therapy (CIMT) trial. <i>BMJ Open</i> , 2016, 6, e008377.	1.9	11
39	Metformin may adversely affect orthostatic blood pressure recovery in patients with type 2 diabetes: substudy from the placebo-controlled Copenhagen Insulin and Metformin Therapy (CIMT) trial. <i>Cardiovascular Diabetology</i> , 2020, 19, 150.	6.8	11
40	Does long-term losartan vs atenolol based antihypertensive treatment influence collagen markers differently in hypertensive patients? A LIFE substudy. <i>Blood Pressure</i> , 2006, 15, 198-206.	1.5	10
41	Isolated systolic hypertension in an elderly Danish population. Prevalence and daytime ambulatory blood pressure. <i>Blood Pressure</i> , 2006, 15, 347-353.	1.5	6
42	Carotid intima media thickness and ankle brachial index are inversely associated in subjects with and without diabetes. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2018, 78, 464-469.	1.2	5
43	Intensive versus conventional blood pressure monitoring in a general practice population. The Blood Pressure Reduction in Danish General Practice trial: a randomized controlled parallel group trial. <i>Family Practice</i> , 2018, 35, 433-439.	1.9	3
44	Flow-Mediated Vasodilatation and Intima-Media Thickness in Patients with Coexisting Heart Failure and Diabetes Receiving Medical Therapy. <i>Diagnostics</i> , 2011, 1, 38-52.	2.6	2
45	Measuring systolic ankle and toe pressure using the strain gauge technique – a comparison study between mercury and indium-gallium strain gauges. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2014, 74, 555-559.	1.2	2
46	Associations between glycaemic deterioration and aortic stiffness and central blood pressure. <i>Journal of Hypertension</i> , 2017, 35, 1832-1840.	0.5	2
47	Progression in risk factors during 36 years of follow-up and prediction of carotid intima-media thickness in a large cohort of adults with and without diabetes. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2020, 80, 491-499.	1.2	2
48	Prediction of carotid intima-media thickness and its relation to cardiovascular events in persons with type 2 diabetes. <i>Journal of Diabetes and Its Complications</i> , 2020, 34, 107681.	2.3	1
49	Conjugated linoleic acid (CLA) and vaccenic acid (VA) have no effect on blood pressure and isobar arterial elasticity in healthy young men. <i>FASEB Journal</i> , 2006, 20, A1016.	0.5	1
50	Comparison of brachial artery diameter measured by oscillometry and ultrasound. <i>American Journal of Hypertension</i> , 2003, 16, A148-A149.	2.0	0
51	Brachial artery compliance and ambulatory pulse pressure in hypertension accompanied by left ventricular hypertrophy. a life study. <i>American Journal of Hypertension</i> , 2004, 17, S136.	2.0	0
52	Brachial artery compliance and neurohormones in hypertension accompanied by left ventricular hypertrophy. a life study. <i>American Journal of Hypertension</i> , 2004, 17, S136-S137.	2.0	0
53	Noninvasive assessment of pulse-wave velocity and flow-mediated vasodilation in anesthetized C57Bl/6J mice. <i>Comparative Medicine</i> , 2014, 64, 471-7.	1.0	0