

Niels Wiinberg

List of Publications by Citations

Source: <https://exaly.com/author-pdf/3345432/niels-wiinberg-publications-by-citations.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

50
papers

1,277
citations

20
h-index

35
g-index

53
ext. papers

1,372
ext. citations

4.1
avg, IF

3.51
L-index

#	Paper	IF	Citations
50	24-h ambulatory blood pressure in 352 normal Danish subjects, related to age and gender. <i>American Journal of Hypertension</i> , 1995 , 8, 978-86	2.3	276
49	N-terminal-pro-B-type natriuretic peptide (NT-pro-BNP) in different thyroid function states. <i>Clinical Endocrinology</i> , 2004 , 60, 54-9	3.4	73
48	Noninvasive 24-hour ambulatory arterial blood pressure monitoring in cirrhosis. <i>Hepatology</i> , 1995 , 22, 88-95	11.2	69
47	Haemodynamic changes following treatment of subclinical and overt hyperthyroidism. <i>European Journal of Endocrinology</i> , 2001 , 145, 391-6	6.5	68
46	Smoking related to 24-h ambulatory blood pressure and heart rate: a study in 352 normotensive Danish subjects. <i>American Journal of Hypertension</i> , 1997 , 10, 483-91	2.3	66
45	Soya isoflavone-enriched cereal bars affect markers of endothelial function in postmenopausal women. <i>British Journal of Nutrition</i> , 2006 , 95, 1120-6	3.6	49
44	Vitamin D levels and asymptomatic coronary artery disease in type 2 diabetic patients with elevated urinary albumin excretion rate. <i>Diabetes Care</i> , 2012 , 35, 168-72	14.6	40
43	Diets rich in conjugated linoleic acid and vaccenic acid have no effect on blood pressure and isobaric arterial elasticity in healthy young men. <i>Journal of Nutrition</i> , 2006 , 136, 992-7	4.1	36
42	Autonomic dysfunction in HIV patients on antiretroviral therapy: studies of heart rate variability. <i>Clinical Physiology and Functional Imaging</i> , 2007 , 27, 363-7	2.4	31
41	Markers of collagen synthesis is related to blood pressure and vascular hypertrophy: a LIFE substudy. <i>Journal of Human Hypertension</i> , 2005 , 19, 301-7	2.6	31
40	Effects of a Ginkgo biloba extract on forearm haemodynamics in healthy volunteers. <i>Clinical Physiology and Functional Imaging</i> , 2002 , 22, 375-8	2.4	30
39	Increased prevalence of coronary artery disease risk markers in patients with chronic hepatitis C--a cross-sectional study. <i>Vascular Health and Risk Management</i> , 2014 , 10, 55-62	4.4	29
38	Additive prognostic value of plasma N-terminal pro-brain natriuretic peptide and coronary artery calcification for cardiovascular events and mortality in asymptomatic patients with type 2 diabetes. <i>Cardiovascular Diabetology</i> , 2015 , 14, 59	8.7	28
37	Carotid intima-media thickness in individuals with and without type 2 diabetes: a reproducibility study. <i>Cardiovascular Diabetology</i> , 2010 , 9, 40	8.7	28
36	Silent ischemic heart disease and pericardial fat volume in HIV-infected patients: a case-control myocardial perfusion scintigraphy study. <i>PLoS ONE</i> , 2013 , 8, e72066	3.7	28
35	Metformin versus placebo in combination with insulin analogues in patients with type 2 diabetes mellitus-the randomised, blinded Copenhagen Insulin and Metformin Therapy (CIMT) trial. <i>BMJ Open</i> , 2016 , 6, e008376	3	28
34	NT-proBNP levels, atherosclerosis and vascular function in asymptomatic type 2 diabetic patients with microalbuminuria: peripheral reactive hyperaemia index but not NT-proBNP is an independent predictor of coronary atherosclerosis. <i>Cardiovascular Diabetology</i> , 2011 , 10, 71	8.7	25

33	Daily consumption for six weeks of a lignan complex isolated from flaxseed does not affect endothelial function in healthy postmenopausal women. <i>Journal of Nutrition</i> , 2006 , 136, 2314-8	4.1	25
32	Study rationale and design of the CIMT trial: the Copenhagen Insulin and Metformin Therapy trial. <i>Diabetes, Obesity and Metabolism</i> , 2009 , 11, 315-22	6.7	24
31	Elevated NT-proBNP and coronary calcium score in relation to coronary artery disease in asymptomatic type 2 diabetic patients with elevated urinary albumin excretion rate. <i>Nephrology Dialysis Transplantation</i> , 2011 , 26, 3242-9	4.3	22
30	Effect of intensive multifactorial treatment compared with routine care on aortic stiffness and central blood pressure among individuals with screen-detected type 2 diabetes: the ADDITION-Denmark study. <i>Diabetes Care</i> , 2012 , 35, 2207-14	14.6	19
29	Interarm difference in blood pressure: reproducibility and association with peripheral vascular disease. <i>International Journal of Vascular Medicine</i> , 2014 , 2014, 841542	1.2	18
28	Carotid intima-media thickness is reduced 12 months after gastric bypass surgery in obese patients with type 2 diabetes or impaired glucose tolerance. <i>Journal of Diabetes and Its Complications</i> , 2014 , 28, 517-22	3.2	18
27	Coronary and peripheral endothelial function in HIV patients studied with positron emission tomography and flow-mediated dilation: relation to hypercholesterolemia. <i>European Journal of Nuclear Medicine and Molecular Imaging</i> , 2008 , 35, 2049-58	8.8	17
26	24-h Ambulatory blood pressure in patients with ECG-determined left ventricular hypertrophy: left ventricular geometry and urinary albumin excretion-a LIFE substudy. <i>Journal of Human Hypertension</i> , 2004 , 18, 391-6	2.6	17
25	Carotid intima-media thickness in HIV patients treated with antiretroviral therapy. <i>Clinical Physiology and Functional Imaging</i> , 2007 , 27, 173-9	2.4	16
24	Oscillometric blood pressure measurement: a simple method in screening for peripheral arterial disease. <i>Clinical Physiology and Functional Imaging</i> , 2008 , 28, 426-9	2.4	15
23	Reduction in coronary and peripheral vasomotor function in patients with HIV after initiation of antiretroviral therapy: a longitudinal study with positron emission tomography and flow-mediated dilation. <i>Nuclear Medicine Communications</i> , 2010 , 31, 874-80	1.6	14
22	High prevalence of peripheral arterial disease in patients with previous cerebrovascular or coronary event. <i>Blood Pressure</i> , 2010 , 19, 308-12	1.7	12
21	Measurement of arm blood pressure using different oscillometry manometers compared to auscultatory readings. <i>Blood Pressure</i> , 2003 , 12, 155-9	1.7	12
20	Increased systolic ambulatory blood pressure and microalbuminuria in treated and non-treated hypertensive smokers. <i>Blood Pressure</i> , 2004 , 13, 362-8	1.7	12
19	Higher physical activity is associated with lower aortic stiffness but not with central blood pressure: the ADDITION-Pro Study. <i>Medicine (United States)</i> , 2015 , 94, e485	1.8	11
18	NT-proBNP, echocardiographic abnormalities and subclinical coronary artery disease in high risk type 2 diabetic patients. <i>Cardiovascular Diabetology</i> , 2012 , 11, 19	8.7	11
17	Vitamin D, carotid intima-media thickness and bone structure in patients with type 2 diabetes. <i>Endocrine Connections</i> , 2015 , 4, 128-35	3.5	10
16	Osteoprotegerin and coronary artery disease in type 2 diabetic patients with microalbuminuria. <i>Cardiovascular Diabetology</i> , 2011 , 10, 70	8.7	10

15	Losartan versus atenolol on 24-hour ambulatory blood pressure. A LIFE substudy. <i>Blood Pressure</i> , 2007 , 16, 392-7	1.7	10
14	Diagnostic value of the Nycocard, Nycomed D-dimer assay for the diagnosis of deep venous thrombosis and pulmonary embolism: a retrospective study. <i>Thrombosis Research</i> , 2000 , 100, 287-92	8.2	10
13	Effects of biphasic, basal-bolus or basal insulin analogue treatments on carotid intima-media thickness in patients with type 2 diabetes mellitus: the randomised Copenhagen Insulin and Metformin Therapy (CIMT) trial. <i>BMJ Open</i> , 2016 , 6, e008377	3	9
12	Does long-term losartan- vs atenolol-based antihypertensive treatment influence collagen markers differently in hypertensive patients? A LIFE substudy. <i>Blood Pressure</i> , 2006 , 15, 198-206	1.7	8
11	Isolated systolic hypertension in an elderly Danish population. Prevalence and daytime ambulatory blood pressure. <i>Blood Pressure</i> , 2006 , 15, 347-53	1.7	6
10	Carotid intima media thickness and ankle brachial index are inversely associated in subjects with and without diabetes. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2018 , 78, 464-469	2	4
9	Metformin may adversely affect orthostatic blood pressure recovery in patients with type 2 diabetes: substudy from the placebo-controlled Copenhagen Insulin and Metformin Therapy (CIMT) trial. <i>Cardiovascular Diabetology</i> , 2020 , 19, 150	8.7	4
8	Associations between glycaemic deterioration and aortic stiffness and central blood pressure: the ADDITION-PRO Study. <i>Journal of Hypertension</i> , 2017 , 35, 1832-1840	1.9	2
7	Measuring systolic ankle and toe pressure using the strain gauge technique--a comparison study between mercury and indium-gallium strain gauges. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2014 , 74, 555-9	2	2
6	Flow-Mediated Vasodilatation and Intima-Media Thickness in Patients with Coexisting Heart Failure and Diabetes Receiving Medical Therapy. <i>Diagnostics</i> , 2011 , 1, 38-52	3.8	2
5	Intensive versus conventional blood pressure monitoring in a general practice population. The Blood Pressure Reduction in Danish General Practice trial: a randomized controlled parallel group trial. <i>Family Practice</i> , 2018 , 35, 433-439	1.9	1
4	Progression in risk factors during 36 years of follow-up and prediction of carotid intima-media thickness in a large cohort of adults with and without diabetes. <i>Scandinavian Journal of Clinical and Laboratory Investigation</i> , 2020 , 80, 491-499	2	1
3	Prediction of carotid intima-media thickness and its relation to cardiovascular events in persons with type 2 diabetes. <i>Journal of Diabetes and Its Complications</i> , 2020 , 34, 107681	3.2	0
2	Noninvasive assessment of pulse-wave velocity and flow-mediated vasodilation in anesthetized Göttingen minipigs. <i>Comparative Medicine</i> , 2014 , 64, 471-7	1.6	
1	Conjugated linoleic acid (CLA) and vaccenic acid (VA) have no effect on blood pressure and isobar arterial elasticity in healthy young men. <i>FASEB Journal</i> , 2006 , 20, A1016	0.9	