

# Long Chen

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3377643/publications.pdf>

Version: 2024-02-01

57  
papers

2,885  
citations

230014

27  
h-index

190340

53  
g-index

61  
all docs

61  
docs citations

61  
times ranked

4107  
citing authors

| #  | ARTICLE  | IF  | CITATIONS |
|----|--|-----|-----------|
| 1  | Introducing the Solvent Co-intercalation Mechanism for Hard Carbon with Ultrafast Sodium Storage. <i>Small</i> , 2022, 18, e2108092.   | 5.2 | 14        |
| 2  | NOX2-derived hydrogen peroxide impedes the AMPK/Akt-mTOR signaling pathway contributing to cell death in neuronal cells. <i>Cellular Signalling</i> , 2022, 94, 110330.  | 1.7 | 4         |
| 3  | Resveratrol induces autophagy impeding BAFF-stimulated B-cell proliferation and survival by inhibiting the Akt/mTOR pathway. <i>Biochemical Pharmacology</i> , 2022, 202, 115139.  | 2.0 | 8         |
| 4  | Cadmium Impairs Autophagy Leading to Apoptosis by Ca <sup>2+</sup> -Dependent Activation of JNK Signaling Pathway in Neuronal Cells. <i>Neurochemical Research</i> , 2021, 46, 2033-2045.  | 1.6 | 11        |
| 5  | Surface enrichment and diffusion enabling gradient-doping and coating of Ni-rich cathode toward Li-ion batteries. <i>Nature Communications</i> , 2021, 12, 4564.   | 5.8 | 153       |
| 6  | Metformin prevents BAFF activation of Erk1/2 from B-cell proliferation and survival by impeding mTOR-PTEN/Akt signaling pathway. <i>International Immunopharmacology</i> , 2021, 96, 107771.   | 1.7 | 7         |
| 7  | Magnesium isoglycyrrhizinate prevents cadmium-induced activation of JNK and apoptotic hepatocyte death by reversing ROS-inactivated PP2A. <i>Journal of Pharmacy and Pharmacology</i> , 2021, 73, 1663-1674.   | 1.2 | 6         |
| 8  | Metformin attenuates cadmium-induced neuronal apoptosis in vitro via blocking ROS-dependent PP5/AMPK-JNK signaling pathway. <i>Neuropharmacology</i> , 2020, 175, 108065.  | 2.0 | 26        |
| 9  | Cadmium induces mitochondrial ROS inactivation of XIAP pathway leading to apoptosis in neuronal cells. <i>International Journal of Biochemistry and Cell Biology</i> , 2020, 121, 105715.  | 1.2 | 30        |
| 10 | Rapamycin inhibits B-cell activating factor (BAFF)-stimulated cell proliferation and survival by suppressing Ca <sup>2+</sup> -CaMKII-dependent PTEN/Akt-Erk1/2 signaling pathway in normal and neoplastic B-lymphoid cells. <i>Cell Calcium</i> , 2020, 87, 102171. | 1.1 | 18        |
| 11 | Resveratrol inhibits Erk1/2-mediated adhesion of cancer cells via activating PP2A-PTEN signaling network. <i>Journal of Cellular Physiology</i> , 2019, 234, 2822-2836.  | 2.0 | 13        |
| 12 | Maduramicin inactivation of Akt impairs autophagic flux leading to accumulated autophagosomes-dependent apoptosis in skeletal myoblast cells. <i>International Journal of Biochemistry and Cell Biology</i> , 2019, 114, 105573.                                     | 1.2 | 3         |
| 13 | Morphologically Controlled Synthesis of Cs <sub>2</sub> SnCl <sub>6</sub> Perovskite Crystals and Their Photoluminescence Activity. <i>Crystals</i> , 2019, 9, 258.  | 1.0 | 13        |
| 14 | Photoluminescence Characteristics of Sn <sup>2+</sup> and Ce <sup>3+</sup> -Doped Cs <sub>2</sub> SnCl <sub>6</sub> Double-Perovskite Crystals. <i>Materials</i> , 2019, 12, 1501.   | 1.3 | 29        |
| 15 | Cadmium results in accumulation of autophagosomes-dependent apoptosis through activating Akt-impaired autophagic flux in neuronal cells. <i>Cellular Signalling</i> , 2019, 55, 26-39.   | 1.7 | 45        |
| 16 | Maduramicin induces cardiac muscle cell death by the ROS-dependent PTEN/Akt-Erk1/2 signaling pathway. <i>Journal of Cellular Physiology</i> , 2019, 234, 10964-10976.  | 2.0 | 18        |
| 17 | BAFF inhibits autophagy promoting cell proliferation and survival by activating Ca <sup>2+</sup> -CaMKII-dependent Akt/mTOR signaling pathway in normal and neoplastic B-lymphoid cells. <i>Cellular Signalling</i> , 2019, 53, 68-79.                               | 1.7 | 29        |
| 18 | Rapamycin attenuates BAFF-extended proliferation and survival via disruption of mTORC1/2 signaling in normal and neoplastic B-lymphoid cells. <i>Journal of Cellular Physiology</i> , 2018, 233, 516-529.  | 2.0 | 20        |

| #  | ARTICLE   | IF  | CITATIONS |
|----|---|-----|-----------|
| 19 | Maduramicin induces apoptosis and necrosis, and blocks autophagic flux in myocardial H9c2 cells. <i>Journal of Applied Toxicology</i> , 2018, 38, 366-375.  | 1.4 | 12        |
| 20 | Celastrol ameliorates Cd <sup>2+</sup> -induced neuronal apoptosis by targeting NOX2 <sup>+</sup> -derived ROS <sup>+</sup> -dependent PP5 <sup>+</sup> /JNK signaling pathway. <i>Journal of Neurochemistry</i> , 2017, 141, 48-62.                              | 2.1 | 37        |
| 21 | Celastrol Attenuates Cadmium <sup>2+</sup> -Induced Neuronal Apoptosis via Inhibiting Ca <sup>2+</sup> -CaMKII <sup>+</sup> -Dependent Akt/mTOR Pathway. <i>Journal of Cellular Physiology</i> , 2017, 232, 2145-2157.  | 2.0 | 34        |
| 22 | Celastrol prevents cadmium <sup>2+</sup> -induced neuronal cell death by blocking reactive oxygen species <sup>+</sup> -mediated mammalian target of rapamycin pathway. <i>British Journal of Pharmacology</i> , 2017, 174, 82-100.                               | 2.7 | 37        |
| 23 | An insight of rapamycin against cadmium <sup>2+</sup> 's neurotoxicity. <i>Oncotarget</i> , 2017, 8, 9013-9014.   | 0.8 | 2         |
| 24 | IL-2, IL-4, IFN- $\gamma$ or TNF- $\alpha$ enhances BAFF-stimulated cell viability and survival by activating Erk1/2 and S6K1 pathways in neoplastic B-lymphoid cells. <i>Cytokine</i> , 2016, 84, 37-46.   | 1.4 | 14        |
| 25 | An investigation into the use of six facially encoded emotions in brain-computer interfacing. <i>Brain-Computer Interfaces</i> , 2016, 3, 59-73.  | 0.9 | 6         |
| 26 | Rapamycin ameliorates cadmium-induced activation of MAPK pathway and neuronal apoptosis by preventing mitochondrial ROS inactivation of PP2A. <i>Neuropharmacology</i> , 2016, 105, 270-284.  | 2.0 | 56        |
| 27 | Crosstalk between Ca <sup>2+</sup> signaling and mitochondrial H <sub>2</sub> O <sub>2</sub> is required for rotenone inhibition of mTOR signaling pathway leading to neuronal apoptosis. <i>Oncotarget</i> , 2016, 7, 7534-7549.                                 | 0.8 | 26        |
| 28 | Resveratrol prevents cadmium activation of Erk1/2 and JNK pathways from neuronal cell death via protein phosphatases 2A and 5. <i>Journal of Neurochemistry</i> , 2015, 135, 466-478.   | 2.1 | 31        |
| 29 | Rapamycin prevents cadmium-induced neuronal cell death via targeting both mTORC1 and mTORC2 pathways. <i>Neuropharmacology</i> , 2015, 97, 35-45.   | 2.0 | 22        |
| 30 | Rapamycin inhibits BAFF-stimulated cell proliferation and survival by suppressing mTOR-mediated PP2A-Erk1/2 signaling pathway in normal and neoplastic B-lymphoid cells. <i>Cellular and Molecular Life Sciences</i> , 2015, 72, 4867-4884.                       | 2.4 | 42        |
| 31 | A deut of mTORC1/2 for cell adhesion. <i>Cell Cycle</i> , 2015, 14, 1131-1132.  | 1.3 | 1         |
| 32 | Rotenone Induction of Hydrogen Peroxide Inhibits mTOR-mediated S6K1 and 4E-BP1/eIF4E Pathways, Leading to Neuronal Apoptosis. <i>Toxicological Sciences</i> , 2015, 143, 81-96.   | 1.4 | 90        |
| 33 | Both mTORC1 and mTORC2 are involved in the regulation of cell adhesion. <i>Oncotarget</i> , 2015, 6, 7136-7150.   | 0.8 | 33        |
| 34 | Rapamycin inhibits Erk1/2-mediated neuronal apoptosis caused by cadmium. <i>Oncotarget</i> , 2015, 6, 21452-21467.  | 0.8 | 11        |
| 35 | N <sup>6</sup> -acetyl-L-cysteine protects against cadmium <sup>2+</sup> -induced neuronal apoptosis by inhibiting ROS <sup>+</sup> -dependent activation of Akt/mTOR pathway in mouse brain. <i>Neuropathology and Applied Neurobiology</i> , 2014, 40, 759-777. | 1.8 | 96        |
| 36 | Celastrol prevents cadmium <sup>2+</sup> -induced neuronal cell death via targeting JNK and PTEN <sup>+</sup> /Akt/mTOR network. <i>Journal of Neurochemistry</i> , 2014, 128, 256-266.   | 2.1 | 44        |

| #  | ARTICLE   | IF  | CITATIONS |
|----|---|-----|-----------|
| 37 | BAFF activates Erk1/2 promoting cell proliferation and survival by Ca <sup>2+</sup> -CaMKII-dependent inhibition of PP2A in normal and neoplastic B-lymphoid cells. <i>Biochemical Pharmacology</i> , 2014, 87, 332-343.                | 2.0 | 20        |
| 38 | Activation of AMPK and inactivation of Akt result in suppression of mTOR-mediated S6K1 and 4E-BP1 pathways leading to neuronal cell death in in vitro models of Parkinson's disease. <i>Cellular Signalling</i> , 2014, 26, 1680-1689.  | 1.7 | 133       |
| 39 | hsBAFF promotes proliferation and survival in cultured B lymphocytes via calcium signaling activation of mTOR pathway. <i>Cytokine</i> , 2013, 62, 310-321.   | 1.4 | 25        |
| 40 | Curcumin inhibits protein phosphatases 2A and 5, leading to activation of mitogen-activated protein kinases and death in tumor cells. <i>Carcinogenesis</i> , 2012, 33, 868-875.  | 1.3 | 68        |
| 41 | hsBAFF regulates proliferation and response in cultured CD4 <sup>+</sup> T lymphocytes by upregulation of intracellular free Ca <sup>2+</sup> homeostasis. <i>Cytokine</i> , 2011, 53, 215-222.   | 1.4 | 18        |
| 42 | CaMKII is involved in cadmium activation of MAPK and mTOR pathways leading to neuronal cell death. <i>Journal of Neurochemistry</i> , 2011, 119, 1108-1118.   | 2.1 | 85        |
| 43 | Cadmium induction of reactive oxygen species activates the mTOR pathway, leading to neuronal cell death. <i>Free Radical Biology and Medicine</i> , 2011, 50, 624-632.  | 1.3 | 214       |
| 44 | Calcium Signaling Is Involved in Cadmium-Induced Neuronal Apoptosis via Induction of Reactive Oxygen Species and Activation of MAPK/mTOR Network. <i>PLoS ONE</i> , 2011, 6, e19052.  | 1.1 | 158       |
| 45 | Hydrogen peroxide inhibits mTOR signaling by activation of AMPK $\pm$ leading to apoptosis of neuronal cells. <i>Laboratory Investigation</i> , 2010, 90, 762-773.  | 1.7 | 207       |
| 46 | Rapamycin Inhibits IGF-1 Stimulated Cell Motility through PP2A Pathway. <i>PLoS ONE</i> , 2010, 5, e10578.  | 1.1 | 36        |
| 47 | Specific IgG activity against diarrheagenic bacteria in bovine immune milk and effect of pH on its antigen-binding activity upon heating. <i>Journal of Dairy Research</i> , 2010, 77, 220-224.   | 0.7 | 5         |
| 48 | Rapamycin Inhibits Cytoskeleton Reorganization and Cell Motility by Suppressing RhoA Expression and Activity. <i>Journal of Biological Chemistry</i> , 2010, 285, 38362-38373.  | 1.6 | 120       |
| 49 | Hydrogen peroxide-induced neuronal apoptosis is associated with inhibition of protein phosphatase 2A and 5, leading to activation of MAPK pathway. <i>International Journal of Biochemistry and Cell Biology</i> , 2009, 41, 1284-1295. | 1.2 | 204       |
| 50 | BAFF enhances B-cell-mediated immune response and vaccine-protection against a very virulent IBDV in chickens. <i>Vaccine</i> , 2009, 27, 1393-1399.  | 1.7 | 16        |
| 51 | MAPK and mTOR pathways are involved in cadmium-induced neuronal apoptosis. <i>Journal of Neurochemistry</i> , 2008, 105, 251-261.   | 2.1 | 134       |
| 52 | Cadmium activates the mitogen-activated protein kinase (MAPK) pathway via induction of reactive oxygen species and inhibition of protein phosphatases 2A and 5. <i>Free Radical Biology and Medicine</i> , 2008, 45, 1035-1044.         | 1.3 | 231       |
| 53 | Specific IgG activity of bovine immune milk against diarrhea bacteria and its protective effects on pathogen-infected intestinal damages. <i>Vaccine</i> , 2008, 26, 5973-5980.   | 1.7 | 11        |
| 54 | Enriched Selenium and Its Effects on Growth and Biochemical Composition in <i>Lactobacillus bulgaricus</i> . <i>Journal of Agricultural and Food Chemistry</i> , 2007, 55, 2413-2417.   | 2.4 | 64        |

| #  | ARTICLE  | IF  | CITATIONS |
|----|--|-----|-----------|
| 55 | Bovine immune colostrum against 17 strains of diarrhea bacteria and in vitro and in vivo effects of its specific IgG. <i>Vaccine</i> , 2006, 24, 2131-2140.                                    | 1.7 | 13        |
| 56 | Protective effect of selenium-enriched lactobacillus on CCl <sub>4</sub> -induced liver injury in mice and its possible mechanisms. <i>World Journal of Gastroenterology</i> , 2005, 11, 5795. | 1.4 | 67        |
| 57 | Action of NO and TNF-alpha release of rats with cadmium loading in malfunction of multiple system organ. <i>Acta Physiologica Sinica</i> , 2003, 55, 535-40.                                   | 0.5 | 11        |