

Arndt's Fä“lafsdÄ³ttir Pgd

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3393101/publications.pdf>

Version: 2024-02-01

10
papers

968
citations

1040056

9
h-index

1372567

10
g-index

10
all docs

10
docs citations

10
times ranked

1244
citing authors

#	ARTICLE	IF	CITATIONS
1	Continuous Glucose Monitoring vs Conventional Therapy for Glycemic Control in Adults With Type 1 Diabetes Treated With Multiple Daily Insulin Injections. JAMA - Journal of the American Medical Association, 2017, 317, 379.	7.4	520
2	A Clinical Trial of the Accuracy and Treatment Experience of the Flash Glucose Monitor FreeStyle Libre in Adults with Type 1 Diabetes. Diabetes Technology and Therapeutics, 2017, 19, 164-172.	4.4	143
3	HbA _{1c} level as a risk factor for retinopathy and nephropathy in children and adults with type 1 diabetes: Swedish population based cohort study. BMJ: British Medical Journal, 2019, 366, 14894.	2.3	109
4	A Randomized Clinical Trial of the Effect of Continuous Glucose Monitoring on Nocturnal Hypoglycemia, Daytime Hypoglycemia, Glycemic Variability, and Hypoglycemia Confidence in Persons with Type 1 Diabetes Treated with Multiple Daily Insulin Injections (GOLD-3). Diabetes Technology and Therapeutics, 2018, 20, 274-284.	4.4	88
5	The Association Between HbA _{1c} and Time in Hypoglycemia During CGM and Self-Monitoring of Blood Glucose in People With Type 1 Diabetes and Multiple Daily Insulin Injections: A Randomized Clinical Trial (GOLD-4). Diabetes Care, 2020, 43, 2017-2024.	8.6	34
6	Sustained Intensive Treatment and Long-term Effects on HbA _{1c} Reduction (SILVER Study) by CGM in People With Type 1 Diabetes Treated With MDI. Diabetes Care, 2021, 44, 141-149.	8.6	19
7	Excess risk of lower extremity amputations in people with type 1 diabetes compared with the general population: amputations and type 1 diabetes. BMJ Open Diabetes Research and Care, 2019, 7, e000602.	2.8	17
8	Risk Factors for Atrial Fibrillation in People With Type 1 Diabetes: An Observational Cohort Study of 36,258 Patients From the Swedish National Diabetes Registry. Diabetes Care, 2019, 42, 1530-1538.	8.6	16
9	Risk factors and incidence over time for lower extremity amputations in people with type 1 diabetes: an observational cohort study of 46,088 patients from the Swedish National Diabetes Registry. Diabetologia, 2021, 64, 2751-2761.	6.3	13
10	The majority of people with type 1 diabetes and multiple daily insulin injections benefit from using continuous glucose monitoring: An analysis based on the GOLD randomized trial (GOLD-5). Diabetes, Obesity and Metabolism, 2021, 23, 619-630.	4.4	9