

Masashi Hyodo

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/33941/publications.pdf>

Version: 2024-02-01

11
papers

50
citations

2258059

3
h-index

1872680

6
g-index

11
all docs

11
docs citations

11
times ranked

42
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 1 | Testing block-diagonal covariance structure for high-dimensional data. <i>Statistica Neerlandica</i> , 2015, 69, 460-482. | 1.6 | 13 |
| 2 | Testing block-diagonal covariance structure for high-dimensional data under non-normality. <i>Journal of Multivariate Analysis</i> , 2017, 155, 305-316. | 1.0 | 8 |
| 3 | A simultaneous testing of the mean vector and the covariance matrix among two populations for high-dimensional data. <i>Test</i> , 2018, 27, 680-699. | 1.1 | 4 |
| 4 | Evaluation of multinomial logistic regression models for predicting causative pathogens of food poisoning cases. <i>Journal of Veterinary Medical Science</i> , 2018, 80, 1223-1227. | 0.9 | 4 |
| 5 | On simultaneous confidence interval estimation for the difference of paired mean vectors in high-dimensional settings. <i>Journal of Multivariate Analysis</i> , 2018, 168, 160-173. | 1.0 | 4 |
| 6 | Asymptotic power comparison of T^2 -type test and likelihood ratio test for a mean vector based on two-step monotone missing data. <i>Communications in Statistics - Theory and Methods</i> , 2020, 49, 4270-4287. | 1.0 | 4 |
| 7 | Simultaneous testing of the mean vector and covariance matrix among k populations for high-dimensional data. <i>Communications in Statistics - Theory and Methods</i> , 2021, 50, 663-684. | 1.0 | 4 |
| 8 | Estimation of the covariance matrix with two-step monotone missing data. <i>Communications in Statistics - Theory and Methods</i> , 2016, 45, 1910-1922. | 1.0 | 3 |
| 9 | Tests for the parallelism and flatness hypotheses of multi-group profile analysis for high-dimensional elliptical populations. <i>Journal of Multivariate Analysis</i> , 2017, 162, 82-92. | 1.0 | 3 |
| 10 | Bartlett correction to the likelihood ratio test for MCAR with two-step monotone sample. <i>Statistica Neerlandica</i> , 2017, 71, 184-199. | 1.6 | 3 |
| 11 | A one-sample location test based on weighted averaging of two test statistics when the dimension and the sample size are large. <i>Communications in Statistics - Theory and Methods</i> , 2017, 46, 3526-3541. | 1.0 | 0 |