

Caroline C Duwaerts

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3394642/publications.pdf>

Version: 2024-02-01

19
papers

422
citations

759233

12
h-index

839539

18
g-index

22
all docs

22
docs citations

22
times ranked

761
citing authors

#	ARTICLE	IF	CITATIONS
1	Targeting acid ceramidase inhibits YAP/TAZ signaling to reduce fibrosis in mice. <i>Science Translational Medicine</i> , 2020, 12, .	12.4	71
2	Macronutrients and the Adipose-Liver Axis in Obesity and Fatty Liver. <i>Cellular and Molecular Gastroenterology and Hepatology</i> , 2019, 7, 749-761.	4.5	63
3	Polycystic ovary syndrome (PCOS) is associated with NASH severity and advanced fibrosis. <i>Liver International</i> , 2020, 40, 355-359.	3.9	50
4	Mechanisms of Liver Injury in Non-Alcoholic Steatohepatitis. <i>Current Hepatology Reports</i> , 2014, 13, 119-129.	0.9	37
5	Targeting the diverse immunological functions expressed by hepatic NKT cells. <i>Expert Opinion on Therapeutic Targets</i> , 2011, 15, 973-988.	3.4	23
6	NK cells suppress experimental cholestatic liver injury by an interleukin-6-mediated, Kupffer cell-dependent mechanism. <i>Journal of Hepatology</i> , 2011, 54, 746-752.	3.7	21
7	Hepatocyte-specific deletion of XBP1 sensitizes mice to liver injury through hyperactivation of IRE1 α . <i>Cell Death and Differentiation</i> , 2021, 28, 1455-1465.	11.2	20
8	Contrasting responses of Kupffer cells and inflammatory mononuclear phagocytes to biliary obstruction in a mouse model of cholestatic liver injury. <i>Liver International</i> , 2013, 33, 255-265.	3.9	19
9	Cross-Activating Invariant NKT Cells and Kupffer Cells Suppress Cholestatic Liver Injury in a Mouse Model of Biliary Obstruction. <i>PLoS ONE</i> , 2013, 8, e79702.	2.5	19
10	Isocaloric manipulation of macronutrients within a high-carbohydrate/moderate-fat diet induces unique effects on hepatic lipogenesis, steatosis and liver injury. <i>Journal of Nutritional Biochemistry</i> , 2016, 29, 12-20.	4.2	16
11	The Association of Hispanic Ethnicity with Nonalcoholic Fatty Liver Disease in Polycystic Ovary Syndrome. <i>Current Opinion in Gynecology and Obstetrics</i> , 2018, 1, 24-33.	0.0	16
12	Effects of recovery from immobilization stress on striatal preprodynorphin- and kappa opioid receptor-mRNA levels of the male rat. <i>Physiology and Behavior</i> , 2011, 104, 972-980.	2.1	15
13	Specific Macronutrients Exert Unique Influences on the Adipose-Liver Axis to Promote Hepatic Steatosis in Mice. <i>Cellular and Molecular Gastroenterology and Hepatology</i> , 2017, 4, 223-236.	4.5	13
14	Induced Pluripotent Stem Cell-derived Hepatocytes From Patients With Nonalcoholic Fatty Liver Disease Display a Disease-specific Gene Expression Profile. <i>Gastroenterology</i> , 2021, 160, 2591-2594.e6.	1.3	13
15	CD18 deficiency improves liver injury in the MCD model of steatohepatitis. <i>PLoS ONE</i> , 2017, 12, e0183912.	2.5	8
16	Doxycycline Significantly Enhances Induction of Induced Pluripotent Stem Cells to Endoderm by Enhancing Survival Through Protein Kinase B Phosphorylation. <i>Hepatology</i> , 2021, 74, 2102-2117.	7.3	5
17	ER Disposal Pathways in Chronic Liver Disease: Protective, Pathogenic, and Potential Therapeutic Targets. <i>Frontiers in Molecular Biosciences</i> , 2021, 8, 804097.	3.5	5
18	Effects of acute diuresis stress on egr-1 (zif268) mRNA levels in brain regions associated with motivated behavior. <i>Brain Research Bulletin</i> , 2010, 81, 114-119.	3.0	4

#	ARTICLE	IF	CITATIONS
19	The Association of Hispanic Ethnicity with Nonalcoholic Fatty Liver Disease in Polycystic Ovary Syndrome. , 2018, 1, 24-33.		3