
Amir Mehrkar

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/3400194/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Severity of Severe Acute Respiratory System Coronavirus 2 (SARS-CoV-2) Alpha Variant (B.1.1.7) in England. Clinical Infectious Diseases, 2022, 75, e1120-e1127.	2.9	71
2	OpenSAFELY NHS Service Restoration Observatory 1: primary care clinical activity in England during the first wave of COVID-19. British Journal of General Practice, 2022, 72, e63-e74.	0.7	22
3	Trends and clinical characteristics of COVID-19 vaccine recipients: a federated analysis of 57.9 million patients' primary care records <i>in situ</i> using OpenSAFELY. British Journal of General Practice, 2022, 72, e51-e62.	0.7	75
4	Mortality among Care Home Residents in England during the first and second waves of the COVID-19 pandemic: an observational study of 4.3 million adults over the age of 65. Lancet Regional Health - Europe, The, 2022, 14, 100295.	3.0	38
5	Overall and cause-specific hospitalisation and death after COVID-19 hospitalisation in England: A cohort study using linked primary care, secondary care, and death registration data in the OpenSAFELY platform. PLoS Medicine, 2022, 19, e1003871.	3.9	39
6	Potentially inappropriate prescribing of DOACs to people with mechanical heart valves: A federated analysis of 57.9 million patients' primary care records in situ using OpenSAFELY. Thrombosis Research, 2022, 211, 150-153.	0.8	6
7	Association between oral anticoagulants and COVID-19-related outcomes: a population-based cohort study. British Journal of General Practice, 2022, 72, e456-e463.	0.7	3
8	Comparison of methods for predicting COVID-19-related death in the general population using the OpenSAFELY platform. Diagnostic and Prognostic Research, 2022, 6, 6.	0.8	2
9	Effect of pre-exposure use of hydroxychloroquine on COVID-19 mortality: a population-based cohort study in patients with rheumatoid arthritis or systemic lupus erythematosus using the OpenSAFELY platform. Lancet Rheumatology, The, 2021, 3, e19-e27.	2.2	49
10	Use of non-steroidal anti-inflammatory drugs and risk of death from COVID-19: an OpenSAFELY cohort analysis based on two cohorts. Annals of the Rheumatic Diseases, 2021, 80, 943-951.	0.5	66
11	HIV infection and COVID-19 death: a population-based cohort analysis of UK primary care data and linked national death registrations within the OpenSAFELY platform. Lancet HIV,the, 2021, 8, e24-e32.	2.1	340
12	Association between living with children and outcomes from covid-19: OpenSAFELY cohort study of 12 million adults in England. BMJ, The, 2021, 372, n628.	3.0	56
13	Hydroxychloroquine treatment does not reduce COVID-19 mortality; underdosing to the wrong patients? – Authors' reply. Lancet Rheumatology, The, 2021, 3, e172-e173.	2.2	1
14	Case fatality risk of the SARS-CoV-2 variant of concern B.1.1.7 in England, 16 November to 5 February. Eurosurveillance, 2021, 26, .	3.9	156
15	Identifying Care Home Residents in Electronic Health Records - An OpenSAFELY Short Data Report. Wellcome Open Research, 2021, 6, 90.	0.9	18
16	Ethnic differences in SARS-CoV-2 infection and COVID-19-related hospitalisation, intensive care unit admission, and death in 17 million adults in England: an observational cohort study using the OpenSAFELY platform. Lancet, The, 2021, 397, 1711-1724.	6.3	332
17	Clinical coding of long COVID in English primary care: a federated analysis of 58 million patient records <i>in situ</i> using OpenSAFELY. British Journal of General Practice, 2021, 71, e806-e814.	0.7	74
18	Risks of covid-19 hospital admission and death for people with learning disability: population based cohort study using the OpenSAFELY platform. BMJ, The, 2021, 374, n1592.	3.0	70

#	Article	IF	CITATIONS
19	Factors associated with deaths due to COVID-19 versus other causes: population-based cohort analysis of UK primary care data and linked national death registrations within the OpenSAFELY platform. Lancet Regional Health - Europe, The, 2021, 6, 100109.	3.0	121
20	Association between warfarin and COVID-19-related outcomes compared with direct oral anticoagulants: population-based cohort study. Journal of Hematology and Oncology, 2021, 14, 172.	6.9	8
21	OpenSAFELY: impact of national guidance on switching anticoagulant therapy during COVID-19 pandemic. Open Heart, 2021, 8, e001784.	0.9	17
22	Factors associated with COVID-19-related death using OpenSAFELY. Nature, 2020, 584, 430-436.	13.7	4,674
23	Risk of COVID-19-related death among patients with chronic obstructive pulmonary disease or asthma prescribed inhaled corticosteroids: an observational cohort study using the OpenSAFELY platform. Lancet Respiratory Medicine,the, 2020, 8, 1106-1120.	5.2	211