
Yongheng Li

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/3401608/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Yap promotes hepatocellular carcinoma metastasis and mobilization via governing cofilin/F-actin/lamellipodium axis by regulation of JNK/Bnip3/SERCA/CaMKII pathways. Redox Biology, 2018, 14, 59-71.	9.0	193
2	Combination of radiotherapy and suppression of Tregs enhances abscopal antitumor effect and inhibits metastasis in rectal cancer. , 2020, 8, e000826.		74
3	Expert consensus on multidisciplinary therapy of colorectal cancer with lung metastases (2019) Tj ETQq1 1 0.7	84314 rgB 17.0	T /Qyerlock 1
4	Chinese Guidelines on the Diagnosis and Treatment of Melanoma (2015 Edition). Chinese Clinical Oncology, 2016, 5, 57-57.	1.2	46
5	Chinese Guidelines on the Diagnosis and Treatment of Melanoma (2015 Edition). Annals of Translational Medicine, 2015, 3, 322.	1.7	32
6	Smart Tumorâ€Cellâ€Derived Microparticles Provide Onâ€Demand Photosensitizer Synthesis and Hypoxia Relief for Photodynamic Therapy. Angewandte Chemie - International Edition, 2021, 60, 25365-25371.	13.8	26
7	Prognostic Factors and Optimal Response Interval for Stereotactic Body Radiotherapy in Patients With Lung Oligometastases or Oligoprogression From Colorectal Cancer. Frontiers in Oncology, 2019, 9, 1080.	2.8	20
8	MRI radiomics independent of clinical baseline characteristics and neoadjuvant treatment modalities predicts response to neoadjuvant therapy in rectal cancer. British Journal of Cancer, 2022, 127, 249-257.	6.4	20
9	Simultaneous Integrated Boost Intensity-Modulated Radiation Therapy Can Benefit the Locally Advanced Rectal Cancer Patients With Clinically Positive Lateral Pelvic Lymph Node. Frontiers in Oncology, 2020, 10, 627572.	2.8	7
10	Establishment of prognostic models for adenocarcinoma of oesophagogastric junction patients with neoadjuvant chemoradiotherapy: a real-world study. Radiation Oncology, 2022, 17, 45.	2.7	7
11	Excluding the ischiorectal fossa irradiation during neoadjuvant chemoradiotherapy with intensity-modulated radiotherapy followed by abdominoperineal resection decreases perineal complications in patients with lower rectal cancer. Radiation Oncology, 2019, 14, 138.	2.7	5
12	Clinical effectiveness of recombinant adenovirus-p53 combined with radiotherapy in advanced soft tissue sarcoma: A report of 37 cases Journal of Clinical Oncology, 2014, 32, e21514-e21514.	1.6	5
13	The feasibility and efficiency of wait and see policy for patients with complete clinical response following neoadjuvant therapy in rectal cancer: A prospective cohort study from China Journal of Clinical Oncology, 2017, 35, 3610-3610.	1.6	5
14	Clinical antiangiogenic effect of recombinant adenovirus-p53 combined with hyperthermia for advanced cancer. Chinese Journal of Cancer Research: Official Journal of China Anti-Cancer Association, Beijing Institute for Cancer Research, 2013, 25, 749-55.	2.2	5
15	Prognostic significance of <scp>albumin–bilirubin</scp> score in patients with unresectable hepatocellular carcinoma undergoing combined immunotherapy and radiotherapy. Journal of Medical Imaging and Radiation Oncology, 2022, 66, 662-670.	1.8	5
16	Surgery may not benefit patients with locally advanced rectal cancer who achieved clinical complete response following neoadjuvant chemoradiotherapy. Asian Journal of Surgery, 2022, 45, 97-104.	0.4	4
17	Effect of Simultaneous Integrated Boost Intensity Modulated Radiation Therapy (SIB-IMRT) and Non-Operative Strategy on Outcomes of Distal Rectal Cancer Patients with Clinically Positive Lateral Pelvic Lymph Node. Cancer Management and Research, 2021, Volume 13, 537-546.	1.9	4
18	ls Elective Inguinal or External Iliac Irradiation During Neoadjuvant (Chemo)radiotherapy Necessary for Locally Advanced Lower Rectal Cancer With Anal Sphincter Invasion?. Practical Radiation Oncology, 2022, 12, 125-134.	2.1	4

YONGHENG LI

#	Article	IF	CITATIONS
19	Stereotactic body radiotherapy prolongs the progressionâ€free survival and delays the change of systemic therapy regimen in patients with lung oligoprogressive metastatic colorectal cancer. Asia-Pacific Journal of Clinical Oncology, 2022, 18, .	1.1	2
20	One to Two Cycles of Consolidation Chemotherapy With Capecitabine After Neoadjuvant Chemoradiotherapy Does Not Benefit Low-Risk Patients With Locally Advanced Middle-Low Rectal Cancer. Frontiers in Oncology, 2021, 11, 695726.	2.8	2
21	Smart Tumorâ€Cellâ€Derived Microparticles Provide Onâ€Demand Photosensitizer Synthesis and Hypoxia Relief for Photodynamic Therapy. Angewandte Chemie, 0, , .	2.0	2
22	Hyperfractionation versus Conventional Fractionation of Preoperative Intensity-Modulated Radiotherapy with Oral Capecitabine in Locally Advanced Mid-Low Rectal Cancer: A Propensity Score Matching Study. Journal of Oncology, 2022, 2022, 1-10.	1.3	2
23	Compliance and safety of neoadjuvant intensity modulated radiotherapy (IMRT) with concurrent capecitabine for locally advanced rectal cancer: Updated results from a phase II trial (ChiCTR-TNC-10001094) Journal of Clinical Oncology, 2014, 32, 3598-3598.	1.6	Ο