Mingxun Chen

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/3411498/publications.pdf

Version: 2024-02-01

26 papers

929 citations

471509 17 h-index 25 g-index

26 all docs

 $\begin{array}{c} 26 \\ \text{docs citations} \end{array}$

times ranked

26

1008 citing authors

#	Article	IF	CITATIONS
1	Overexpression of $\langle i \rangle$ BnaAGL11 $\langle i \rangle$, a MADS-Box Transcription Factor, Regulates Leaf Morphogenesis and Senescence in $\langle i \rangle$ Brassica napus $\langle i \rangle$. Journal of Agricultural and Food Chemistry, 2022, 70, 3420-3434.	5.2	9
2	Novel parameters characterizing size distribution of A and B starch granules in the gluten network: Effects on dough stability in bread wheat. Carbohydrate Polymers, 2021, 257, 117623.	10.2	23
3	Linum usitatissimum FAD2A and FAD3A enhance seed polyunsaturated fatty acid accumulation and seedling cold tolerance in Arabidopsis thaliana. Plant Science, 2021, 311, 111014.	3.6	17
4	Transcriptome analysis reveals key genes in response to salinity stress during seed germination in Setaria italica. Environmental and Experimental Botany, 2021, 191, 104604.	4.2	8
5	Genome-wide association study reveals a patatin-like lipase relating to the reduction of seed oil content in Brassica napus. BMC Plant Biology, 2021, 21, 6.	3.6	11
6	The possible role of BnaA10.SOI.a in seed fatty acid biosynthesis of rapeseed. Plant Breeding, 2020, 139, 167-175.	1.9	0
7	Melatonin Represses Oil and Anthocyanin Accumulation in Seeds. Plant Physiology, 2020, 183, 898-914.	4.8	25
8	Genome-Wide Identification of Direct Targets of the TTG1–bHLH–MYB Complex in Regulating Trichome Formation and Flavonoid Accumulation in Arabidopsis Thaliana. International Journal of Molecular Sciences, 2019, 20, 5014.	4.1	25
9	Satellite Observations of the Recovery of Forests and Grasslands in Western China. Journal of Geophysical Research G: Biogeosciences, 2019, 124, 1905-1922.	3.0	20
10	Drought-Induced Carbon and Water Use Efficiency Responses in Dryland Vegetation of Northern China. Frontiers in Plant Science, 2019, 10, 224.	3.6	17
11	Functional characterization of a heterologously expressed Brassica napus WRKY41-1 transcription factor in regulating anthocyanin biosynthesis in Arabidopsis thaliana. Plant Science, 2018, 268, 47-53.	3.6	90
12	<i>TRANSPARENT TESTA 4</i> i>â€mediated flavonoids negatively affect embryonic fatty acid biosynthesis in <i>Arabidopsis</i> Plant, Cell and Environment, 2018, 41, 2773-2790.	5.7	26
13	Genome-wide identification of GLABRA3 downstream genes for anthocyanin biosynthesis and trichome formation in Arabidopsis. Biochemical and Biophysical Research Communications, 2017, 485, 360-365.	2.1	11
14	Functional characterization of Brassica napus DNA topoisomerase $\hat{ll}\pm -1$ and its effect on flowering time when expressed in Arabidopsis thaliana. Biochemical and Biophysical Research Communications, 2017, 486, 124-129.	2.1	2
15	The effect of BnTT8 on accumulation of seed storage reserves and tolerance to abiotic stresses during Arabidopsis seedling establishment. Plant Growth Regulation, 2017, 82, 271-280.	3.4	8
16	MYB89 Transcription Factor Represses Seed Oil Accumulation. Plant Physiology, 2017, 173, 1211-1225.	4.8	87
17	TRANSPARENT TESTA GLABRA 1 ubiquitously regulates plant growth and development from Arabidopsis to foxtail millet (Setaria italica). Plant Science, 2017, 254, 60-69.	3.6	22
18	MYB76 Inhibits Seed Fatty Acid Accumulation in Arabidopsis. Frontiers in Plant Science, 2017, 8, 226.	3.6	30

#	Article	IF	CITATION
19	MYC2, MYC3, and MYC4 function redundantly in seed storage protein accumulation in Arabidopsis. Plant Physiology and Biochemistry, 2016, 108, 63-70.	5.8	40
20	Comparison of vitality between seedlings germinated from black-coated and yellow-coated seeds of a turnip rape (Brassica rapa L.) subjected to NaCl and CdCl2 stresses. Plant Growth Regulation, 2015, 76, 61-70.	3.4	11
21	<i>TRANSPARENT TESTA GLABRA1</i> Regulates the Accumulation of Seed Storage Reserves in Arabidopsis. Plant Physiology, 2015, 169, 391-402.	4.8	71
22	<i><scp>TRANSPARENT TESTA</scp>2</i> regulates embryonic fatty acid biosynthesis by targeting <i><scp>FUSCA</scp>3</i> during the early developmental stage of <scp>A</scp> rabidopsis seeds. Plant Journal, 2014, 77, 757-769.	5.7	63
23	TRANSPARENT TESTA8 Inhibits Seed Fatty Acid Accumulation by Targeting Several Seed Development Regulators in Arabidopsis. Plant Physiology, 2014, 165, 905-916.	4.8	78
24	Removal of DELLA repression promotes leaf senescence in Arabidopsis. Plant Science, 2014, 219-220, 26-34.	3.6	63
25	The Effect of <i>TRANSPARENT TESTA2</i> on Seed Fatty Acid Biosynthesis and Tolerance to Environmental Stresses during Young Seedling Establishment in Arabidopsis Â. Plant Physiology, 2012, 160, 1023-1036.	4.8	79
26	<i>Seed Fatty Acid Reducer</i> acts downstream of gibberellin signalling pathway to lower seed fatty acid storage in <i>Arabidopsis</i> . Plant, Cell and Environment, 2012, 35, 2155-2169.	5 . 7	93