

List of Publications by Citations

Source: <https://exaly.com/author-pdf/3412860/karel-kostev-publications-by-citations.pdf>
Version: 2024-04-20

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.
The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

406 papers	3,974 citations	29 h-index	45 g-index
506 ext. papers	5,329 ext. citations	3.4 avg, IF	6.39 L-index

#	Paper	IF	Citations
406	Validity and representativeness of the "Disease Analyzer" patient database for use in pharmacoepidemiological and pharmacoeconomic studies. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2009 , 47, 617-26	2	193
405	Basic characteristics and representativeness of the German Disease Analyzer database. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2018 , 56, 459-466	2	136
404	GRAND: the German retrospective cohort analysis on compliance and persistence and the associated risk of fractures in osteoporotic women treated with oral bisphosphonates. <i>Osteoporosis International</i> , 2012 , 23, 223-31	5.3	96
403	Prevalence, utilization, and costs of antiepileptic drugs for epilepsy in Germany--a nationwide population-based study in children and adults. <i>Journal of Neurology</i> , 2012 , 259, 2376-84	5.5	93
402	Treatment persistence, hypoglycaemia and clinical outcomes in type 2 diabetes patients with dipeptidyl peptidase-4 inhibitors and sulphonylureas: a primary care database analysis. <i>Diabetes, Obesity and Metabolism</i> , 2013 , 15, 55-61	6.7	69
401	No reduced risk of overall, colorectal, lung, breast, and prostate cancer with metformin therapy in diabetic patients: database analyses from Germany and the UK. <i>Pharmacoepidemiology and Drug Safety</i> , 2015 , 24, 865-74	2.6	69
400	Persistence in patients with breast cancer treated with tamoxifen or aromatase inhibitors: a retrospective database analysis. <i>Breast Cancer Research and Treatment</i> , 2013 , 138, 185-91	4.4	69
399	Amputation rate and risk factors in type 2 patients with diabetic foot syndrome under real-life conditions in Germany. <i>Primary Care Diabetes</i> , 2012 , 6, 241-6	2.4	62
398	Persistence with antihypertensive treatments: results of a 3-year follow-up cohort study. <i>European Journal of Clinical Pharmacology</i> , 2007 , 63, 1055-61	2.8	58
397	Risk factors for dementia diagnosis in German primary care practices. <i>International Psychogeriatrics</i> , 2016 , 28, 1059-65	3.4	56
396	Diabetic retinopathy at diagnosis of type 2 diabetes in the UK: a database analysis. <i>Diabetologia</i> , 2013 , 56, 109-11	10.3	55
395	Persistence and compliance of medications used in the treatment of osteoporosis--analysis using a large scale, representative, longitudinal German database. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2012 , 50, 315-22	2	54
394	Isopropanolic black cohosh extract and recurrence-free survival after breast cancer. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2007 , 45, 143-54	2	53
393	Prevalence of diabetic foot syndrome and its risk factors in the UK. <i>Journal of Wound Care</i> , 2010 , 19, 333-7	2.2	51
392	Basic characteristics and representativeness of the German Disease Analyzer database?. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2018 , 56, 459-466	2	50
391	Nonadherence to antiepileptic drugs in Germany: A retrospective, population-based study. <i>Neurology</i> , 2016 , 87, 466-72	6.5	46
390	Adherence to GLP-1 receptor agonist therapy administered by once-daily or once-weekly injection in patients with type 2 diabetes in Germany. <i>Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy</i> , 2016 , 9, 201-5	3.4	45

389	Use of dipeptidyl peptidase-4 inhibitors and risk of bone fracture in patients with type 2 diabetes in Germany-A retrospective analysis of real-world data. <i>Osteoporosis International</i> , 2017 , 28, 2421-2428	5.3	42
388	Impact of tobacco smoking on the risk of developing 25 different cancers in the UK: a retrospective study of 422,010 patients followed for up to 30 years. <i>Oncotarget</i> , 2018 , 9, 17420-17429	3.3	38
387	Are sulfonylurea and insulin therapies associated with a larger risk of cancer than metformin therapy? A retrospective database analysis. <i>Diabetes Care</i> , 2015 , 38, 59-65	14.6	36
386	Fracture risk in patients with newly diagnosed type 2 diabetes: a retrospective database analysis in primary care. <i>Journal of Diabetes and Its Complications</i> , 2015 , 29, 766-70	3.2	35
385	Impact of NAFLD on the Incidence of Cardiovascular Diseases in a Primary Care Population in Germany. <i>Digestive Diseases and Sciences</i> , 2020 , 65, 2112-2119	4	35
384	Prevalence of depression, anxiety and their risk factors in German women with breast cancer in general and gynecological practices. <i>Journal of Cancer Research and Clinical Oncology</i> , 2016 , 142, 447-52	4.9	33
383	Panic buying or good adherence? Increased pharmacy purchases of drugs from wholesalers in the last week prior to Covid-19 lockdown. <i>Journal of Psychiatric Research</i> , 2020 , 130, 19-21	5.2	33
382	Prevalence and risk factors of neuropathy in newly diagnosed type 2 diabetes in primary care practices: a retrospective database analysis in Germany and U.K. <i>Primary Care Diabetes</i> , 2014 , 8, 250-5	2.4	32
381	Antiemetic medications in children with presumed infectious gastroenteritis--pharmacoepidemiology in Europe and Northern America. <i>Journal of Pediatrics</i> , 2008 , 153, 659-62, 662.e1-3	3.6	32
380	Impact of the COVID-19 Pandemic on Cancer Diagnoses in General and Specialized Practices in Germany. <i>Cancers</i> , 2021 , 13,	6.6	32
379	Depression Risk in Patients with Rheumatoid Arthritis in the United Kingdom. <i>Rheumatology and Therapy</i> , 2017 , 4, 195-200	4.4	31
378	Nonalcoholic Fatty Liver Disease Increases the Risk of Anxiety and Depression. <i>Hepatology Communications</i> , 2020 , 4, 1293-1301	6	29
377	Switching the manufacturer of antiepileptic drugs is associated with higher risk of seizures: A nationwide study of prescription data in Germany. <i>Annals of Neurology</i> , 2018 , 84, 918-925	9.4	29
376	Treatment patterns and low-density lipoprotein cholesterol (LDL-C) goal attainment among patients receiving high- or moderate-intensity statins. <i>Clinical Research in Cardiology</i> , 2018 , 107, 380-388	6.1	28
375	The impact of depot medroxyprogesterone acetate on fracture risk: a case-control study from the UK. <i>Osteoporosis International</i> , 2017 , 28, 291-297	5.3	28
374	Impact of rotavirus vaccination in Germany: rotavirus surveillance, hospitalization, side effects and comparison of vaccines. <i>Pediatric Infectious Disease Journal</i> , 2014 , 33, e299-304	3.4	28
373	Real-world treatment patterns of sacubitril/valsartan: a longitudinal cohort study in Germany. <i>European Journal of Heart Failure</i> , 2019 , 21, 588-597	12.3	27
372	Predictors of hypoglycaemia in insulin-treated type 2 diabetes patients in primary care: a retrospective database analysis. <i>Primary Care Diabetes</i> , 2014 , 8, 127-31	2.4	27

371	Association Between the Use of Antihyperglycemic Drugs and Dementia Risk: A Case-Control Study. <i>Journal of Alzheimer's Disease</i> , 2018 , 66, 725-732	4.3	26
370	The impact of treatment compliance on fracture risk in women with breast cancer treated with aromatase inhibitors in the United Kingdom. <i>Breast Cancer Research and Treatment</i> , 2016 , 155, 151-7	4.4	25
369	Effect of the COVID-19 lockdown on disease recognition and utilisation of healthcare services in the older population in Germany: a cross-sectional study. <i>Age and Ageing</i> , 2021 , 50, 317-325	3	25
368	Incidence of depression and anxiety among women newly diagnosed with breast or genital organ cancer in Germany. <i>Psycho-Oncology</i> , 2017 , 26, 1535-1540	3.9	23
367	Antihypertensive treatment and risk of dementia: a retrospective database study. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2012 , 50, 195-201	2	23
366	Continuous treatment with antidementia drugs in Germany 2003-2013: a retrospective database analysis. <i>International Psychogeriatrics</i> , 2015 , 27, 1335-42	3.4	22
365	Medical Record Databases	224-243	22
364	Prevalence of depression in type 2 diabetes patients in German primary care practices. <i>Journal of Diabetes and Its Complications</i> , 2016 , 30, 432-7	3.2	21
363	Non-alcoholic fatty liver disease increases the risk of incident chronic kidney disease. <i>United European Gastroenterology Journal</i> , 2020 , 8, 942-948	5.3	21
362	Differences in persistency with teriparatide in patients with osteoporosis according to gender and health care provider. <i>Osteoporosis International</i> , 2014 , 25, 2721-8	5.3	20
361	Cost comparison of outpatient treatment with granulocyte colony-stimulating factors (G-CSF) in Germany. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2012 , 50, 281-9	2	20
360	Impact of metformin on metastases in patients with breast cancer and type 2 diabetes. <i>Journal of Diabetes and Its Complications</i> , 2016 , 30, 1056-9	3.2	20
359	Long-term use of benzodiazepines in older patients in Germany: a retrospective analysis. <i>Therapeutic Advances in Psychopharmacology</i> , 2017 , 7, 191-200	4.9	19
358	The tamoxifen paradox-influence of adjuvant tamoxifen on fracture risk in pre- and postmenopausal women with breast cancer. <i>Osteoporosis International</i> , 2018 , 29, 2557-2564	5.3	19
357	Discontinuation of treatment using anticholinergic medications in patients with urinary incontinence. <i>Obstetrics and Gynecology</i> , 2014 , 124, 794-800	4.9	19
356	Fracture risk in patients with type 2 diabetes under different antidiabetic treatment regimens: a retrospective database analysis in primary care. <i>Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy</i> , 2016 , 9, 17-23	3.4	19
355	Impact of prior breast cancer on mode of delivery and pregnancy-associated disorders: a retrospective analysis of subsequent pregnancy outcomes. <i>Journal of Cancer Research and Clinical Oncology</i> , 2017 , 143, 1069-1074	4.9	18
354	Association of dipeptidyl peptidase 4 inhibitors with risk of metastases in patients with type 2 diabetes and breast, prostate or digestive system cancer. <i>Journal of Diabetes and Its Complications</i> , 2017 , 31, 687-692	3.2	17

353	Sociodemographic disparities in administration of antiepileptic drugs to adults with epilepsy in Germany: a retrospective, database study of drug prescriptions. <i>CNS Drugs</i> , 2014 , 28, 753-9	6.7	17
352	Changes in time to insulin initiation in type 2 diabetes patients: a retrospective database analysis in Germany and UK (2005-2010). <i>Primary Care Diabetes</i> , 2013 , 7, 229-33	2.4	17
351	Prevalence of chronic diseases among older patients in German general practices. <i>GMS German Medical Science</i> , 2016 , 14, Doc03	3.2	17
350	Gender- and age-related treatment compliance in patients with osteoporosis in Germany. <i>Patient Preference and Adherence</i> , 2016 , 10, 2379-2385	2.4	17
349	Oral contraceptive use and fracture risk-a retrospective study of 12,970 women in the UK. <i>Osteoporosis International</i> , 2017 , 28, 2349-2355	5.3	16
348	Prevalence of and risk factors associated with cesarean section in Lebanon - A retrospective study based on a sample of 29,270 women. <i>Women and Birth</i> , 2017 , 30, e265-e271	3.3	16
347	Trends and patterns of hormonal contraceptive prescribing for adolescents in primary care in the U.K. <i>Journal of Family Planning and Reproductive Health Care</i> , 2015 , 41, 216-22		16
346	Adherence to antiepileptic drugs in children and adolescents: A retrospective study in primary care settings in Germany. <i>Epilepsy and Behavior</i> , 2017 , 75, 36-41	3.2	16
345	Glycemic control after initiating basal insulin therapy in patients with type 2 diabetes: a primary care database analysis. <i>Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy</i> , 2015 , 8, 45-8	3.4	16
344	Risk of venous thrombosis in users of hormonal contraceptives in German gynaecological practices: a patient database analysis. <i>Archives of Gynecology and Obstetrics</i> , 2014 , 289, 413-9	2.5	16
343	Prevalence of vulvovaginal candidiasis in gynecological practices in Germany: A retrospective study of 954,186 patients. <i>Current Medical Mycology</i> , 2018 , 4, 6-11	1.1	16
342	Induction of labor in breech presentations at term: a retrospective observational study. <i>Archives of Gynecology and Obstetrics</i> , 2016 , 293, 549-55	2.5	15
341	Treatment persistence after initiating basal insulin in type 2 diabetes patients: A primary care database analysis. <i>Primary Care Diabetes</i> , 2015 , 9, 377-84	2.4	15
340	Depression risk in patients with late-onset rheumatoid arthritis in Germany. <i>Quality of Life Research</i> , 2017 , 26, 437-443	3.7	15
339	Rotavirus vaccination in Germany: analysis of nationwide surveillance data 2006 to 2010. <i>Pediatric Infectious Disease Journal</i> , 2011 , 30, e244-7	3.4	15
338	Changes in HbA1c, body weight, and systolic blood pressure in type 2 diabetes patients initiating dapagliflozin therapy: a primary care database study. <i>Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy</i> , 2016 , 9, 337-345	3.4	15
337	Treatment of depression in cancer and non-cancer patients in German neuropsychiatric practices. <i>Psycho-Oncology</i> , 2016 , 25, 1324-1328	3.9	15
336	Impact of disease management programs on women with breast cancer in Germany. <i>Breast Cancer Research and Treatment</i> , 2015 , 153, 391-5	4.4	14

335	Multimorbidity and polypharmacy among elderly people followed in general practices in Germany. <i>European Journal of Internal Medicine</i> , 2018 , 55, 66-68	3.9	14
334	Lower incidence of recorded cardiovascular outcomes in patients with type 2 diabetes using insulin aspart vs. those on human regular insulin: observational evidence from general practices. <i>Diabetes, Obesity and Metabolism</i> , 2013 , 15, 358-63	6.7	14
333	Characteristics of diabetic patients visiting a podiatry practice in Germany. <i>Journal of Wound Care</i> , 2010 , 19, 140, 142, 144 passim	2.2	14
332	Depression risk in patients with coronary heart disease in Germany. <i>World Journal of Cardiology</i> , 2016 , 8, 547-552	2.1	14
331	Association Between Migraine Headaches and Dementia in More than 7,400 Patients Followed in General Practices in the United Kingdom. <i>Journal of Alzheimer's Disease</i> , 2019 , 71, 353-360	4.3	13
330	Real-life treatment patterns, compliance, persistence, and medication costs in patients with hypertension in Germany. <i>Journal of Medical Economics</i> , 2012 , 15, 155-65	2.4	13
329	Relation of the first hypertension-associated event with medication, compliance and persistence in naïve hypertensive patients after initiating monotherapy. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2010 , 48, 173-83	2	13
328	Association Between Hearing and Vision Impairment and Risk of Dementia: Results of a Case-Control Study Based on Secondary Data. <i>Frontiers in Aging Neuroscience</i> , 2019 , 11, 363	5.3	13
327	Association between chronic somatic conditions and depression in children and adolescents: A retrospective study of 13,326 patients. <i>Journal of Affective Disorders</i> , 2019 , 245, 697-701	6.6	13
326	Compliance with vaccination against tick-borne encephalitis virus in Germany. <i>Clinical Microbiology and Infection</i> , 2017 , 23, 460-463	9.5	12
325	Association between induced abortion, spontaneous abortion, and infertility respectively and the risk of psychiatric disorders in 57,770 women followed in gynecological practices in Germany. <i>Journal of Affective Disorders</i> , 2019 , 251, 107-113	6.6	12
324	Discontinuation rates of menopausal hormone therapy among postmenopausal women in the post-WHI study era. <i>Climacteric</i> , 2015 , 18, 737-42	3.1	12
323	The use of proton pump inhibitors is positively associated with osteoporosis in postmenopausal women in Germany. <i>Climacteric</i> , 2016 , 19, 478-81	3.1	12
322	Persistence with bisphosphonates in patients with metastatic breast cancer: a retrospective database analysis. <i>Journal of Cancer Research and Clinical Oncology</i> , 2013 , 139, 1149-55	4.9	12
321	Risk of psychiatric and neurological diseases in patients with workplace mobbing experience in Germany: a retrospective database analysis. <i>GMS German Medical Science</i> , 2014 , 12, Doc10	3.2	12
320	Do patients with lactose intolerance exhibit more frequent comorbidities than patients without lactose intolerance? An analysis of routine data from German medical practices. <i>Annals of Gastroenterology</i> , 2016 , 29, 174-9	2.2	12
319	Prevalence of pain medication prescriptions in France, Germany, and the UK - a cross-sectional study including 4,270,142 patients. <i>Postgraduate Medicine</i> , 2018 , 130, 32-36	3.7	11
318	Changes in Type 2 Diabetes Mellitus Patients in German Primary Care Prior to (2006) and After (2010, 2014) Launch of New Drugs. <i>Journal of Diabetes Science and Technology</i> , 2016 , 10, 414-20	4.1	11

317	Psoriasis risk in patients with type 2 diabetes in German primary care practices. <i>Primary Care Diabetes</i> , 2017 , 11, 52-56	2.4	11
316	Prevalence of depression, anxiety, and adjustment disorders in women with spontaneous abortion in Germany - A retrospective cohort study. <i>Psychiatry Research</i> , 2017 , 258, 382-386	9.9	11
315	Drug utilization patterns and reported health status in ethnic German migrants (Aussiedler) in Germany: a cross-sectional study. <i>BMC Public Health</i> , 2011 , 11, 509	4.1	11
314	The risk of developing depression when suffering from neurological diseases. <i>GMS German Medical Science</i> , 2013 , 11, Doc02	3.2	11
313	Adherence in tamoxifen therapy after conversion to a rebate pharmaceutical in breast cancer patients in Germany. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2013 , 51, 969-75	2	11
312	Change in glycated haemoglobin levels after initiating second-line therapy in type 2 diabetes: a primary care database study. <i>Diabetes, Obesity and Metabolism</i> , 2016 , 18, 840-3	6.7	11
311	Depression risk in female patients with osteoporosis in primary care practices in Germany. <i>Osteoporosis International</i> , 2016 , 27, 2739-2744	5.3	11
310	Hip fracture risk in patients with dementia in German primary care practices. <i>Dementia</i> , 2017 , 16, 853-864		10
309	Risk factors and outcomes in "well-selected" vaginal breech deliveries: a retrospective observational study. <i>Journal of Perinatal Medicine</i> , 2017 , 45, 291-297	2.7	10
308	Risk Factors for Mild Cognitive Impairment in German Primary Care Practices. <i>Journal of Alzheimer's Disease</i> , 2017 , 56, 379-384	4.3	10
307	Early discontinuation and related treatment costs after initiation of Basal insulin in type 2 diabetes patients: a German primary care database analysis. <i>Journal of Diabetes Science and Technology</i> , 2015 , 9, 644-50	4.1	10
306	Diagnoses of Dizziness- and Vertigo-related Disorders in ENT Practices in Germany. <i>Otology and Neurotology</i> , 2018 , 39, 474-480	2.6	10
305	Prevalence of medically treated children with ADHD and type 1 diabetes in Germany - Analysis of two representative databases. <i>Journal of Pediatric Endocrinology and Metabolism</i> , 2016 , 29, 1293-1297	1.6	10
304	Increased treatment persistence and its determinants in women with osteoporosis with prior fracture compared to those without fracture. <i>Osteoporosis International</i> , 2016 , 27, 963-969	5.3	10
303	Risk of stillbirth in pregnant women with obesity in the United Kingdom. <i>Obesity Research and Clinical Practice</i> , 2016 , 10, 574-579	5.4	10
302	Association of laryngeal cancer with vocal cord leukoplakia and associated risk factors in 1,184 patients diagnosed in otorhinolaryngology practices in Germany. <i>Molecular and Clinical Oncology</i> , 2018 , 8, 689-693	1.6	10
301	Impact of Osteoporosis on the Risk of Dementia in Almost 60,000 Patients Followed in General Practices in Germany. <i>Journal of Alzheimer's Disease</i> , 2018 , 65, 401-407	4.3	10
300	Prescription Patterns and the Cost of Antihyperglycemic Drugs in Patients With Type 2 Diabetes Mellitus in Germany. <i>Journal of Diabetes Science and Technology</i> , 2017 , 11, 123-127	4.1	10

299	Glycemic durability of monotherapy for diabetes. <i>New England Journal of Medicine</i> , 2007 , 356, 1378-9; author reply 1380	59.2	10
298	Risk factors for epistaxis in patients followed in general practices in Germany. <i>Rhinology</i> , 2017 , 55, 312-318		10
297	Impact of the coronavirus 2019 (COVID-19) pandemic on anxiety diagnosis in general practices in Germany. <i>Journal of Psychiatric Research</i> , 2021 , 143, 528-533	5.2	10
296	Comorbid disorders associated with asthma in children in Germany - National analysis of pediatric primary care data. <i>Pediatric Allergy and Immunology</i> , 2016 , 27, 861-866	4.2	10
295	Incidence of newly diagnosed diabetes after Covid-19.. <i>Diabetologia</i> , 2022 , 65, 949	10.3	10
294	Impact of Disease Management Programs on HbA1c Values in Type 2 Diabetes Patients in Germany. <i>Journal of Diabetes Science and Technology</i> , 2017 , 11, 117-122	4.1	9
293	Association between Anti-Dementia Treatment Persistence and Daily Dosage of the First Prescription: A Retrospective Analysis in Neuropsychiatric Practices in Germany. <i>Journal of Alzheimer's Disease</i> , 2017 , 58, 37-44	4.3	9
292	Effects of selected antidiabetics on weight loss--a retrospective database analysis. <i>Primary Care Diabetes</i> , 2015 , 9, 74-7	2.4	9
291	Risk of depression, anxiety, and adjustment disorders in women with a suspected but unconfirmed diagnosis of breast or genital organ cancer in Germany. <i>Cancer Causes and Control</i> , 2017 , 28, 1021-1026	2.8	9
290	Which adverse effects influence the dropout rate in selective serotonin reuptake inhibitor (SSRI) treatment? Results for 50,824 patients. <i>GMS German Medical Science</i> , 2014 , 12, Doc15	3.2	9
289	Persistence with antipsychotics in dementia patients in Germany?. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2016 , 54, 835-840	2	9
288	Incidence of epilepsy and associated factors in elderly patients in Germany. <i>Epilepsy and Behavior</i> , 2019 , 90, 107-111	3.2	9
287	Impact of dronedarone on the risk of myocardial infarction and stroke in atrial fibrillation patients followed in general practices in Germany. <i>International Journal of Cardiology</i> , 2019 , 278, 126-132	3.2	9
286	Association Between Low-Density Lipoprotein Cholesterol Levels, Statin Use, and Dementia in Patients followed in German General Practices. <i>Journal of Alzheimer's Disease</i> , 2021 , 79, 37-46	4.3	9
285	Association between sleep disorders and the presence of breast cancer metastases in gynecological practices in Germany: a case-control study of 11,412 women. <i>Breast Cancer Research and Treatment</i> , 2018 , 171, 443-448	4.4	9
284	Persistence with antiepileptic drugs in epilepsy patients treated in neurological practices in Germany. <i>Epilepsy and Behavior</i> , 2017 , 73, 204-207	3.2	8
283	The Relationship Between the Use of Antihypertensive Drugs and the Incidence of Dementia in General Practices in Germany. <i>Journal of Alzheimer's Disease</i> , 2019 , 70, 91-97	4.3	8
282	Cancer risk in stroke survivors followed for up to 10 years in general practices in Germany. <i>Journal of Cancer Research and Clinical Oncology</i> , 2019 , 145, 1013-1020	4.9	8

281	Incidence of fractures in patients with type 1 diabetes mellitus-a retrospective study with 4420 patients. <i>Osteoporosis International</i> , 2020 , 31, 1315-1322	5.3	8
280	Increased depression symptom score in newly diagnosed type 2 diabetes patients. <i>Psychiatry Research</i> , 2018 , 261, 259-263	9.9	8
279	Second-line therapy with biological drugs in rheumatoid arthritis patients in German rheumatologist practices: a retrospective database analysis. <i>Rheumatology International</i> , 2016 , 36, 1113-8	3.6	8
278	Impact of attention deficit hyperactivity disorder therapy on fracture risk in children treated in German pediatric practices. <i>Osteoporosis International</i> , 2017 , 28, 1265-1269	5.3	8
277	Longitudinal change in HbA1c after insulin initiation in primary care patients with type 2 diabetes: a database analysis in UK and Germany. <i>Primary Care Diabetes</i> , 2012 , 6, 47-52	2.4	8
276	Tumor Incidence in Patients with Non-Alcoholic Fatty Liver Disease. <i>Deutsches A&#x0308;rztblatt International</i> , 2020 , 117, 719-724	2.5	8
275	Treatment persistence evaluation of tamoxifen and aromatase inhibitors in breast cancer patients in early and late stage disease. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2014 , 52, 933-9	2	8
274	Persistence with antidepressant drugs in patients with dementia: a retrospective database analysis. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2016 , 54, 323-9	2	8
273	Prescription patterns and drug costs in German patients with dementia in nursing homes and home-care settings?. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2017 , 55, 9-15	2	8
272	Prescription rates of cardiovascular and diabetes therapies prior to and during the COVID-19 lockdown in Germany. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2020 , 58, 475-481	2	8
271	Antiherpetic medication and incident dementia: Observational cohort studies in four countries. <i>European Journal of Neurology</i> , 2021 , 28, 1840-1848	6	8
270	Heart failure is associated with an increased incidence of cancer diagnoses. <i>ESC Heart Failure</i> , 2021 , 8, 3628-3633	3.7	8
269	Risk Factors for Tinnitus in 37,692 Patients Followed in General Practices in Germany. <i>Otology and Neurotology</i> , 2019 , 40, 436-440	2.6	8
268	Trends in valproate use in patients in Germany between 2009 and 2017. <i>Epilepsy and Behavior</i> , 2019 , 92, 26-30	3.2	8
267	Time to pregnancy in subfertile women in German gynecological practices: analysis of a representative cohort of more than 60,000 patients. <i>Archives of Gynecology and Obstetrics</i> , 2015 , 291, 657-62	2.5	7
266	Persistency with estrogen replacement therapy among hysterectomized women after the Women's Health Initiative study. <i>Climacteric</i> , 2015 , 18, 826-34	3.1	7
265	Reduced antibiotic use after initial treatment of acute respiratory infections with phytopharmaceuticals- a retrospective cohort study. <i>Postgraduate Medicine</i> , 2020 , 132, 412-418	3.7	7
264	20-year risk of breast cancer recurrence. <i>Breast Cancer Research and Treatment</i> , 2018 , 168, 765-766	4.4	7

263	Caesarean section and its impact on fertility and time to a subsequent pregnancy in Germany: a database analysis in gynecological practices. <i>Archives of Gynecology and Obstetrics</i> , 2016 , 294, 1005-1010 ^{2.5}	7
262	Age-related differences in persistence in women with breast cancer treated with tamoxifen or aromatase inhibitors in Germany. <i>Journal of Geriatric Oncology</i> , 2016 , 7, 169-75	3.6 7
261	Incidence of fractures in young women with breast cancer - a retrospective cohort study. <i>Journal of Bone Oncology</i> , 2019 , 18, 100254	4.5 7
260	Prevalence and treatment profile of patients with grass pollen and house dust mite allergy. <i>JDDG - Journal of the German Society of Dermatology</i> , 2013 , 11, 653-61	1.2 7
259	Use of antidiabetic agents in the treatment of gestational diabetes mellitus in Germany, 2008-2012. <i>Journal of Obstetrics and Gynaecology Research</i> , 2014 , 40, 1592-7	1.9 7
258	Risk of breast cancer in patients on long-acting insulin analogues in comparison with those on human insulin. <i>Diabetologia</i> , 2012 , 55, 1554-5	10.3 7
257	Unlicensed use of metformin in children and adolescents in Germany and France. <i>British Journal of Clinical Pharmacology</i> , 2012 , 73, 307-8	3.8 7
256	Different injection frequencies of basal insulins in type 2 diabetes patients under real-life conditions: a retrospective database analysis. <i>Journal of Diabetes Science and Technology</i> , 2013 , 7, 1354-8 ^{4.1}	7
255	Potential delay in the diagnosis of vulvar cancer and associated risk factors in women treated in German gynecological practices. <i>Oncotarget</i> , 2018 , 9, 8725-8730	3.3 7
254	Fewer non-COVID-19 respiratory tract infections and gastrointestinal infections during the COVID-19 pandemic. <i>Journal of Medical Virology</i> , 2022 , 94, 298-302	19.7 7
253	Predictors of early discontinuation of basal insulin therapy in type 2 diabetes in primary care. <i>Primary Care Diabetes</i> , 2016 , 10, 142-7	2.4 6
252	Diabetes treatment in people with type 2 diabetes and schizophrenia: Retrospective primary care database analyses. <i>Primary Care Diabetes</i> , 2016 , 10, 36-40	2.4 6
251	Analysis of the Effects of Selective Serotonin (and Noradrenaline) Reuptake Inhibitors on the Risk of Dementia in Patients with Depression. <i>Journal of Alzheimer's Disease</i> , 2019 , 69, 577-583	4.3 6
250	Persistence with opioid treatment in Germany in patients suffering from chronic non-malignant or cancer pain. <i>Current Medical Research and Opinion</i> , 2015 , 31, 1157-63	2.5 6
249	Association between thyroid gland diseases and breast cancer: a case-control study. <i>Breast Cancer Research and Treatment</i> , 2020 , 182, 207-213	4.4 6
248	Urinary and fecal incontinence in stroke survivors followed in general practice: A retrospective cohort study. <i>Annals of Physical and Rehabilitation Medicine</i> , 2020 , 63, 488-494	3.8 6
247	Drug treatment for patients with bipolar disorders in psychiatric practices in Germany in 2009 and 2018. <i>Psychiatry Research</i> , 2020 , 289, 112965	9.9 6
246	Changes in patient characteristics, glucose lowering treatment, glycemic control and complications in type 2 diabetes in general practices (Disease Analyzer, Germany: 2008-2016). <i>Postgraduate Medicine</i> , 2018 , 130, 244-250	3.7 6

245	The impact of physician-level drug budgets on prescribing behavior. <i>European Journal of Health Economics</i> , 2018 , 19, 213-222	3.6	6
244	The impact of non-persistence on the direct and indirect costs in patients treated with subcutaneous tumour necrosis factor-alpha inhibitors in Germany. <i>Rheumatology</i> , 2018 , 57, 1276-1281	3.9	6
243	The Influence of Polypharmacy on the Initiation of Anti-Dementia Therapy in Germany. <i>Journal of Alzheimer's Disease</i> , 2018 , 64, 827-833	4.3	6
242	Demographic data and seasonal variation in peripheral vestibular disorders in ENT practices in Germany. <i>Journal of Vestibular Research: Equilibrium and Orientation</i> , 2019 , 29, 181-190	2.5	6
241	The prescribing of contraceptives for adolescents in German gynecologic practices in 2007 and 2011: a retrospective database analysis. <i>Journal of Pediatric and Adolescent Gynecology</i> , 2013 , 26, 261-4	2	6
240	Prescription Patterns and the Cost of Migraine Treatments in German General and Neurological Practices. <i>Pain Practice</i> , 2017 , 17, 747-752	3	6
239	Macro- and microvascular outcomes in patients with type 2 diabetes treated with rapid-acting insulin analogues or human regular insulin: a retrospective database analysis. <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2014 , 122, 92-9	2.3	6
238	Predictors for the initiation of a basal supported oral therapy (BOT) in type 2 diabetic patients under real-life conditions in Germany. <i>Primary Care Diabetes</i> , 2012 , 6, 329-35	2.4	6
237	Prescription of insulin glargine in primary care practices in Germany. <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2007 , 115, 252-6	2.3	6
236	Clinical and Patient-Related Variables Associated with Initiating GLP-1 Receptor Agonist Therapy in Type 2 Diabetes Patients in Primary Care in Germany. <i>PLoS ONE</i> , 2016 , 11, e0152281	3.7	6
235	Use of a German longitudinal prescription database (LRx) in pharmacoepidemiology. <i>GMS German Medical Science</i> , 2015 , 13, Doc14	3.2	6
234	Impact of caesarean section on mode of delivery, pregnancy-induced and pregnancy-associated disorders, and complications in the subsequent pregnancy in Germany. <i>GMS German Medical Science</i> , 2016 , 14, Doc06	3.2	6
233	Risk factors for ectopic pregnancy in Germany: a retrospective study of 100,197 patients. <i>GMS German Medical Science</i> , 2017 , 15, Doc19	3.2	6
232	Physicians' influence on breast cancer patient compliance. <i>GMS German Medical Science</i> , 2014 , 12, Doc03	3.2	6
231	A retrospective database analysis on persistence with inhaled corticosteroid therapy: comparison of two dry powder inhalers during asthma treatment in Germany. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2012 , 50, 257-64	2	6
230	Impact of comorbidities on the treatment of atopic dermatitis in clinical practice. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2014 , 52, 726-31	2	6
229	Risk of dementia in German patients treated with antidepressants in general or psychiatric practices?. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2017 , 55, 322-328	2	6
228	Patterns of benzodiazepine prescribing by neuropsychiatrists and general practitioners for elderly patients in Germany in 2014?. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2017 , 55, 466-471	2	6

227	Incident Dementia in Elderly Patients with Nonalcoholic Fatty Liver Disease in Germany. <i>Digestive Diseases and Sciences</i> , 2021 , 66, 3179-3185	4	6
226	Association Between Inflammatory Bowel Disease and Dementia: A Retrospective Cohort Study. <i>Journal of Alzheimer's Disease</i> , 2021 , 80, 1471-1478	4.3	6
225	The impact of the coronavirus disease (COVID-19) pandemic on outpatient epilepsy care: An analysis of physician practices in Germany. <i>Epilepsy and Behavior</i> , 2021 , 117, 107833	3.2	6
224	Physical and mental health comorbidities of patients with epilepsy in Germany - A retrospective cohort study. <i>Epilepsy and Behavior</i> , 2021 , 117, 107857	3.2	6
223	Persistence with biological drugs in patients treated in rheumatology practices in Germany. <i>Rheumatology International</i> , 2019 , 39, 525-531	3.6	6
222	Change in glucose-lowering medication regimens in individuals with type 2 diabetes mellitus during the COVID-19 pandemic in Germany. <i>Diabetes, Obesity and Metabolism</i> , 2021 , 23, 910-915	6.7	6
221	Consultations Decline for Stroke, Transient Ischemic Attack, and Myocardial Infarction during the COVID-19 Pandemic in Germany. <i>Neuroepidemiology</i> , 2021 , 1-8	5.4	6
220	Prevalence and risk factors for delirium diagnosis in patients followed in general practices in Germany. <i>International Psychogeriatrics</i> , 2018 , 30, 511-518	3.4	6
219	An elevated FIB-4 score predicts liver cancer development: A longitudinal analysis from 29,999 patients with NAFLD. <i>Journal of Hepatology</i> , 2021 ,	13.4	6
218	Is There an Association Between Antiepileptic Drug Use and Dementia Risk? A Case-Control Study. <i>Journal of Alzheimer's Disease</i> , 2019 , 68, 97-103	4.3	5
217	Relationship between induced abortion and the incidence of depression, anxiety disorder, adjustment disorder, and somatoform disorder in Germany. <i>Journal of Psychiatric Research</i> , 2019 , 114, 75-79	5.2	5
216	Prevalence of Use of Cardiovascular Drugs in Dementia Patients Treated in General Practices in Germany. <i>Journal of Alzheimer's Disease</i> , 2017 , 56, 1519-1524	4.3	5
215	Association of characteristics of people with type 2 diabetes mellitus with discordant values of fasting glucose and HbA1c. <i>Journal of Diabetes</i> , 2018 , 10, 934-941	3.8	5
214	Influence of macro- and microvascular comorbidity on time to insulin initiation in type 2 diabetes patients: a retrospective database analysis in Germany, France, and UK. <i>Primary Care Diabetes</i> , 2013 , 7, 167-71	2.4	5
213	Resource consumption and costs of treatment in patients with type 1 diabetes under intensified conventional therapy under German real-life conditions. <i>Journal of Diabetes Science and Technology</i> , 2013 , 7, 736-42	4.1	5
212	Prevalence of high-risk cardiovascular patients with therapy-resistant hypercholesterolemia. <i>Cardiovascular Endocrinology</i> , 2017 , 6, 81-85		5
211	Treatment Outcomes and Tolerability Following Initiation of GLP-1 Receptor Agonists Among Type 2 Diabetes Patients in Primary Care Practices in Germany. <i>Journal of Diabetes Science and Technology</i> , 2017 , 11, 272-277	4.1	5
210	Predictors of insulin initiation in metformin and sulfonylurea users in primary care practices: the role of kidney function. <i>Journal of Diabetes Science and Technology</i> , 2014 , 8, 1023-8	4.1	5

209	Prevalence of female subfertility in German gynecological practices. <i>Gynecological Endocrinology</i> , 2013 , 29, 767-70	2.4	5
208	Prevalence and predictors of prescription of antibiotics in pregnant women treated by gynecologists in Germany?. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2017 , 55, 643-649	2	5
207	Prevalence of switching from two anti-TNF biosimilars back to biologic reference products in Germany?. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2019 , 57, 323-328	2	5
206	The association between antihypertensive therapy and the incidence of Parkinson's disease in patients followed in general practices in Germany?. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2019 , 57, 483-488	2	5
205	Achalasia is associated with a higher incidence of depression in outpatients in Germany. <i>PLoS ONE</i> , 2021 , 16, e0250503	3.7	5
204	Hepatobiliary Adverse Drug Reactions Associated With Remdesivir: The WHO International Pharmacovigilance Study. <i>Clinical Gastroenterology and Hepatology</i> , 2021 , 19, 1970-1972.e3	6.9	5
203	Depression risk in patients with heart failure in primary care practices in Germany. <i>International Psychogeriatrics</i> , 2016 , 28, 1889-1894	3.4	5
202	Age-related differences in persistence with bisphosphonates in women with metastatic breast cancer. <i>Journal of Bone Oncology</i> , 2016 , 5, 63-6	4.5	5
201	Progression of mild cognitive impairment to dementia in German specialist practices. <i>Dementia</i> , 2019 , 18, 380-390	3	5
200	Association between antiseizure medication use and risk of urinary tract infection: A case-control study. <i>Epilepsy and Behavior</i> , 2021 , 115, 107502	3.2	5
199	Low annual frequency of HbA testing in people with Type 2 diabetes in primary care practices in Germany. <i>Diabetic Medicine</i> , 2018 , 35, 249-254	3.5	5
198	Changes in Glycemic Control and Body Weight After Initiation of Dapagliflozin or Basal Insulin Supported Oral Therapy in Type 2 Diabetes: A Primary Care Database Study. <i>Journal of Diabetes Science and Technology</i> , 2017 , 11, 590-596	4.1	4
197	Risk of Nonfatal Stroke in Type 2 Diabetes Mellitus Patients: A Retrospective Comparison Between Disease Management Programs and Standard Care. <i>Journal of Diabetes Science and Technology</i> , 2017 , 11, 808-813	4.1	4
196	Increased risk for cancer after stroke at a young age: etiological relevance or incidental finding?. <i>Journal of Cancer Research and Clinical Oncology</i> , 2019 , 145, 3047-3054	4.9	4
195	Emergency peripartum hysterectomy and risk factors by mode of delivery and obstetric history: a 10-year review from Helsinki University Central Hospital. <i>Journal of Perinatal Medicine</i> , 2015 , 43, 721-8	2.7	4
194	Comparative Analysis of the 3-Year Persistence Rate with Second-Line Vedolizumab and Tumor Necrosis Factor- α Inhibitors in Patients with Inflammatory Bowel Disease Followed in Gastroenterology Practices in Germany. <i>Digestive Diseases</i> , 2020 , 38, 466-473	3.2	4
193	Seasonal Variation of Epistaxis in Germany. <i>Journal of Craniofacial Surgery</i> , 2018 , 29, e365-e367	1.2	4
192	Tramadol prescription patterns in patients followed by general practitioners and orthopedists in Germany in the year 2015. <i>Postgraduate Medicine</i> , 2018 , 130, 37-41	3.7	4

191	Association between depression and persistence with oral antihyperglycemic drugs in type 2 diabetes mellitus patients in Germany. <i>Psychiatry Research</i> , 2018 , 261, 90-93	9.9	4
190	Persistence With Oral Antihyperglycemic Drugs in Type 2 Diabetes Mellitus Patients With Dementia in Germany. <i>Journal of Diabetes Science and Technology</i> , 2018 , 12, 140-144	4.1	4
189	Which Adverse Effects Influence the Dropout Rate in Selective Serotonin Reuptake Inhibitor (Ssri) Treatment?. <i>Value in Health</i> , 2014 , 17, A460	3.3	4
188	Persistence with Antihypertensive Drugs in Patients with Hypertension and Dementia in Germany. <i>Journal of Alzheimer's Disease</i> , 2017 , 60, 505-510	4.3	4
187	Risk factors for discontinuation of insulin pump therapy in pediatric and young adult patients. <i>Primary Care Diabetes</i> , 2014 , 8, 346-51	2.4	4
186	Basal Supported Oral Therapy with Insulin Glargine Results in Longer Persistence and Lower Costs Compared with Insulin Detemir in Type 2 Diabetics in Germany. <i>Health Outcomes Research in Medicine</i> , 2011 , 2, e39-e50		4
185	Längere Verweildauer unter einer basalunterstützten oralen Therapie mit Insulin Glargin (BOT) im Vergleich zu einer Kombinationstherapie aus NPH-Insulin und oralen Antidiabetika. <i>Diabetologie Und Stoffwechsel</i> , 2009 , 4, 166-171	0.7	4
184	Prevalence of depression, anxiety, adjustment disorders, and somatoform disorders in patients with age-related macular degeneration in Germany. <i>GMS German Medical Science</i> , 2017 , 15, Doc04	3.2	4
183	To Be Continued? Long-Term Treatment Effects of Antidepressant Drug Classes and Individual Antidepressants on the Risk of Developing Dementia: A German Case-Control Study. <i>Journal of Clinical Psychiatry</i> , 2020 , 81,	4.6	4
182	Micro- and macrovascular outcomes in Type 2 diabetic patients treated with insulin glulisine or human regular insulin: a retrospective database analysis. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2012 , 50, 821-9	2	4
181	Regional differences in insulin therapy regimens in five European countries?. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2017 , 55, 403-408	2	4
180	Diagnosing and treating dementia in German primary and specialized care between 2011 and 2015?. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2018 , 56, 301-309	2	4
179	Factors associated with fracture after stroke and TIA: a long-term follow-up. <i>Osteoporosis International</i> , 2020 , 31, 2395-2402	5.3	4
178	Proton pump inhibitors increase risk of bone fractures in men with cirrhosis: a population-based study. <i>Alimentary Pharmacology and Therapeutics</i> , 2020 , 52, 1042-1050	6.1	4
177	Proton pump inhibitor use is associated with a variety of infections in patients with liver cirrhosis. <i>Medicine (United States)</i> , 2020 , 99, e23436	1.8	4
176	Prevalence of overuse of short-acting beta-2 agonists (SABA) and associated factors among patients with asthma in Germany. <i>Respiratory Research</i> , 2021 , 22, 108	7.3	4
175	Treatment of Urinary Tract Infections with Canephron in Germany: A Retrospective Database Analysis. <i>Antibiotics</i> , 2021 , 10,	4.9	4
174	Osteoarthritis and incidence of stroke and transient ischemic attack in 320,136 adults followed in general practices in the United Kingdom. <i>Joint Bone Spine</i> , 2021 , 88, 105104	2.9	4

173	Trends in Dementia and Mild Cognitive Impairment Prevalence and Incidence in German General and Specialist Practices Between 2015 and 2019. <i>Journal of Alzheimer's Disease</i> , 2021 , 79, 1683-1690	4.3	4
172	Current healthcare pathways in the treatment of rhinosinusitis in Germany. <i>Acta Oto-Laryngologica</i> , 2018 , 138, 1086-1091	1.6	4
171	Pulmonary hypertension is associated with an increased incidence of NAFLD: A retrospective cohort study of 18,910 patients. <i>Journal of Internal Medicine</i> , 2021 , 290, 886-893	10.8	4
170	Seasonality of Insulin Use in German Outpatients With Diabetes: A Retrospective Analysis. <i>Journal of Diabetes Science and Technology</i> , 2017 , 11, 996-1000	4.1	3
169	Duration of sick leave in patients with depression treated by general practitioners and psychiatrists. <i>Psychiatry Research</i> , 2019 , 279, 382-383	9.9	3
168	Increased prevalence of depression, anxiety, and adjustment and somatoform disorders in transsexual individuals. <i>Journal of Affective Disorders</i> , 2020 , 274, 482-485	6.6	3
167	Prevalence of insulin as a first-line therapy and associated factors in people with type 2 diabetes in German primary care practices. <i>Diabetic Medicine</i> , 2020 , 37, 1333-1339	3.5	3
166	Association between various thyroid gland diseases, TSH values and thyroid cancer: a case-control study. <i>Journal of Cancer Research and Clinical Oncology</i> , 2020 , 146, 2989-2994	4.9	3
165	Frequency of blood pressure and estimated glomerular filtration rate monitoring in patients affected by hypertension: a retrospective study with 176 565 patients in Germany. <i>Blood Pressure Monitoring</i> , 2018 , 23, 85-90	1.3	3
164	Long-term use of basal insulin and the risk of breast cancer. <i>Breast Cancer Research and Treatment</i> , 2018 , 168, 763-764	4.4	3
163	Prescription Patterns and Disease Control in Type 2 Diabetes Mellitus Patients in Nursing Home and Home Care Settings: A Retrospective Analysis in Germany. <i>Journal of Diabetes Science and Technology</i> , 2018 , 12, 136-139	4.1	3
162	Cancer is associated with intraoperative and postprocedural complications and disorders. <i>Journal of Cancer Research and Clinical Oncology</i> , 2016 , 142, 777-81	4.9	3
161	Prescription-based prevalence of biological therapy in patients with psoriasis, rheumatoid arthritis, and inflammatory bowel diseases. <i>Biologicals</i> , 2019 , 61, 52-54	1.8	3
160	One-Year Persistence with Donepezil, Memantine, and Rivastigmine in More than 66,000 Elderly Patients Followed in Poland. <i>Journal of Alzheimer's Disease</i> , 2019 , 70, 899-905	4.3	3
159	Effects of a low-dose oral estrogen only treatment on bone mineral density and quantitative ultrasonometry in postmenopausal women. <i>Gynecological Endocrinology</i> , 2012 , 28, 1002-5	2.4	3
158	Different persistence on initial basal supported oral therapy in Type 2 diabetics is associated with unequal distributions of insulin treatment regimens under real-life conditions in Germany. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2010 , 48, 761-6	2	3
157	Suspected cancer diagnoses made by general practitioners in a population with subsequently confirmed cancer diagnoses in Germany: a retrospective study of 31,628 patients. <i>Oncotarget</i> , 2017 , 8, 84540-84545	3.3	3
156	Risk of hypoglycaemia in type 2 diabetes patients under different insulin regimens: a primary care database analysis. <i>GMS German Medical Science</i> , 2015 , 13, Doc01	3.2	3

155	Conflicts at work are associated with a higher risk of cardiovascular disease. <i>GMS German Medical Science</i> , 2017 , 15, Doc08	3.2	3
154	The negative impact of rebate contracts on the health care of patients with depression in Germany. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2011 , 49, 397-402	2	3
153	Frequency of hospitalizations prior to and after conversion to a rebate pharmaceutical in depression patients in Germany. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2013 , 51, 416-22	2	3
152	Differences in baseline characteristics between type 2 diabetes mellitus patients treated with dipeptidyl peptidase-4 inhibitors in randomized controlled trials and those receiving the same treatment in real-world settings?. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2013 , 51, 416-22	2	3
151	Persistence and adherence to dutasteride/tamsulosin fixed-dose versus free-combination alpha blocker/5ARI therapy in patients with benign prostate hyperplasia in Germany?. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2020 , 58, 37-49	2	3
150	Comparative Analysis of 3-Year Persistence With Vedolizumab Compared With Antibodies Against Tumor Necrosis Factor-Alpha in Patients With Inflammatory Bowel Disease in Germany: Retrospective Analysis of a Large Prescription Database. <i>Journal of Clinical Gastroenterology</i> , 2021 , 55, 141-147	3	3
149	Compliance with vaccination against hepatitis A virus in Germany: A retrospective analysis?. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2017 , 55, 740-745	2	3
148	Adherence to neuroleptic treatment in psychiatric practices: A retrospective study of 55 practices with more than 5000 bipolar and schizophrenic patients in Germany. <i>Psychiatry Research</i> , 2020 , 284, 112758	2.9	3
147	Fecal and urinary incontinence are major problems associated with rectal cancer. <i>International Journal of Colorectal Disease</i> , 2020 , 35, 35-40	3	3
146	Real-world treatment patterns in patients with nontuberculous mycobacterial lung disease in general and pneumologist practices in Germany. <i>Journal of Clinical Tuberculosis and Other Mycobacterial Diseases</i> , 2020 , 20, 100178	2.1	3
145	Association between skin disorders and depression in children and adolescents: A retrospective case-control study. <i>Journal of Affective Disorders</i> , 2021 , 282, 939-944	6.6	3
144	Incidence and Risk of Various Disorders of the External Ear in Patients With Hearing Aids Treated in ENT Practices in Germany. <i>Otology and Neurotology</i> , 2019 , 40, 1076-1081	2.6	3
143	Prescription patterns of antiepileptic drugs in Kazakhstan in 2018: A retrospective study of 57,959 patients. <i>Epilepsy and Behavior</i> , 2019 , 99, 106445	3.2	3
142	Osteoarthritis and the incidence of fracture in the United Kingdom: a retrospective cohort study of 258,696 patients. <i>Osteoarthritis and Cartilage</i> , 2021 , 29, 215-221	6.2	3
141	Medication use in patients with chronic rhinosinusitis in Germany - a large retrospective patient-based study. <i>Rhinology</i> , 2019 , 57, 94-100	7	3
140	Prevalence of and factors associated with COVID-19 diagnosis in symptomatic patients followed in general practices in Germany between March 2020 and March 2021. <i>International Journal of Infectious Diseases</i> , 2021 , 111, 37-42	10.5	3
139	Effect of the COVID-19 pandemic on stress, anxiety, and depressive disorders in German primary care: A cross-sectional study. <i>Journal of Psychiatric Research</i> , 2021 , 143, 43-49	5.2	3
138	Calculated Daily Insulin Dosages Overestimate Prescribed Insulin Doses in Type 2 Diabetes: A Primary Care Database Study. <i>Journal of Diabetes Science and Technology</i> , 2017 , 11, 597-601	4.1	2

137	Persistence with selective serotonin (norepinephrine) reuptake inhibitors in Germany-A retrospective database analysis. <i>Journal of Affective Disorders</i> , 2019 , 247, 156-160	6.6	2
136	Coded Prevalence of Mild Cognitive Impairment in General and Neuropsychiatrists Practices in Germany Between 2007 and 2017. <i>Journal of Alzheimer's Disease</i> , 2019 , 67, 1313-1318	4.3	2
135	Longitudinal changes in HbA1c following therapy intensification after inadequate response to two oral antidiabetic agents in patients with type 2 diabetes in Germany. <i>Diabetes, Obesity and Metabolism</i> , 2019 , 21, 1757	6.7	2
134	Diagnostic Behavior for Mild Cognitive Impairment in General and Neuropsychiatric Practices in Germany. <i>Journal of Alzheimer's Disease</i> , 2019 , 68, 925-930	4.3	2
133	Time to Insulin Initiation in Type 2 Diabetes Patients in 2010/2011 and 2016/2017 in Germany. <i>Journal of Diabetes Science and Technology</i> , 2019 , 13, 1129-1134	4.1	2
132	Schwangerschaften nach Brustkrebs in Deutschland [Ergebnisse einer retrospektiven Datenbankanalyse. <i>Tumor Diagnostik Und Therapie</i> , 2015 , 36, 226-229	0.1	2
131	Prescription of hormone replacement therapy prior to and after the diagnosis of gynecological cancers in German patients. <i>Journal of Cancer Research and Clinical Oncology</i> , 2020 , 146, 1567-1573	4.9	2
130	Topical therapy of hand eczema - analysis of the prescription profile from dermatologists in private practice. <i>JDDG - Journal of the German Society of Dermatology</i> , 2012 , 10, 180-4	1.2	2
129	Fracture risk reduction with use of dipeptidyl peptidase-4 inhibitors: response to Driessen et al. <i>Osteoporosis International</i> , 2017 , 28, 2431	5.3	2
128	Risk of diabetic foot ulceration during treatment with insulin glargine and NPH insulin. <i>Journal of Wound Care</i> , 2012 , 21, 483-4, 486-9	2.2	2
127	PCV108 RISK AND COSTS OF THE FIRST HYPERTENSION-ASSOCIATED EVENT, COMPLIANCE AND PERSISTENCE IN NAIVE HYPERTENSIVE PATIENTS AFTER INITIATING MONOTHERAPY. <i>Value in Health</i> , 2008 , 11, A415	3.3	2
126	Decrease in the number of patients diagnosed with cancer during the COVID-19 pandemic in Germany.. <i>Journal of Cancer Research and Clinical Oncology</i> , 2022 , 1	4.9	2
125	Investigation of the prevalence of non-COVID-19 infectious diseases during the COVID-19 pandemic.. <i>Public Health</i> , 2021 , 203, 53-57	4	2
124	Gender-based differences in the antidepressant treatment of patients with depression in German psychiatric practices. <i>GMS German Medical Science</i> , 2016 , 14, Doc02	3.2	2
123	Prevalence and treatment of sexually transmitted infections in men followed by urologists in Germany - a cross sectional study with 347,090 men. <i>GMS German Medical Science</i> , 2018 , 16, Doc03	3.2	2
122	Gender differences in persistency to bisphosphonates in patients with metastatic breast and prostate cancer. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2014 , 52, 352-9	2	2
121	Patterns of insulin therapy and insulin daily doses in children and adolescents with type 1 diabetes in Germany. <i>Diabetes, Obesity and Metabolism</i> , 2021 ,	6.7	2
120	Antiherpetic medication and incident dementia: observational cohort studies in four countries		2

119	Depression and sleep disorders are associated with early mortality in women with breast cancer in the United Kingdom. <i>Journal of Psychiatric Research</i> , 2021 , 143, 481-484	5.2	2
118	Multimorbidity Among Adult Outpatients With Type 1 Diabetes in Germany. <i>Journal of Diabetes Science and Technology</i> , 2020 , 1932296820965261	4.1	2
117	Incidences of hypothyroidism and autoimmune thyroiditis are increased in patients with nonalcoholic fatty liver disease. <i>European Journal of Gastroenterology and Hepatology</i> , 2021 ,	2.2	2
116	Mild cognitive impairment is associated with fall-related injury among adults aged 65 years in low- and middle-income countries. <i>Experimental Gerontology</i> , 2021 , 146, 111222	4.5	2
115	Association between sleep disorders and osteoarthritis: A case-control study of 351,932 adults in the UK. <i>Journal of Sleep Research</i> , 2021 , 30, e13367	5.8	2
114	Variables associated with increased incidence of non-alcoholic fatty liver disease (NAFLD) in patients with type 2 diabetes. <i>BMJ Open Diabetes Research and Care</i> , 2021 , 9,	4.5	2
113	Impact of the COVID-19 Pandemic on Consultations and Diagnoses in Gastroenterology Practices in Germany. <i>Frontiers in Medicine</i> , 2021 , 8, 684032	4.9	2
112	Incidence of antiepileptic drug therapy and factors associated with their prescribing in outpatients with diabetic polyneuropathy. <i>Primary Care Diabetes</i> , 2021 , 15, 535-540	2.4	2
111	Anti-seizure medication is not associated with an increased risk to develop cancer in epilepsy patients. <i>Journal of Neurology</i> , 2021 , 268, 2185-2191	5.5	2
110	Impact of Non-Alcoholic Fatty Liver Disease on Metabolic Comorbidities in Type 2 Diabetes Mellitus. <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2021 ,	2.3	2
109	Relevance of Coded Prodromal Mild Cognitive Impairment in the Routine Treatment of Patients with Dementia in Germany. <i>Journal of Alzheimer's Disease</i> , 2018 , 65, 393-399	4.3	2
108	Diagnoses of suspected cancer in otolaryngology practices in Germany. <i>Molecular and Clinical Oncology</i> , 2018 , 9, 459-463	1.6	2
107	What have we learned in the past year? A study on pharmacy purchases of psychiatric drugs from wholesalers in the days prior to the first and second COVID-19 lockdowns in Germany. <i>Journal of Psychiatric Research</i> , 2021 , 140, 346-349	5.2	2
106	The spectrum of comorbidities at the initial diagnosis of heart failure a case control study.. <i>Scientific Reports</i> , 2022 , 12, 2670	4.9	2
105	Prevalence of diabetes mellitus among 80,193 gastrointestinal cancer patients in five European and three Asian countries.. <i>Journal of Cancer Research and Clinical Oncology</i> , 2021 , 148, 1057	4.9	2
104	Low blood levels of high-density lipoprotein (HDL) cholesterol are positively associated with cancer.. <i>Journal of Cancer Research and Clinical Oncology</i> , 2021 , 1	4.9	2
103	Persistence and treatment-free interval in patients being prescribed biological drugs in rheumatology practices in Germany. <i>European Journal of Clinical Pharmacology</i> , 2019 , 75, 717-722	2.8	1
102	Real-world prevalence and incidence of lower back pain in Germany. <i>Rheumatology International</i> , 2019 , 39, 1113	3.6	1

101	Treatment Patterns of Patients with All-Cause Dementia in Russia. <i>Journal of Alzheimer's Disease Reports</i> , 2020 , 4, 9-14	3.3	1
100	Effect of progestogen-only contraception on premenopausal fracture risk: a case-control study. <i>Osteoporosis International</i> , 2020 , 31, 1801-1806	5.3	1
99	Relationship between physical conditions and attempted or completed suicide in more than 9,300 individuals from the United Kingdom: a case-control study. <i>Journal of Affective Disorders</i> , 2020 , 274, 457-463	6.6	1
98	Usage of antiepileptic drugs in different diseases in Germany - A retrospective study. <i>Epilepsy and Behavior</i> , 2020 , 111, 107210	3.2	1
97	Sex Differences in People Aging With HIV in Germany. <i>Journal of Acquired Immune Deficiency Syndromes (1999)</i> , 2020 , 84, e11	3.1	1
96	Association between gout and osteoporosis risk in the United Kingdom. <i>Osteoporosis International</i> , 2018 , 29, 1685	5.3	1
95	Frequency of Thyroid-Stimulating Hormone Monitoring in Patients After Thyroidectomy: A Retrospective Study in Primary Care Practices in Germany. <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2018 , 126, 640-644	2.3	1
94	Time between suspected and confirmed diagnoses of Crohn's disease and ulcerative colitis in patients followed in gastroenterological practices in Germany. <i>International Journal of Colorectal Disease</i> , 2018 , 33, 967-971	3	1
93	Are data from general practices suitable for survival analyses in the field of breast cancer? A retrospective study conducted in the United Kingdom, France and Germany. <i>Molecular and Clinical Oncology</i> , 2019 , 11, 177-180	1.6	1
92	Late Detection of Atrial Fibrillation after Stroke: Implications for the Secondary Prevention. <i>European Neurology</i> , 2019 , 81, 262-269	2.1	1
91	Comparing Quality Effects Of Patient Care In Integrated And Regular Care For Patients With Hypertension. <i>Value in Health</i> , 2014 , 17, A498	3.3	1
90	Secondary analysis of electronic databases: potentials and limitations. Reply to Asghari S, Mahdavian M [letter]. <i>Diabetologia</i> , 2013 , 56, 2098-9	10.3	1
89	Longitudinal change in HbA1c in patients with type 2 diabetes: comparison of short-acting, biphasic and basal insulin. <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2009 , 117, 230-3	2.3	1
88	Modellierung der Langzeit-Behandlungskosten insulinpflichtiger Typ-2-Diabetiker in Abhängigkeit von der Art des Insulineinstiegs: Vergleich zwischen Insulin glargin und NPH-Insulin. <i>Gesundheitsökonomie Und Qualitätsmanagement</i> , 2011 , 16, 182-188	0.3	1
87	Association between ivy leaves dry extract EA 575 prescriptions and antibiotic use, sick leave duration and repeated infections in adult patients. <i>Postgraduate Medicine</i> , 2021 ,	3.7	1
86	Overlap between irritable bowel syndrome and common gastrointestinal diagnoses: a retrospective cohort study of 29,553 outpatients in Germany.. <i>BMC Gastroenterology</i> , 2022 , 22, 48	3	1
85	Age-Related Changes in the Association Between Traumatic Brain Injury and Dementia in Older Men and Women. <i>Journal of Head Trauma Rehabilitation</i> , 2021 , 36, E139-E146	3	1
84	Diabetes mellitus is associated with an increased incidence of aortic valve stenosis. <i>Diabetes and Vascular Disease Research</i> , 2021 , 18, 14791641211033819	3.3	1

83	Diabetes mellitus is associated with an increased incidence of aortic valve stenosis. <i>Diabetes and Vascular Disease Research</i> , 2021 , 18, 147916412110338	3.3	1
82	Increase in depression and anxiety disorder diagnoses during the COVID-19 pandemic in children and adolescents followed in pediatric practices in Germany. <i>European Child and Adolescent Psychiatry</i> , 2021 , 1	5.5	1
81	No significant association between COVID-19 diagnosis and the incidence of depression and anxiety disorder? A retrospective cohort study conducted in Germany.. <i>Journal of Psychiatric Research</i> , 2022 , 147, 79-84	5.2	1
80	Economic prescribing of corticosteroid nasal sprays in Germany: comparison of mometasone and budesonide nasal sprays on the basis of the DDD, the PDD and reference prices. <i>International Journal of Clinical Pharmacology and Therapeutics</i> , 2013 , 51, 12-8	2	1
79	Non-alcoholic fatty liver disease (NAFLD) is associated with an increased incidence of osteoporosis and bone fractures. <i>Zeitschrift Fur Gastroenterologie</i> , 2021 ,	1.6	1
78	Epilepsy is associated with an increased incidence of heart failure diagnoses. <i>Epilepsy and Behavior</i> , 2021 , 125, 108393	3.2	1
77	Association Between Migraine Headaches and Dry Eye Disease in Patients Studied in General Practices in Germany. <i>JAMA Ophthalmology</i> , 2020 , 138, 223	3.9	1
76	Long-term risk of stroke and its predictors in transient ischaemic attack patients in Germany. <i>European Journal of Neurology</i> , 2020 , 27, 723-728	6	1
75	Comment on Mizokami-Stout et al. The Contemporary Prevalence of Diabetic Neuropathy in Type 1 Diabetes: Findings From the T1D Exchange. <i>Diabetes Care</i> 2020;43:806-812. <i>Diabetes Care</i> , 2020 , 43, e113	14.6	1
74	Associations of physical and psychiatric conditions with chronic fatigue syndrome in Germany: an exploratory case-control study. <i>Psychological Medicine</i> , 2020 , 1-7	6.9	1
73	Changes in the utilization of blood glucose test strips among patients using intermittent-scanning continuous glucose monitoring in Germany. <i>Diabetes, Obesity and Metabolism</i> , 2020 , 22, 922-928	6.7	1
72	Influence of chemotherapy and endocrine treatment on fractures in postmenopausal women with breast cancer - a retrospective cohort study. <i>Journal of Bone Oncology</i> , 2020 , 22, 100292	4.5	1
71	Association between inflammatory central nervous system diseases and epilepsy: A retrospective cohort study of 4252 patients in Germany. <i>Epilepsy and Behavior</i> , 2021 , 117, 107879	3.2	1
70	Prevalence of and Factors Associated with Antibiotic Prescriptions in Patients with Acute Lower and Upper Respiratory Tract Infections-A Case-Control Study. <i>Antibiotics</i> , 2021 , 10,	4.9	1
69	(Absence of) Association Between Non-Vitamin K Antagonist Oral Anticoagulant Therapy and Urinary Tract Infection in Patients With Atrial Fibrillation. <i>Journal of Cardiovascular Pharmacology</i> , 2021 , 77, 830-834	3.1	1
68	Multimorbidity and Subjective Cognitive Complaints: Findings from 48 Low- and Middle-Income Countries of the World Health Survey 2002-2004. <i>Journal of Alzheimer's Disease</i> , 2021 , 81, 1737-1747	4.3	1
67	Relationship between chronic gingivitis and subsequent depression in 13,088 patients followed in general practices. <i>Journal of Psychiatric Research</i> , 2021 , 138, 103-106	5.2	1
66	Impact of comorbidities on the cost of depression drug therapy in general practices in Germany. <i>Journal of Psychiatric Research</i> , 2016 , 83, 130-136	5.2	1

65	Frequency of Blood Pressure and Estimated Glomerular Filtration Rate Testing in Type 2 Diabetes Mellitus: A Retrospective Study with 43,509 Patients. <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2019 , 127, 455-460	2.3	1
64	Burn injury and incidence of psychiatric disorders: A retrospective cohort study of 18,198 patients from Germany. <i>Burns</i> , 2021 , 47, 1110-1117	2.3	1
63	Incidence of Inner Ear Disorders in Various Forms of Acute Otitis Media in ENT Practices in Germany. <i>Ear, Nose and Throat Journal</i> , 2021 , 100, 325S-332S	1	1
62	Prevalence of urinary system, pelvic organ, and genital tract disorders among women with type 1 diabetes in Germany. <i>Primary Care Diabetes</i> , 2021 , 15, 257-261	2.4	1
61	P1930Dronedarone is associated with a lower risk of myocardial infarction in Germany - a retrospective database analysis. <i>European Heart Journal</i> , 2018 , 39,	9.5	1
60	Impact of thyroid disorders on the incidence of non-alcoholic fatty liver disease in Germany. <i>United European Gastroenterology Journal</i> , 2021 , 9, 829	5.3	1
59	Association between type 2 diabetes and chronic low back pain in general practices in Germany. <i>BMJ Open Diabetes Research and Care</i> , 2021 , 9,	4.5	1
58	Prevalence of, and factors associated with, long-term COVID-19 sick leave in working-age patients followed in general practices in Germany. <i>International Journal of Infectious Diseases</i> , 2021 , 109, 203-208 ^{10.5}	10.5	1
57	Increased Incidence of Stroke and Transient Ischemic Attack in Patients with Rheumatoid Arthritis and Ankylosing Spondylitis in Germany. <i>Neuroepidemiology</i> , 2021 , 55, 162-170	5.4	1
56	Predicting the Risk of Ischemic Stroke in Patients Treated with Novel Oral Anticoagulants: A Machine Learning Approach. <i>Neuroepidemiology</i> , 2021 , 55, 387-392	5.4	1
55	Obesity and lipid metabolism disorders determine the risk for development of long COVID syndrome: a cross-sectional study from 50,402 COVID-19 patients.. <i>Infection</i> , 2022 , 1	5.8	1
54	Prevalence of valproate prescriptions in women of childbearing age in certain regions of Russia. <i>Epilepsy and Behavior</i> , 2019 , 101, 106584	3.2	0
53	Prevalence of Lung Metastases among 19,321 Metastatic Colorectal Cancer Patients in Eight Countries of Europe and Asia.. <i>Current Oncology</i> , 2021 , 28, 5035-5040	2.8	0
52	Association between osteoarthritis and the incidence of Parkinson's disease in the United Kingdom. <i>Clinical Parkinsonism & Related Disorders</i> , 2021 , 5, 100120	0.9	0
51	Multimorbidity and Anxiety Symptoms among Adults Aged 50 Years and Over from Six Low- and Middle-Income Countries. <i>Journal of Ageing and Longevity</i> , 2021 , 1, 36-47		0
50	Age effects on treatment patterns in 138,097 patients with unipolar depression followed in general practices in Germany. <i>Journal of Psychiatric Research</i> , 2021 , 144, 208-216	5.2	0
49	Long-term use of proton pump inhibitors (PPIs) is associated with an increased risk of type 2 diabetes. <i>Gut</i> , 2021 ,	19.2	0
48	Incidence of and factors associated with manic episodes and bipolar disorder in the decade following depression onset in Germany. <i>Journal of Affective Disorders</i> , 2020 , 266, 534-539	6.6	0

47	Is loneliness associated with mild cognitive impairment in low- and middle-income countries?. <i>International Journal of Geriatric Psychiatry</i> , 2021 , 36, 1345-1353	3.9	o
46	Is there an association between multiple sclerosis and osteoarthritis in Germany? A retrospective cohort study of 8,600 patients from Germany. <i>Multiple Sclerosis Journal - Experimental, Translational and Clinical</i> , 2021 , 7, 20552173211022784	2	o
45	Diverticular disease is associated with an increased incidence rate of depression and anxiety disorders. <i>International Journal of Colorectal Disease</i> , 2021 , 36, 2437-2443	3	o
44	Inflammatory Bowel Disease as a Precondition for Stroke or TIA: A Matter of Crohn's Disease Rather than Ulcerative Colitis. <i>Journal of Stroke and Cerebrovascular Diseases</i> , 2021 , 30, 105787	2.8	o
43	Prevalence of Hospitalization and Readmission in Patients with Inflammatory Bowel Diseases. <i>Digestive Diseases and Sciences</i> , 2021 , 1	4	o
42	Clinical characteristics and sick leave associated with infectious mononucleosis in a real-world setting in Germany. <i>International Journal of Clinical Practice</i> , 2021 , 75, e14690	2.9	o
41	Sarcopenia and Mild Cognitive Impairment in Older Adults from Six Low- and Middle-Income Countries. <i>Journal of Alzheimer's Disease</i> , 2021 , 82, 1745-1754	4.3	o
40	Effects of the COVID-19 lockdown on primary health care for persons with type 2 diabetes - Results from the German Disease Analyzer database. <i>Diabetes Research and Clinical Practice</i> , 2021 , 179, 109002	7.4	o
39	Predicting the risk of stroke in patients with late-onset epilepsy: A machine learning approach. <i>Epilepsy and Behavior</i> , 2021 , 122, 108211	3.2	o
38	Factors associated with the diagnosis of depression in women followed in gynecological practices in Germany. <i>Journal of Psychiatric Research</i> , 2021 , 141, 358-363	5.2	o
37	Bipolar Disorder and Outcomes of Monotherapy with Lithium, Valproate, Quetiapine, Olanzapine, Venlafaxine, and Citalopram. <i>Pharmacopsychiatry</i> , 2021 , 54, 126-130	2	o
36	Multiple Sclerosis and Incidence of Urinary and Fecal Incontinence in Almost 9,000 Patients Followed Up for up to 10 Years in Germany. <i>Neuroepidemiology</i> , 2021 , 55, 92-99	5.4	o
35	Prevalence of and factors associated with long-term sick leave in working-age adults with chronic low back pain in Germany.. <i>International Archives of Occupational and Environmental Health</i> , 2022 , 1	3.2	o
34	No increased incidence of venous thrombosis or pulmonary embolism after SARS-CoV-2 vaccination in Germany.. <i>Public Health</i> , 2022 , 207, 14-18	4	o
33	Effects of the COVID-19 pandemic on clinically diagnosed psychiatric disorders in persons with type 2 diabetes.. <i>Diabetic Medicine</i> , 2022 , e14852	3.5	o
32	Omega-3 fatty acids supplementation on major cardiovascular outcomes: an umbrella review of meta-analyses of observational studies and randomized controlled trials. <i>European Review for Medical and Pharmacological Sciences</i> , 2021 , 25, 2079-2092	2.9	o
31	Mild Cognitive Disorder in Post-COVID-19 Syndrome: A Retrospective Cohort Study of 67,000 Primary Care Post-COVID Patients. <i>Journal of Alzheimer's Disease Reports</i> , 1-9	3.3	o
30	Effektivität der glukosesenkenden Therapie bei Patienten mit Typ-2-Diabetes in primärärztlichen Praxen. <i>Diabetologe</i> , 2019 , 15, 89-95	0.2	

29	Discontinuation of Antihypertensive Therapy and Dementia Incidence in the United Kingdom. <i>Journal of the American Medical Directors Association</i> , 2019 , 20, 790-791	5.9
28	Einfluss des Disease-Management-Programms auf den HbA1c-Wert bei Typ-2-Diabetes-mellitus-Patienten: Ein retrospektiver Vergleich zwischen Disease Management Programm und Regelversorgung. <i>Diabetologie Und Stoffwechsel</i> , 2017 , 12, 455-462	0.7
27	Referral for occupational therapy after diagnosis of developmental disorder by German child psychiatrists. <i>Early Child Development and Care</i> , 2016 , 186, 959-970	0.9
26	Prevalence of Chronic Diseases Among Older Patients (>65 Years) in German General Practitioner Practices. <i>Value in Health</i> , 2014 , 17, A505	3.3
25	Treatment Discontinuation In Patients With Urinary Incontinence Suffering From Glaucoma. <i>Value in Health</i> , 2014 , 17, A471	3.3
24	Duration of first prescribed long acting insulin therapy in type 2 diabetes. <i>Diabetes Research and Clinical Practice</i> , 2012 , 97, e3	7.4
23	Valide Kosten-Nutzenbewertung im Gesundheitssystem [Fehlsteuerung] vermeiden durch Abbildung der Versorgungswirklichkeit (eine Real-World-Studie am Beispiel der Refluxerkrankung). <i>Gesundheitsökonomie Und Qualitätsmanagement</i> , 2008 , 13, 228-233	0.3
22	Authors' reply: Pulmonary hypertension is associated with an increased incidence of NAFLD.. <i>Journal of Internal Medicine</i> , 2022 ,	10.8
21	Autoimmune disorders are positively associated with achalasia.. <i>Neurogastroenterology and Motility</i> , 2022 , e14327	4
20	Frequency of over- and under-treatment with levothyroxine in primary care in Germany.. <i>European Journal of Endocrinology</i> , 2022 , 186, L5	6.5
19	No association between antiviral treatment and risk of Alzheimer's disease in German outpatients. <i>Alzheimer's and Dementia: Translational Research and Clinical Interventions</i> , 2021 , 7, e12216	6
18	Burden of depression and anxiety disorders per disease codes in patients with lymphoma in Germany. <i>Supportive Care in Cancer</i> , 2021 , 1	3.9
17	Factor V Leiden and the 10-year incidence of depression: A retrospective cohort study conducted in Germany.. <i>Journal of Psychiatric Research</i> , 2021 , 146, 87-91	5.2
16	Direct Oral Anticoagulants Versus Preprohormone in Patients With Cirrhosis and the Risk of Gastrointestinal Bleeding. <i>Annals of Pharmacotherapy</i> , 2021 , 10600280211048008	2.9
15	Prevalence of and Factors Associated With the Prescription of Fibrates Among Patients Receiving Lipid-Lowering Drugs in Germany. <i>Journal of Cardiovascular Pharmacology</i> , 2021 , 78, 885-890	3.1
14	Importance of durational hormone therapy in breast cancer patients in Germany: Retrospective database analysis.. <i>Journal of Clinical Oncology</i> , 2014 , 32, 1603-1603	2.2
13	Association of diabetes and cancer-diagnoses in primary care practices in Germany.. <i>Journal of Clinical Oncology</i> , 2014 , 32, 1604-1604	2.2
12	Examples of Existing Automated Databases	123-177

11	Association between sexual orientation and subjective cognitive complaints in the general population in England. <i>Journal of Psychiatric Research</i> , 2021 , 136, 1-6	5.2
10	In Reply. <i>Deutsches Arzteblatt International</i> , 2021 , 118, 271-272	2.5
9	Comment on Eilam et al. Incidence of acute ischemic cerebrovascular events in a cohort of HIV-infected patients. <i>Acta Neurologica Belgica</i> , 2021 , 1	1.5
8	Epidemiology of Gynaecological and Breast Cancers. <i>Advances in Medical Diagnosis, Treatment, and Care</i> , 2021 , 1-21	0.2
7	Manufacturer switch of anti-seizure drugs may not increase the risk of seizure recurrence in Children: A nationwide study of prescription data in Germany. <i>Epilepsy and Behavior</i> , 2021 , 115, 107705	3.2
6	Effect of SARS-CoV-2 vaccination among health care workers in a geriatric care unit after a B.1.1.7-variant outbreak. <i>Public Health</i> , 2021 , 198, e20-e22	4
5	An elevated FIB-4 score is associated with an increased incidence of liver cancer: A longitudinal analysis among 248,224 outpatients in Germany.. <i>European Journal of Cancer</i> , 2022 , 168, 41-50	7.5
4	Portal vein thrombosis is associated with an increased risk of bone fractures.. <i>PLoS ONE</i> , 2022 , 17, e0267535	3.7
3	Atrial fibrillation is associated with a subsequent epilepsy diagnosis independent of stroke: A retrospective matched administrative cohort study on 149,632 patients.. <i>Epilepsy and Behavior</i> , 2022 , 132, 108721	3.2
2	Epidemiology of Gynaecological and Breast Cancers 2022 , 187-212	
1	Sex-Differential Associations Between Body Mass Index and the Incidence of Dementia. <i>Journal of Alzheimer's Disease</i> , 2022 , 1-9	4.3