Kirsten Sundby Hall

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/3413852/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Comparison of MAPIE versus MAP in patients with a poor response to preoperative chemotherapy for newly diagnosed high-grade osteosarcoma (EURAMOS-1): an open-label, international, randomised controlled trial. Lancet Oncology, The, 2016, 17, 1396-1408.	10.7	356
2	Survival and prognosis with osteosarcoma: outcomes in more than 2000 patients in the EURAMOS-1 (European and American Osteosarcoma Study) cohort. European Journal of Cancer, 2019, 109, 36-50.	2.8	354
3	Methotrexate, Doxorubicin, and Cisplatin (MAP) Plus Maintenance Pegylated Interferon Alfa-2b Versus MAP Alone in Patients With Resectable High-Grade Osteosarcoma and Good Histologic Response to Preoperative MAP: First Results of the EURAMOS-1 Good Response Randomized Controlled Trial. Iournal of Clinical Oncology, 2015, 33, 2279-2287.	1.6	329
4	Adjuvant Imatinib for High-Risk GI Stromal Tumor: Analysis of a Randomized Trial. Journal of Clinical Oncology, 2016, 34, 244-250.	1.6	174
5	Effect of <i>KIT</i> and <i>PDGFRA</i> Mutations on Survival in Patients With Gastrointestinal Stromal Tumors Treated With Adjuvant Imatinib. JAMA Oncology, 2017, 3, 602.	7.1	141
6	Radiotherapy to Improve Local Control Regardless of Surgical Margin and Malignancy Grade in Extremity and Trunk Wall Soft Tissue Sarcoma: A Scandinavian Sarcoma Group Study. International Journal of Radiation Oncology Biology Physics, 2008, 71, 1196-1203.	0.8	129
7	Clinical course of nonvisceral soft tissue leiomyosarcoma in 225 patients from the Scandinavian Sarcoma Group. Cancer, 2007, 109, 282-291.	4.1	123
8	Survival Outcomes Associated With 3 Years vs 1 Year of Adjuvant Imatinib for Patients With High-Risk Gastrointestinal Stromal Tumors. JAMA Oncology, 2020, 6, 1241.	7.1	111
9	High dose methotrexate chemotherapy: pharmacokinetics, folate and toxicity in osteosarcoma patients. British Journal of Clinical Pharmacology, 2012, 73, 106-114.	2.4	101
10	Ultraâ€rare sarcomas: A consensus paper from the Connective Tissue Oncology Society community of experts on the incidence threshold and the list of entities. Cancer, 2021, 127, 2934-2942.	4.1	96
11	Radiation-induced sarcoma: 25-year experience from The Norwegian Radium Hospital. Acta Oncológica, 2008, 47, 1475-1482.	1.8	93
12	EURO-B.O.S.S.: A European study on chemotherapy in bone-sarcoma patients aged over 40: Outcome in primary high-grade osteosarcoma. Tumori, 2018, 104, 30-36.	1.1	84
13	Anthracycline, Gemcitabine, and Pazopanib in Epithelioid Sarcoma. JAMA Oncology, 2018, 4, e180219.	7.1	63
14	Activity of Pazopanib and Trabectedin in Advanced Alveolar Soft Part Sarcoma. Oncologist, 2018, 23, 62-70.	3.7	62
15	Results of the Scandinavian Sarcoma Group XIV protocol for classical osteosarcoma. Monthly Notices of the Royal Astronomical Society: Letters, 2011, 82, 211-216.	3.3	53
16	Pembrolizumab in advanced osteosarcoma: results of a single-arm, open-label, phase 2 trial. Cancer Immunology, Immunotherapy, 2021, 70, 2617-2624.	4.2	45
17	Young survivors of malignant bone tumours in the extremities: a comparative study of quality of life, fatigue and mental distress. Supportive Care in Cancer, 2007, 15, 1087-1096.	2.2	44
18	Recurrence-Free Survival After Resection of Gastric Gastrointestinal Stromal Tumors Classified According to a Strict Definition of Tumor Rupture: A Population-Based Study. Annals of Surgical Oncology, 2018, 25, 1133-1139.	1.5	40

#	Article	IF	CITATIONS
19	Highâ€dose chemotherapy with stem cell rescue in the primary treatment of metastatic and pelvic osteosarcoma: Final results of the ISC/SSG II study. Pediatric Blood and Cancer, 2014, 61, 840-845.	1.5	39
20	Needle biopsy through the abdominal wall for the diagnosis of gastrointestinal stromal tumour – Does it increase the risk for tumour cell seeding and recurrence?. European Journal of Cancer, 2016, 59, 128-133.	2.8	39
21	The Scandinavian Sarcoma Group Central Register: 6,000 patients after 25 years of monitoring of referral and treatment of extremity and trunk wall soft-tissue sarcoma. Monthly Notices of the Royal Astronomical Society: Letters, 2017, 88, 341-347.	3.3	38
22	Time dependence of prognostic factors for patients with soft tissue sarcoma. Cancer, 2004, 100, 2233-2239.	4.1	36
23	Prognostic Factors and Treatment Results of High-Grade Osteosarcoma in Norway: A Scope Beyond the "Classical―Patient. Sarcoma, 2015, 2015, 1-14.	1.3	36
24	Results of the second interim assessment of rEECur, an international randomized controlled trial of chemotherapy for the treatment of recurrent and primary refractory Ewing sarcoma (RR-ES) Journal of Clinical Oncology, 2020, 38, 11502-11502.	1.6	34
25	Protein expression of BIRC5, TK1, and TOP2A in malignant peripheral nerve sheath tumours – A prognostic test after surgical resection. Molecular Oncology, 2015, 9, 1129-1139.	4.6	32
26	Management of high-grade bone sarcomas over two decades: The Norwegian Radium Hospital experience. Acta OncolÃ ³ gica, 2006, 45, 38-46.	1.8	30
27	Small Round Cell Sarcomas. Seminars in Oncology, 2009, 36, 338-346.	2.2	28
28	Adjuvant chemotherapy and postoperative radiotherapy in high-risk soft tissue sarcoma patients defined by biological risk factors—A Scandinavian Sarcoma GroupÂstudy (SSG XX). European Journal of Cancer, 2018, 99, 78-85.	2.8	26
29	Denosumab in patients with giant-cell tumor of bone in Norway: results from a nationwide cohort. Acta Oncológica, 2017, 56, 479-483.	1.8	23
30	Chemotherapy in soft tissue sarcoma. Acta Orthopaedica, 1999, 70, 62-68.	1.4	20
31	Five-Year Results From a Scandinavian Sarcoma Group Study (SSG XIII) of Adjuvant Chemotherapy Combined With Accelerated Radiotherapy in High-Risk Soft Tissue Sarcoma of Extremities and Trunk Wall. International Journal of Radiation Oncology Biology Physics, 2011, 81, 1359-1366.	0.8	20
32	Results of the first interim assessment of rEECur, an international randomized controlled trial of chemotherapy for the treatment of recurrent and primary refractory Ewing sarcoma Journal of Clinical Oncology, 2019, 37, 11007-11007.	1.6	20
33	Methylated RASSF1A in malignant peripheral nerve sheath tumors identifies neurofibromatosis type 1 patients with inferior prognosis. Neuro-Oncology, 2015, 17, 63-69.	1.2	17
34	Expression of cell cycle regulators and frequency of TP53 mutations in high risk gastrointestinal stromal tumors prior to adjuvant imatinib treatment. PLoS ONE, 2018, 13, e0193048.	2.5	17
35	The Scandinavian Sarcoma Group. Monthly Notices of the Royal Astronomical Society: Letters, 2009, 80, 1-104.	3.3	16
36	Multimodal treatment of craniofacial osteosarcoma with high-grade histology. A single-center experience over 35Âyears. Neurosurgical Review, 2017, 40, 449-460.	2.4	16

KIRSTEN SUNDBY HALL

#	Article	IF	CITATIONS
37	DW MRI for evaluation of treatment response to imatinib in a rectal gastrointestinal stromal tumour. Acta Oncológica, 2011, 50, 148-150.	1.8	15
38	Prediction of long-term survival in patients with metastatic gastrointestinal stromal tumor: analysis of a large, single-institution cohort. Acta Oncológica, 2017, 56, 1317-1323.	1.8	15
39	MAP plus maintenance pegylated interferon α-2b (MAPIfn) versus MAP alone in patients with resectable high-grade osteosarcoma and good histologic response to preoperative MAP: First results of the EURAMOS-1 "good response―randomization Journal of Clinical Oncology, 2013, 31, LBA10504-LBA10504.	1.6	14
40	Referral patterns, treatment and outcome of highâ€grade malignant bone sarcoma in Scandinavia—SSG Central Register 25 years' experience. Journal of Surgical Oncology, 2015, 112, 853-860.	1.7	13
41	SLUG transcription factor: a pro-survival and prognostic factor in gastrointestinal stromal tumour. British Journal of Cancer, 2017, 116, 1195-1202.	6.4	13
42	High-dose cis-platinum combination chemotherapy in advanced nonseminomatous malignant germ cell tumours with emphasis on nephrotoxicity. Cancer Chemotherapy and Pharmacology, 1986, 18, 74-77.	2.3	12
43	Metastatic mesenteric dedifferentiated leiomyosarcoma: a case report and a review of literature. Clinical Sarcoma Research, 2016, 6, 2.	2.3	12
44	Inferior survival for patients with malignant peripheral nerve sheath tumors defined by aberrant TP53. Modern Pathology, 2018, 31, 1694-1707.	5.5	11
45	A randomized, double-blind, placebo-controlled, phase III study of crenolanib in advanced or metastatic GIST patients bearing a D842V mutation in <i>PDGFRA</i> : The CrenoGIST study Journal of Clinical Oncology, 2017, 35, TPS11080-TPS11080.	1.6	11
46	Clinical epidemiology and treatment outcomes of spindle cell non-osteogenic bone sarcomas – A nationwide population-based study. Journal of Bone Oncology, 2019, 14, 100207.	2.4	10
47	Striving towards Normality in Daily Life: A Qualitative Study of Patients Living with Metastatic Gastrointestinal Stromal Tumour in Long-Term Clinical Remission. Sarcoma, 2020, 2020, 1-9.	1.3	7
48	The Role of Adjuvant Treatment in Craniofacial Malignancy: A Critical Review. Frontiers in Oncology, 2020, 10, 1402.	2.8	7
49	The influence of partial hepatectomy on the pharmacokinetics of preoperatively injected 4′-epidoxorubicin in rats. Cancer Chemotherapy and Pharmacology, 1990, 26, 444-448.	2.3	4
50	Event-free survival and overall survival in 2,253 patients with osteosarcoma registered to EURAMOS-1 Journal of Clinical Oncology, 2015, 33, 10512-10512.	1.6	4
51	MAP plus maintenance pegylated interferon α-2b (MAP-IFN) versus MAP alone in patients (pts) with resectable high-grade osteosarcoma and good histologic response to preoperative MAP: First results of the EURAMOS-1 good response randomization Journal of Clinical Oncology, 2013, 31, LBA10504-LBA10504.	1.6	4
52	Verapamil Potentiates 4′â€Epidoxorubicin Cytotoxicity in a Rat Hepatoma Cell Line and in a Substrain Resistant to 4′â€Epidoxorubicin. Basic and Clinical Pharmacology and Toxicology, 1990, 67, 402-405.	0.0	3
53	Preoperative accelerated radiotherapy combined with chemotherapy in a defined cohort of patients with high risk soft tissue sarcoma: a Scandinavian Sarcoma Group study. Clinical Sarcoma Research, 2020, 10, 22.	2.3	3
54	EURAMOS-1 study: Recruitment, characteristics, and initial treatment of more than 2,000 patients (pts) with high-grade osteosarcoma Journal of Clinical Oncology, 2012, 30, 10081-10081.	1.6	3

#	Article	IF	CITATIONS
55	Fibrinogenâ€like protein 2 in gastrointestinal stromal tumour. Journal of Cellular and Molecular Medicine, 2022, 26, 1083-1094.	3.6	3
56	Validity and completeness of the Scandinavian Sarcoma Group Central Register by comparison with a nationwide cohort of patients with osteosarcoma in Norway. Journal of Surgical Oncology, 2018, 118, 246-247.	1.7	2
57	Real-world evidence on perioperative chemotherapy in localized soft tissue sarcoma of the extremities and trunk wall; a population-based study. Acta Oncológica, 2022, 61, 793-800.	1.8	1
58	Failure rate of standard rescue with leucovorin for high-dose methotrexate (HDMTX) in osteosarcoma Journal of Clinical Oncology, 2017, 35, 11028-11028.	1.6	0