

Fernando Rogério de Carvalho

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3418866/publications.pdf>

Version: 2024-02-01

30

papers

334

citations

840776

11

h-index

888059

17

g-index

30

all docs

30

docs citations

30

times ranked

356

citing authors

#	ARTICLE	IF	CITATIONS
1	<scp>NEOTROPICAL FRESHWATER FISHES</scp>: A dataset of occurrence and abundance of freshwater fishes in the Neotropics. <i>Ecology</i> , 2023, 104, e3713.	3.2	7
2	Broad and fine-scale threats on threatened Brazilian freshwater fish: variability across hydrographic regions and taxonomic groups. <i>Biota Neotropica</i> , 2021, 21, .	0.5	7
3	<i>Gymnotus paraguensis</i>, a Good Example of Phenotypic Plasticity in the Pantanal Biome, Brazil. <i>Zebrafish</i> , 2021, 18, 162-173.	1.1	1
4	Human impacts and the loss of Neotropical freshwater fish diversity. <i>Neotropical Ichthyology</i> , 2021, 19, .	1.0	14
5	Museum specimens reveal a rare new characid fish genus, helping to refine the interrelationships of the Probolodini (Characidae: Stethaprioninae). <i>Systematics and Biodiversity</i> , 2021, 19, 1135-1148.	1.2	0
6	Stream fish metacommunity organisation across a Neotropical ecoregion: The role of environment, anthropogenic impact and dispersal-based processes. <i>PLoS ONE</i> , 2020, 15, e0233733.	2.5	23
7	Stream fish from recently deforested basins in the Meridional Amazon, Mato Grosso, Brazil. <i>Biota Neotropica</i> , 2020, 20, .	0.5	3
8	Freshwater fish richness baseline from the SÃ£o Francisco Interbasin Water Transfer Project in the Brazilian Semi-arid. <i>Neotropical Ichthyology</i> , 2020, 18, .	1.0	5
9	First record of Jupiaba acanthogaster (Eigenmann, 1911) (Ostariophysi, Characidae) in the upper Paraná river basin, Brazil. <i>Check List</i> , 2020, 16, 889-893.	0.4	2
10	Taxonomic review of the Hyphessobrycon panamensis species-group (Characiformes: Characidae). <i>Zootaxa</i> , 2020, 4751, zootaxa.4751.3.1.	0.5	4
11	A new species of Astyanax Baird & Girard (Characiformes: Characidae) from western Chapada Diamantina, Bahia, Brazil. <i>Zootaxa</i> , 2019, 4604, zootaxa.4604.2.9.	0.5	9
12	Redescription of Hyphessobrycon itaparicensis, a senior synonym of H. sergipanus (Characiformes: Characidae). <i>Zootaxa</i> , 2019, 4751, zootaxa.4751.3.1.	1.0	0
13	New species of <i>Hyphessobrycon</i> from the Rio Teles Pires, Rio Tapajós basin, Brazil (Ostariophysi, Characidae). <i>Zootaxa</i> , 2019, 4604, zootaxa.4604.2.9.	1.6	0
14	Inventory of the freshwater fishes from a densely collected area in South America—a case study of the current knowledge of Neotropical fish diversity. <i>Zootaxa</i> , 2016, 4138, 401-40.	0.5	61
15	Redescription and osteology of Hyphessobrycon compressus (Meek) (Teleostei: Characidae), type species of the genus. <i>Neotropical Ichthyology</i> , 2015, 13, 513-540.	1.0	10
16	Taxonomic status of Tetragonopterus anomalus Steindachner, 1891, and inquirenda species in Characidae (Teleostei: Characiformes). <i>Zootaxa</i> , 2015, 4018, 146.	0.5	0
17	Redescription of Moenkhausia doceana (Steindachner, 1877) (Ostariophysi: Characidae): a characid from the Northeastern Mata Atlântica ecoregion, Brazil. <i>Neotropical Ichthyology</i> , 2014, 12, 377-388.	1.0	6
18	Redescription of Hyphessobrycon flammeus Myers, 1924 (Ostariophysi: Characidae), a threatened species from Brazil. <i>Neotropical Ichthyology</i> , 2014, 12, 247-256.	1.0	8

#	ARTICLE	IF	CITATIONS
19	Hypessobrycon uaiso: new characid fish from the rio Grande, upper rio Paraná basin, Minas Gerais State (Ostariophysi: Characidae), with a brief comment about some types of Hypessobrycon. <i>Neotropical Ichthyology</i> , 2013, 11, 525-536.	1.0	24
20	Revalidation of the genus Ectrepopterus Fowler (Teleostei: Characiformes), with the redescription of its type species, <i>E. uruguayensis</i> . <i>Zootaxa</i> , 2012, 3204, 47.	0.5	17
21	Geographic distribution of <i>Hemigrammus ora</i> (Ostariophysi: Characiformes: Characidae) in the Amazon basin, Brazil. <i>Zoologia</i> , 2011, 28, 545-550.	0.5	5
22	New species of <i>Moenkhausia</i> Eigenmann (Ostariophysi: Characidae) from the upper rio Tocantins basin in Central Brazil. <i>Neotropical Ichthyology</i> , 2011, 9, 57-63.	1.0	8
23	New species of <i>Hasemania</i> (Characiformes: Characidae) from Central Brazil, with comments on the endemism of upper rio Tocantins basin, Goiás State. <i>Neotropical Ichthyology</i> , 2010, 8, 27-32.	1.0	19
24	<i>Astyanax goyanensis</i> (Miranda-Ribeiro, 1944), new combination and <i>Astyanax courensis</i> , new species (Ostariophysi: Characiformes): two Characidae from the upper rio Tocantins basin, Central Brazil. <i>Neotropical Ichthyology</i> , 2010, 8, 265-275.	1.0	12
25	<i>Hemigrammus tocantinsi</i> : a new species from the upper rio Tocantins basin, Central Brazil (Characiformes: Characidae). <i>Neotropical Ichthyology</i> , 2010, 8, 247-254.	1.0	17
26	Fish assemblages in stream stretches occupied by cattail (Typhaceae, Angiospermae) stands in Southeast Brazil. <i>Neotropical Ichthyology</i> , 2009, 7, 241-250.	1.0	18
27	A new <i>Hemigrammus</i> Gill from upper rio Parana system, Southeastern Brazil (Characiformes): Tj ETQq1 1 0.784314 _{0.8} rgBT /Overlock 10 T		
28	Feeding association between benthic and nektonic Neotropical stream fishes. <i>Neotropical Ichthyology</i> , 2008, 6, 109-111.	1.0	11
29	Ichthyofauna of the rio Itatinga in the Parque das Neblinas, Bertioga, São Paulo State: composition and biogeography. <i>Biota Neotropica</i> , 2007, 7, 81-86.	1.0	31
30	Life history traits of parental care in <i>Crenicichla lepidota</i> (Cichliformes, Geophagini) in the upper Paraguay River basin, Brazil. <i>Papeis Avulsos De Zoologia</i> , 0, 61, e20216103.	0.4	0