

Hirohito Sone

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/3421735/hirohito-sone-publications-by-year.pdf>

Version: 2024-04-25

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

435
papers

13,600
citations

57
h-index

105
g-index

475
ext. papers

15,689
ext. citations

5.1
avg, IF

5.7
L-index

#	Paper	IF	Citations
435	Morphological and functional adaptation of pancreatic islet blood vessels to insulin resistance is impaired in diabetic db/db mice.. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2022 , 1868, 166339	6.9	1
434	Intensive oral care can reduce bloodstream infection with coagulase-negative staphylococci after neutrophil engraftment in allogeneic hematopoietic stem-cell transplantation. <i>Supportive Care in Cancer</i> , 2022 , 30, 475-485	3.9	0
433	Dietary intake and physical activity in Japanese patients with type 2 diabetes: the Japan Diabetes Complication and its Prevention prospective study (JDCP study 8).. <i>Diabetology International</i> , 2022 , 13, 344-357	2.3	
432	Differences in occupational stress by smoking intensity and gender in cross-sectional study of 59 355 Japanese employees using the Brief Job Stress Questionnaire (BJSQ): the Niigata Wellness Study.. <i>BMJ Open</i> , 2022 , 12, e055577	3	1
431	Severity of hypertension as a predictor of initiation of dialysis among study participants with and without diabetes mellitus. <i>Journal of Investigative Medicine</i> , 2021 , 69, 724-729	2.9	0
430	Network Meta-Analysis of Drug Therapies for Lowering Uric Acid and Mortality Risk in Patients with Heart Failure. <i>Cardiovascular Drugs and Therapy</i> , 2021 , 35, 1217-1225	3.9	1
429	Meta-analytic research of the dose-response relationship between salt intake and risk of heart failure. <i>Hypertension Research</i> , 2021 , 44, 885-887	4.7	
428	Accuracy of Japanese claims data in identifying diabetes-related complications. <i>Pharmacoepidemiology and Drug Safety</i> , 2021 , 30, 594-601	2.6	6
427	Physical Fitness and Dyslipidemia Among Japanese: A Cohort Study From the Niigata Wellness Study. <i>Journal of Epidemiology</i> , 2021 , 31, 287-296	3.4	3
426	Distinct effects of chondroitin sulfate on hematopoietic cells and the stromal microenvironment in bone marrow hematopoiesis. <i>Experimental Hematology</i> , 2021 , 96, 52-62.e5	3.1	2
425	Rates and risk factors for amputation in people with diabetes in Japan: a historical cohort study using a nationwide claims database. <i>Journal of Foot and Ankle Research</i> , 2021 , 14, 29	3.2	1
424	Starvation-induced transcription factor CREBH negatively governs body growth by controlling GH signaling. <i>FASEB Journal</i> , 2021 , 35, e21663	0.9	2
423	Dipeptidyl peptidase-4 inhibitor, anagliptin, alters hepatic insulin clearance in relation to the glycemic status in Japanese individuals with type2 diabetes. <i>Journal of Diabetes Investigation</i> , 2021 , 12, 1805-1815	3.9	0
422	Association of estimated plasma volume and weight loss after long-term administration and subsequent discontinuation of the sodium-glucose cotransporter-2 inhibitor tofogliflozin. <i>Diabetes, Obesity and Metabolism</i> , 2021 , 23, 1660-1665	6.7	3
421	Associations of Systolic Blood Pressure and Diastolic Blood Pressure With the Incidence of Coronary Artery Disease or Cerebrovascular Disease According to Glucose Status. <i>Diabetes Care</i> , 2021 , 44, 2124-2131	14.6	4
420	Body flexibility and incident hypertension: The Niigata wellness study. <i>Scandinavian Journal of Medicine and Science in Sports</i> , 2021 , 31, 702-709	4.6	2
419	A Prospective Cohort Study of Muscular and Performance Fitness and Risk of Hearing Loss: The Niigata Wellness Study. <i>American Journal of Medicine</i> , 2021 , 134, 235-242.e4	2.4	2

418	Enterohepatic Transcription Factor CREB3L3 Protects Atherosclerosis via SREBP Competitive Inhibition. <i>Cellular and Molecular Gastroenterology and Hepatology</i> , 2021 , 11, 949-971	7.9	3
417	WT1-specific CD8 + cytotoxic T cells with the capacity for antigen-specific expansion accumulate in the bone marrow in MDS. <i>International Journal of Hematology</i> , 2021 , 113, 723-734	2.3	0
416	A 52-week randomized controlled trial of ipragliflozin or sitagliptin in type 2 diabetes combined with metformin: The N-ISM study. <i>Diabetes, Obesity and Metabolism</i> , 2021 , 23, 811-821	6.7	3
415	Machine Learning Approach to Decision Making for Insulin Initiation in Japanese Patients With Type 2 Diabetes (JDDM 58): Model Development and Validation Study. <i>JMIR Medical Informatics</i> , 2021 , 9, e22148	3.6	2
414	Ability of Current Machine Learning Algorithms to Predict and Detect Hypoglycemia in Patients With Diabetes Mellitus: Meta-analysis. <i>JMIR Diabetes</i> , 2021 , 6, e22458	2.7	7
413	Le Carbone prevents liver damage in non-alcoholic steatohepatitis-hepatocellular carcinoma mouse model via AMPK/SIRT1 signaling pathway activation. <i>Heliyon</i> , 2021 , 7, e05888	3.6	1
412	Sodium Intake and Incidence of Diabetes Complications in Elderly Patients with Type 2 Diabetes-Analysis of Data from the Japanese Elderly Diabetes Intervention Study (J-EDIT). <i>Nutrients</i> , 2021 , 13,	6.7	2
411	Developing a health economic model for Asians with type 2 diabetes based on the Japan Diabetes Complications Study and the Japanese Elderly Diabetes Intervention Trial. <i>BMJ Open Diabetes Research and Care</i> , 2021 , 9,	4.5	2
410	Impact of prior cerebrovascular disease and glucose status on incident cerebrovascular disease in Japanese. <i>Cardiovascular Diabetology</i> , 2021 , 20, 174	8.7	0
409	Altered microbiota by a high-fat diet accelerates lethal myeloid hematopoiesis associated with systemic SOCS3 deficiency. <i>IScience</i> , 2021 , 24, 103117	6.1	
408	DA-EPOCH-R therapy for high-grade B-cell lymphoma with MYC and BCL2 and/or BCL6 rearrangements in a patient with renal dysfunction. <i>Journal of Clinical and Experimental Hematopathology: JCEH</i> , 2021 , 61, 42-47	1.9	1
407	Successful Treatment with Edoxaban for Disseminated Intravascular Coagulation in a Case of Aortic Dissection Complicated with Immune Thrombocytopenic Purpura. <i>Internal Medicine</i> , 2020 , 59, 2035-2039	1.1	0
406	Brain adaptations of insulin signaling kinases, GLUT 3, p-BADser155 and nitrotyrosine expression in various hypoglycemic models of mice. <i>Neurochemistry International</i> , 2020 , 137, 104745	4.4	2
405	Association between Low Protein Intake and Mortality in Patients with Type 2 Diabetes. <i>Nutrients</i> , 2020 , 12,	6.7	7
404	Skipping breakfast, late-night eating and current smoking are associated with medication adherence in Japanese patients with diabetes. <i>Primary Care Diabetes</i> , 2020 , 14, 753-759	2.4	3
403	CREBH Improves Diet-Induced Obesity, Insulin Resistance, and Metabolic Disturbances by FGF21-Dependent and FGF21-Independent Mechanisms. <i>IScience</i> , 2020 , 23, 100930	6.1	8
402	Prevention of postprandial hypotension-related syncope by caffeine in a patient with long-standing diabetes mellitus. <i>Endocrine Journal</i> , 2020 , 67, 585-592	2.9	0
401	Influence of an SGLT2 inhibitor, tofogliflozin, on the resting heart rate in relation to adipose tissue insulin resistance. <i>Diabetic Medicine</i> , 2020 , 37, 1316-1325	3.5	5

400	Clinicopathological analysis of splenic red pulp low-grade B-cell lymphoma. <i>Pathology International</i> , 2020 , 70, 280-286	1.8	3
399	Association of increased hepatic insulin clearance and change in serum triglycerides or hydroxybutyrate concentration via the sodium/glucose-cotransporter 2 inhibitor tofogliflozin. <i>Diabetes, Obesity and Metabolism</i> , 2020 , 22, 947-956	6.7	6
398	The Glasgow prognostic score divides high-risk hematopoietic cell transplantation-specific comorbidity index patients into stratified subgroups in allogeneic hematopoietic cell transplantation. <i>Annals of Hematology</i> , 2020 , 99, 671-673	3	
397	Genetic Manipulation Resulting in Decreased Donor Chondroitin sulfate Synthesis Mitigates Gvhd Following Allogeneic Hematopoietic Cell Transplantation in a Murine Model.. <i>Blood</i> , 2020 , 136, 25-26	2.2	
396	Applications of physical performance measures to routine diabetes care for frailty prevention concept: fundamental data with grip strength, gait speed, timed chair stand speed, standing balance, and knee extension strength. <i>BMJ Open Diabetes Research and Care</i> , 2020 , 8,	4.5	5
395	A Prospective Cohort Study of Muscular and Performance Fitness and Incident Glaucoma: The Niigata Wellness Study. <i>Journal of Physical Activity and Health</i> , 2020 , 17, 1171-1178	2.5	2
394	6-OR: Blood Pressure as Predictor of Coronary Artery Disease (CAD)/Cerebrovascular Disease (CVD) According to Glucose Tolerance Status (GTS): Implications for Updated Guidelines. <i>Diabetes</i> , 2020 , 69, 6-OR	0.9	1
393	518-P: Association between Plasma Metabolites in the Urea Cycle and Diabetic Kidney Disease (DKD) in a Japanese Population. <i>Diabetes</i> , 2020 , 69, 518-P	0.9	
392	1133-P: Association of Higher Baseline BNP Levels with a Greater Reduction in Plasma Volume and Increase in Beta-Hydroxybutyrate via the SGLT2 Inhibitor Tofogliflozin in Type 2 Diabetes. <i>Diabetes</i> , 2020 , 69, 1133-P	0.9	
391	841-P: Predictive Ability of Incident Type 2 Diabetes Mellitus (T2DM) Using Machine Learning Algorithms: A Meta-analysis. <i>Diabetes</i> , 2020 , 69, 841-P	0.9	
390	449-P: Impact of Body Mass Index (BMI) and Waist Circumference (WC) on Coronary Artery Disease (CAD) in Japanese with and without Diabetes Mellitus (DM). <i>Diabetes</i> , 2020 , 69, 449-P	0.9	2
389	1125-P: Association of Plasma Volume with Body Weight and BNP after Long-Term Administration and Subsequent Withdrawal of the SGLT2 Inhibitor Tofogliflozin. <i>Diabetes</i> , 2020 , 69, 1125-P	0.9	
388	1513-P: Severity of Hypertension (HT) as a Predictor of Initiation of Dialysis among Study Participants with and without Diabetes Mellitus (DM). <i>Diabetes</i> , 2020 , 69, 1513-P	0.9	
387	1126-P: Randomized Controlled Trial of Ipragliflozin or Sitagliptin Combined with Metformin in Type 2 Diabetes: NISM Study. <i>Diabetes</i> , 2020 , 69, 1126-P	0.9	
386	1267-P: Physical Fitness (PF), Weight Status, and Metabolic Risk in Japanese Adolescents. <i>Diabetes</i> , 2020 , 69, 1267-P	0.9	
385	1988-P: Association between Screen Time Including Smartphone Use and Obesity and Its Associated Lifestyles in School Children in Japan. <i>Diabetes</i> , 2020 , 69, 1988-P	0.9	
384	1553-P: Combination of Diabetes Mellitus and Lack of Habitual Physical Activity Is a Risk Factor for Functional Disability in Japanese. <i>Diabetes</i> , 2020 , 69, 1553-P	0.9	
383	389-P: Ability for Detecting or Predicting Hypoglycemia with the Aid of Machine Learning Techniques: A Meta-analysis. <i>Diabetes</i> , 2020 , 69, 389-P	0.9	

382	1903-P: Low Level of and Decrease in Serum Amylase (AMY) Increases the Risk of Type 2 Diabetes Mellitus (T2DM). <i>Diabetes</i> , 2020 , 69, 1903-P	0.9	0
381	663-P: Medication Adherence (MA) Is an Independent Predictor for Glycemic Control Even after Adjustment for Lifestyle Confounders in Japanese Patients with Type 2 Diabetes. <i>Diabetes</i> , 2020 , 69, 663-P	0.9	
380	A Prospective Cohort Study Of Physical Fitness And Incident Hearing Loss: The Niigata Wellness Study. <i>Medicine and Science in Sports and Exercise</i> , 2020 , 52, 421-421	1.2	
379	Efficacy and safety of empagliflozin as add-on to insulin in Japanese patients with type 2 diabetes: A randomized, double-blind, placebo-controlled trial. <i>Diabetes, Obesity and Metabolism</i> , 2020 , 22, 417-426	6.7	7
378	Diabetes mellitus and risk of new-onset and recurrent heart failure: a systematic review and meta-analysis. <i>ESC Heart Failure</i> , 2020 , 7, 2146-2174	3.7	13
377	Marker chromosome is a strong poor prognosis factor after allogeneic HSCT for adverse-risk AML patients. <i>European Journal of Haematology</i> , 2020 , 105, 616-625	3.8	1
376	Association of treatment-achieved HbA with incidence of coronary artery disease and severe eye disease in diabetes patients. <i>Diabetes and Metabolism</i> , 2020 , 46, 331-334	5.4	2
375	Vitamin B6 intake and incidence of diabetic retinopathy in Japanese patients with type 2 diabetes: analysis of data from the Japan Diabetes Complications Study (JDACS). <i>European Journal of Nutrition</i> , 2020 , 59, 1585-1594	5.2	5
374	Hepatocyte ELOVL Fatty Acid Elongase 6 Determines Ceramide Acyl-Chain Length and Hepatic Insulin Sensitivity in Mice. <i>Hepatology</i> , 2020 , 71, 1609-1625	11.2	15
373	Combination of diabetes mellitus and lack of habitual physical activity is a risk factor for functional disability in Japanese. <i>BMJ Open Diabetes Research and Care</i> , 2020 , 8,	4.5	2
372	Weight and cardiometabolic risk among adolescents in Agano city, Japan: NICE EVIDENCE Study-Agano 1. <i>Asia Pacific Journal of Clinical Nutrition</i> , 2020 , 29, 856-866	1	
371	Relationship Between Number of Multiple Risk Factors and Coronary Artery Disease Risk With and Without Diabetes Mellitus. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2019 , 104, 5084-5090	5.6	12
370	Association between Helicobacter pylori infection, eradication and diabetes mellitus. <i>Journal of Diabetes Investigation</i> , 2019 , 10, 1341-1346	3.9	8
369	Combined Effects of Energy Intake and Physical Activity on Obesity in Japanese Patients with Type 2 Diabetes (JDDM 50): A Cross-Sectional Study. <i>Diabetes Therapy</i> , 2019 , 10, 1133-1138	3.6	1
368	Pulse Pressure is a Stronger Predictor Than Systolic Blood Pressure for Severe Eye Diseases in Diabetes Mellitus. <i>Journal of the American Heart Association</i> , 2019 , 8, e010627	6	9
367	Anaplastic large cell lymphoma, with 1,25(OH)D-mediated hypercalcemia: A case report. <i>Journal of Clinical and Experimental Hematopathology: JCEH</i> , 2019 , 59, 22-28	1.9	2
366	Physical Fitness Tests and Type 2 Diabetes Among Japanese: A Longitudinal Study From the Niigata Wellness Study. <i>Journal of Epidemiology</i> , 2019 , 29, 139-146	3.4	27
365	Patient-based prediction algorithm of relapse after allo-HSCT for acute Leukemia and its usefulness in the decision-making process using a machine learning approach. <i>Cancer Medicine</i> , 2019 , 8, 5058-5067	4.8	15

364	Attenuation of Weight Loss Through Improved Antilipolytic Effect in Adipose Tissue Via the SGLT2 Inhibitor Tofogliflozin. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2019 , 104, 3647-3660	5.6	12
363	Overt Proteinuria, Moderately Reduced eGFR and Their Combination Are Predictive of Severe Diabetic Retinopathy or Diabetic Macular Edema in Diabetes 2019 , 60, 2685-2689		9
362	Risk of coronary artery disease according to glucose abnormality status and prior coronary artery disease in Japanese men. <i>Metabolism: Clinical and Experimental</i> , 2019 , 101, 153991	12.7	8
361	Isolated Adrenocorticotrophic Hormone Deficiency Presenting with Severe Hyponatremia and Rhabdomyolysis: A Case Report and Literature Review. <i>American Journal of Case Reports</i> , 2019 , 20, 1857-1863	1.3	0
360	767-P: Family Support for Medical Nutritional Therapy and Dietary Intake among Japanese Patients with Type 2 Diabetes. <i>Diabetes</i> , 2019 , 68, 767-P	0.9	1
359	634-P: Poor Adherence to Medication and HbA1c Level Predict Risk of Amputation in Patients with Diabetes Mellitus: Historical Cohort Study Using a Nationwide Claims Database. <i>Diabetes</i> , 2019 , 68, 634-P	0.9	
358	1335-P: Weight Status and Cardiometabolic Risk Factors among Adolescents in Japan. <i>Diabetes</i> , 2019 , 68, 1335-P	0.9	
357	1579-P: Higher Calcium Intake Is Associated with Lower Incidence of Diabetic Nephropathy in Japanese Patients with Type 2 Diabetes. <i>Diabetes</i> , 2019 , 68, 1579-P	0.9	
356	1219-P: Enhanced Response in Serum Ketone Level in Men Compared with Women by Administration of SGLT2 Inhibitor. <i>Diabetes</i> , 2019 , 68, 1219-P	0.9	
355	1205-P: Baseline Lactate Level Is a Useful Predictor for Weight Loss after Long-Term SGLT2 Inhibitor Treatment. <i>Diabetes</i> , 2019 , 68, 1205-P	0.9	
354	2450-PUB: Association between Intake of Magnesium and Obesity in Japanese Patients with Type 2 Diabetes Mellitus. <i>Diabetes</i> , 2019 , 68, 2450-PUB	0.9	
353	598-P: Lower Hematocrit Is Predictive of Treatment-Required Eye Diseases in Japanese Patients with Diabetes Mellitus. <i>Diabetes</i> , 2019 , 68, 598-P	0.9	
352	698-P: Factors Significantly Associated with Adherence to Diabetes Medications: Findings from a Large Japanese Claims Database. <i>Diabetes</i> , 2019 , 68, 698-P	0.9	
351	450-P: Impact of Prior Coronary Artery Disease (CAD)/Cerebrovascular Disease (CVD) and Diabetes Mellitus (DM) on Incident CAD/CVD in Japanese. <i>Diabetes</i> , 2019 , 68, 450-P	0.9	
350	1583-P: Intake of Fish and Related Nutrients in Association with Obesity in Japanese Patients with Type 2 Diabetes (T2D). <i>Diabetes</i> , 2019 , 68, 1583-P	0.9	
349	775-P: Significant Association of Food Group Intake with Obesity among Patients with Type 2 Diabetes Mellitus in Japan. <i>Diabetes</i> , 2019 , 68, 775-P	0.9	
348	448-P: Predictors and Their Impact on Coronary Artery Disease (CAD) According to Glucose Tolerance Status (GTS) and Prior CAD: Historical Cohort Study in Japan. <i>Diabetes</i> , 2019 , 68, 448-P	0.9	
347	1478-P: Impact of Prior Cerebrovascular Disease (CVD) and Glucose Tolerance Status on Incident CVD in Japanese. <i>Diabetes</i> , 2019 , 68, 1478-P	0.9	

346	1215-P: Adipose Tissue Insulin Resistance Predicts Ketosis via an SGLT2 Inhibitor. <i>Diabetes</i> , 2019 , 68, 1215-P	0.9	
345	1910-P: SOCS3 Deficiency on a High-Fat Diet Accelerates Systemic Inflammation and Results in Lethal Myeloid Hematopoiesis without Obesity and Adiposity. <i>Diabetes</i> , 2019 , 68, 1910-P	0.9	
344	446-P: Associations of Blood Pressure (BP) with Incidence of Coronary Artery Disease (CAD)/Cerebrovascular Disease (CVD) According to Glucose Tolerance Status (GTS). <i>Diabetes</i> , 2019 , 68, 446-P	0.9	
343	A Prospective Cohort Study of Physical Fitness and Incident Glaucoma: The Niigata Wellness Study. <i>Medicine and Science in Sports and Exercise</i> , 2019 , 51, 222-222	1.2	
342	789-P: Association of Zinc Intake with Obesity in Japanese Patients with Type 2 Diabetes Mellitus (T2DM). <i>Diabetes</i> , 2019 , 68, 789-P	0.9	1
341	764-P: Low Dietary Energy Density (DED) Diet Is Associated with Favorable Dietary Pattern in Japanese Patients with Type 2 Diabetes (T2DM). <i>Diabetes</i> , 2019 , 68, 764-P	0.9	
340	1514-P: Serum Amylase (AMY) Level and Its Chronological Change as a Predictor of Incident Type 2 Diabetes Mellitus (T2DM). <i>Diabetes</i> , 2019 , 68, 1514-P	0.9	
339	854-P: Personality, Self-Management Behaviors, and Glycemic Control among Japanese Patients with Type 2 Diabetes Mellitus (T2DM). <i>Diabetes</i> , 2019 , 68, 854-P	0.9	
338	Depletion of Pre-Transplant Skeletal Muscle Is a Significant Poor Prognostic Factor in Allogeneic Hematopoietic Cell Transplantation. <i>Blood</i> , 2019 , 134, 3322-3322	2.2	
337	Safety and effective salvage regimen comprising a novel combination of brentuximab vedotin, L-asparaginase, and dexamethasone for refractory anaplastic large cell lymphoma, anaplastic lymphoma kinase negative. <i>Hematological Oncology</i> , 2019 , 37, 212-214	1.3	1
336	Effect of family-oriented diabetes programs on glycemic control: A meta-analysis. <i>Family Practice</i> , 2019 , 36, 387-394	1.9	3
335	Higher pulse pressure predicts initiation of dialysis in Japanese patients with diabetes. <i>Diabetes/Metabolism Research and Reviews</i> , 2019 , 35, e3120	7.5	6
334	Gemcitabine, Dexamethasone, and Cisplatin Regimen as an Effective Salvage Therapy for High-grade B-cell Lymphoma with MYC and BCL2 and/or BCL6 Rearrangements. <i>Internal Medicine</i> , 2019 , 58, 575-580	1.1	0
333	Meat intake and incidence of cardiovascular disease in Japanese patients with type 2 diabetes: analysis of the Japan Diabetes Complications Study (JDACS). <i>European Journal of Nutrition</i> , 2019 , 58, 281-290	5.3	9
332	Cardiovascular Disease in Japanese Patients with Type 2 Diabetes Mellitus. <i>Annals of Vascular Diseases</i> , 2018 , 11, 2-14	0.9	4
331	Expression of programmed death ligand 1 is associated with poor prognosis in myeloid sarcoma patients. <i>Hematological Oncology</i> , 2018 , 36, 591-599	1.3	7
330	Relationships among cardiorespiratory fitness, muscular fitness, and cardiometabolic risk factors in Japanese adolescents: Niigata screening for and preventing the development of non-communicable disease study-Agano (NICE EVIDENCE Study-Agano) 2. <i>Pediatric Diabetes</i> , 2018 , 19, 593-602	3.6	7
329	Role of fatty liver in the association between obesity and reduced hepatic insulin clearance. <i>Diabetes and Metabolism</i> , 2018 , 44, 135-142	5.4	12

328	MAGI-1 expression is decreased in several types of human T-cell leukemia cell lines, including adult T-cell leukemia. <i>International Journal of Hematology</i> , 2018 , 107, 337-344	2.3	9
327	Frequent expression of CD30 in extranodal NK/T-cell lymphoma: Potential therapeutic target for anti-CD30 antibody-based therapy. <i>Hematological Oncology</i> , 2018 , 36, 166-173	1.3	17
326	Quantitative Relationship Between Cumulative Risk Alleles Based on Genome-Wide Association Studies and Type 2 Diabetes Mellitus: A Systematic Review and Meta-analysis. <i>Journal of Epidemiology</i> , 2018 , 28, 3-18	3.4	8
325	Relationship between intake of fruit separately from vegetables and triglycerides - A meta-analysis. <i>Clinical Nutrition ESPEN</i> , 2018 , 27, 53-58	1.3	5
324	Refinement of the Glasgow Prognostic Score as a pre-transplant risk assessment for allogeneic hematopoietic cell transplantation. <i>International Journal of Hematology</i> , 2018 , 108, 282-289	2.3	3
323	Transgenic Mice Overexpressing SREBP-1a in Male ob/ob Mice Exhibit Lipodystrophy and Exacerbate Insulin Resistance. <i>Endocrinology</i> , 2018 , 159, 2308-2323	4.8	11
322	Glutamic Acid Decarboxylase Autoantibody-negative Slowly Progressive Type 1 Diabetes Mellitus: A Case Report and Literature Review. <i>Internal Medicine</i> , 2018 , 57, 3581-3587	1.1	
321	Influence of SGLT2 Inhibitor on Resting Heart Rate (RHR) and Factors Related to Its Changes. <i>Diabetes</i> , 2018 , 67, 1154-P	0.9	4
320	Higher Dietary Intake of Vitamin D Is Associated with Lower Incidence of Diabetic Nephropathy in Japanese Patients with Type 2 Diabetes. <i>Diabetes</i> , 2018 , 67, 1561-P	0.9	1
319	Test Strip Positive Proteinuria and Its Combination with Low eGFR Are Predictive of Treatment-Required Eye Diseases in Japanese Patients with Diabetes Mellitus. <i>Diabetes</i> , 2018 , 67, 594-P ^{0.9}		
318	Genetically Reduced Chondroitin Sulfate Prevents the Progression of Diabetic Neuropathy. <i>Diabetes</i> , 2018 , 67, 547-P	0.9	
317	Impact of Prior Coronary Artery Disease (CAD) and Glucose Tolerance Status (GTS) on Incident CAD in Japanese Men. <i>Diabetes</i> , 2018 , 67, 1488-P	0.9	
316	Impact of Vitamin B6 Intake on the Risk of Diabetic Retinopathy: Analysis from Multicenter Prospective Study of Japanese Patients with Type 2 Diabetes. <i>Diabetes</i> , 2018 , 67, 597-P	0.9	
315	Impact of Carbohydrate Intake on Obesity in Japanese Patients with Type 2 Diabetes: An Analysis of the JDCP Registry. <i>Diabetes</i> , 2018 , 67, 787-P	0.9	
314	Dipstick Proteinuria as a Predictor of End-Stage Renal Disease in Japanese Adults With and Without Diabetes Mellitus (DM). <i>Diabetes</i> , 2018 , 67, 2384-PUB	0.9	
313	Incidence and Risk Factors for Amputation in Patients with Diabetes in Japan: Historical Cohort Study Using a Nationwide Claims Database. <i>Diabetes</i> , 2018 , 67, 637-P	0.9	0
312	Education for Family Members Is Effective for Improved Glycemic Control of Patients with Type 2 Rather than Type 1 Diabetes Mellitus: A Meta-analysis. <i>Diabetes</i> , 2018 , 67, 823-P	0.9	
311	Effects of Treatment-Achieved HbA1c on Incidence of Micro-/Macrovascular Complications in Patients with Diabetes Mellitus. <i>Diabetes</i> , 2018 , 67, 444-P	0.9	

310	Effect of Number of Achieved Targets for Risk Factors on Coronary Artery Disease (CAD) in Those With and Without Diabetes Mellitus (DM). <i>Diabetes</i> , 2018 , 67, 442-P	0.9	
309	Developing a Health Economic Model for Asians with Type 2 Diabetes Based on the Japan Diabetes Complications Study and the Japanese Elderly Diabetes Intervention Trial. <i>Diabetes</i> , 2018 , 67, 2319-PUB ^{0.9}		
308	Association between Serum Amylase Level and Incidence of Type 2 Diabetes. <i>Diabetes</i> , 2018 , 67, 2368-PUB		
307	Reduced Postprandial Hepatic Insulin Clearance via the DPP-4 Inhibitor Anagliptin Contributed to Improvement in Hyperglycemia in Patients with Type 2 Diabetes Mellitus. <i>Diabetes</i> , 2018 , 67, 1176-P	0.9	
306	Distinct Effects of Chondroitin Sulfate on Hematopoietic Cells and the Stromal Microenvironment in Bone Marrow Hematopoiesis. <i>Blood</i> , 2018 , 132, 3852-3852	2.2	1
305	Marker Chromosomes Are a New Cytogenetic Adverse Risk Factor in AML after Allo-HCT. <i>Blood</i> , 2018 , 132, 5260-5260	2.2	
304	Clinical Features and Risk Factors of Post-Engraftment Bloodstream Infection in Allogeneic HCT. <i>Blood</i> , 2018 , 132, 5712-5712	2.2	
303	A distinct subtype of Epstein-Barr virus-positive T/NK-cell lymphoproliferative disorder: adult patients with chronic active Epstein-Barr virus infection-like features. <i>Haematologica</i> , 2018 , 103, 1018-1028	6.6	3 ¹
302	Predictors of the response of HbA1c and body weight after SGLT2 inhibition. <i>Diabetes and Metabolism</i> , 2018 , 44, 172-174	5.4	7
301	Cladribine treatment for Erdheim-Chester disease involving the central nervous system and concomitant polycythemia vera: A case report. <i>Journal of Clinical and Experimental Hematopathology: JCEH</i> , 2018 , 58, 161-165	1.9	5
300	Heterogeneity of intrahepatic iron deposition in transfusion-dependent iron overload patients with hematological malignancies. <i>Leukemia Research</i> , 2018 , 70, 41-44	2.7	1
299	Unstable bodyweight and incident type 2 diabetes mellitus: A meta-analysis. <i>Journal of Diabetes Investigation</i> , 2017 , 8, 501-509	3.9	16
298	Potential protective effect of lactation against incidence of type 2 diabetes mellitus in women with previous gestational diabetes mellitus: A systematic review and meta-analysis. <i>Diabetes/Metabolism Research and Reviews</i> , 2017 , 33, e2875	7.5	37
297	Comparison of clinicopathological characteristics between T-cell prolymphocytic leukemia and peripheral T-cell lymphoma, not otherwise specified. <i>European Journal of Haematology</i> , 2017 , 98, 459-466	2.8	10
296	Elovl6 Deficiency Improves Glycemic Control in Diabetic / Mice by Expanding βCell Mass and Increasing Insulin Secretory Capacity. <i>Diabetes</i> , 2017 , 66, 1833-1846	0.9	15
295	Association between all-cause mortality and severity of depressive symptoms in patients with type 2 diabetes: Analysis from the Japan Diabetes Complications Study (JDACS). <i>Journal of Psychosomatic Research</i> , 2017 , 99, 34-39	4.1	7
294	Carbohydrate intake during early pregnancy is inversely associated with abnormal glucose challenge test results in Japanese pregnant women. <i>Diabetes/Metabolism Research and Reviews</i> , 2017 , 33, e2898	7.5	6
293	Clinicopathological features of cryptococcal lymphadenitis and a review of literature. <i>Journal of Clinical and Experimental Hematopathology: JCEH</i> , 2017 , 57, 26-30	1.9	6

292	Impact of body mass index and metabolic phenotypes on coronary artery disease according to glucose tolerance status. <i>Diabetes and Metabolism</i> , 2017 , 43, 543-546	5.4	12
291	Internal deletion of BCOR reveals a tumor suppressor function for BCOR in T lymphocyte malignancies. <i>Journal of Experimental Medicine</i> , 2017 , 214, 2901-2913	16.6	35
290	Successful 5-azacytidine treatment of myeloid sarcoma and leukemia cutis associated with myelodysplastic syndrome: A case report and literature review. <i>Medicine (United States)</i> , 2017 , 96, e7975 ^{1.8}	1.8	10
289	Functional dissection of hematopoietic stem cell populations with a stemness-monitoring system based on NS-GFP transgene expression. <i>Scientific Reports</i> , 2017 , 7, 11442	4.9	8
288	Comparative evaluation of torasemide and spironolactone on adverse cardiac remodeling in a rat model of dilated cardiomyopathy. <i>Cardiovascular Therapeutics</i> , 2017 , 35, e12283	3.3	
287	Severe hypoglycaemia is a major predictor of incident diabetic retinopathy in Japanese patients with type 2 diabetes. <i>Diabetes and Metabolism</i> , 2017 , 43, 424-429	5.4	3
286	Advanced glycation end products induce brain-derived neurotrophic factor release from human platelets through the Src-family kinase activation. <i>Cardiovascular Diabetology</i> , 2017 , 16, 20	8.7	10
285	Comparison of baseline characteristics and clinical course in Japanese patients with type 2 diabetes among whom different types of oral hypoglycemic agents were chosen by diabetes specialists as initial monotherapy (JDDM 42). <i>Medicine (United States)</i> , 2017 , 96, e6122	1.8	15
284	Patients with type 2 diabetes having higher glomerular filtration rate showed rapid renal function decline followed by impaired glomerular filtration rate: Japan Diabetes Complications Study. <i>Journal of Diabetes and Its Complications</i> , 2017 , 31, 473-478	3.2	18
283	Comparative effects of torasemide and furosemide on gap junction proteins and cardiac fibrosis in a rat model of dilated cardiomyopathy. <i>BioFactors</i> , 2017 , 43, 187-194	6.1	4
282	Impact of glucose tolerance status on the development of coronary artery disease among working-age men. <i>Diabetes and Metabolism</i> , 2017 , 43, 261-264	5.4	18
281	The Glasgow Prognostic Score as a pre-transplant risk assessment for allogeneic hematopoietic cell transplantation. <i>Clinical Transplantation</i> , 2017 , 31, e13103	3.8	4
280	Is the Proportion of Carbohydrate Intake Associated with the Incidence of Diabetes Complications?-An Analysis of the Japan Diabetes Complications Study. <i>Nutrients</i> , 2017 , 9,	6.7	8
279	Dietary salt intake and diabetes complications in patients with diabetes: An overview. <i>Journal of General and Family Medicine</i> , 2017 , 18, 16-20	1	3
278	Utility of nonblood-based risk assessment for predicting type 2 diabetes mellitus: A meta-analysis. <i>Preventive Medicine</i> , 2016 , 91, 180-187	4.3	2
277	Hyperlipidemia and hepatitis in liver-specific CREB3L3 knockout mice generated using a one-step CRISPR/Cas9 system. <i>Scientific Reports</i> , 2016 , 6, 27857	4.9	25
276	Impact of individual components and their combinations within a family history of hypertension on the incidence of hypertension: Toranomon hospital health management center study 22. <i>Medicine (United States)</i> , 2016 , 95, e4564	1.8	9
275	Clinicopathological, Cytogenetic, and Prognostic Analysis of 131 Myeloid Sarcoma Patients. <i>American Journal of Surgical Pathology</i> , 2016 , 40, 1473-1483	6.7	54

274	The SIL index is a simple and objective prognostic indicator in diffuse large B-cell lymphoma. <i>Leukemia and Lymphoma</i> , 2016 , 57, 2763-2770	1.9	9
273	Comparison of lipid parameters to predict cardiovascular events in Japanese mild-to-moderate hypercholesterolemic patients with and without type 2 diabetes: Subanalysis of the MEGA study. <i>Diabetes Research and Clinical Practice</i> , 2016 , 113, 14-22	7.4	14
272	Development and evaluation of the Japanese version of the Audit of Diabetes-Dependent Quality of Life for patients with diabetes. <i>Diabetology International</i> , 2016 , 7, 384-390	2.3	4
271	Assessment of kidney dysfunction with cystatin C- and creatinine-based estimated glomerular filtration rate and predicting type 2 diabetes: Toranomon Hospital Health Management Center Study 21. <i>Diabetes Research and Clinical Practice</i> , 2016 , 113, 60-8	7.4	
270	In Search of the Ideal Resistance Training Program to Improve Glycemic Control and its Indication for Patients with Type 2 Diabetes Mellitus: A Systematic Review and Meta-Analysis. <i>Sports Medicine</i> , 2016 , 46, 67-77	10.6	45
269	Different Effects of Eicosapentaenoic and Docosahexaenoic Acids on Atherogenic High-Fat Diet-Induced Non-Alcoholic Fatty Liver Disease in Mice. <i>PLoS ONE</i> , 2016 , 11, e0157580	3.7	40
268	Evaluation of Liver Iron Deposition in Transfusion-Dependent Patients By Dual-Energy CT. <i>Blood</i> , 2016 , 128, 3619-3619	2.2	
267	The Predictive Factors of Favorable Prognosis after Allo-HSCT for Refractory Acute Leukemia. <i>Blood</i> , 2016 , 128, 2297-2297	2.2	
266	Meta-analytic research on the relationship between cumulative risk alleles and risk of type 2 diabetes mellitus. <i>Diabetes/Metabolism Research and Reviews</i> , 2016 , 32, 178-86	7.5	1
265	The Dinakara equation for adjusting DLCO for hemoglobin in the HCT-CI is superior to the Cotes equation for predicting high-risk patients in allogeneic hematopoietic stem cell transplantation. <i>American Journal of Hematology</i> , 2016 , 91, E296	7.1	1
264	Fulminant Type 1 Diabetes Mellitus Associated with Coxsackie Virus Type A2 Infection: A Case Report and Literature Review. <i>Internal Medicine</i> , 2016 , 55, 643-6	1.1	16
263	Fibromyalgia in a Patient with Cushing's Disease Accompanied by Central Hypothyroidism. <i>Internal Medicine</i> , 2016 , 55, 3185-3190	1.1	1
262	Rapid Normalization of High Glutamic Acid Decarboxylase Autoantibody Titers and Preserved Endogenous Insulin Secretion in a Patient with Diabetes Mellitus: A Case Report and Literature Review. <i>Internal Medicine</i> , 2016 , 55, 485-9	1.1	1
261	CREB3L3 controls fatty acid oxidation and ketogenesis in synergy with PPAR α . <i>Scientific Reports</i> , 2016 , 6, 39182	4.9	27
260	Association of eating three meals irregularly with changes in BMI and weight among young Japanese men and women: A 2-year follow-up. <i>Physiology and Behavior</i> , 2016 , 163, 81-87	3.5	4
259	Maximum BMI and microvascular complications in a cohort of Japanese patients with type 2 diabetes: the Japan Diabetes Complications Study. <i>Journal of Diabetes and Its Complications</i> , 2016 , 30, 790-7	3.2	5
258	Intestinal CREBH overexpression prevents high-cholesterol diet-induced hypercholesterolemia by reducing expression. <i>Molecular Metabolism</i> , 2016 , 5, 1092-1102	8.8	19
257	Quantitative assessment of genetic testing for type 2 diabetes mellitus based on findings of genome-wide association studies. <i>Annals of Epidemiology</i> , 2016 , 26, 816-818.e6	6.4	0

256	Cluster-randomized trial to improve the quality of diabetes management: The study for the efficacy assessment of the standard diabetes manual (SEAS-DM). <i>Journal of Diabetes Investigation</i> , 2016 , 7, 539-43	3.9	5
255	MYC translocation and/or BCL 2 protein expression are associated with poor prognosis in diffuse large B-cell lymphoma. <i>Cancer Science</i> , 2016 , 107, 853-61	6.9	32
254	Comparison of clinical characteristics in patients with type 2 diabetes among whom different antihyperglycemic agents were prescribed as monotherapy or combination therapy by diabetes specialists. <i>Journal of Diabetes Investigation</i> , 2016 , 7, 260-9	3.9	10
253	Correlation between SNP genotypes and periodontitis in Japanese type II diabetic patients: a preliminary study. <i>Odontology / the Society of the Nippon Dental University</i> , 2015 , 103, 233-40	3.6	2
252	Risk factors associated with abnormal cognition in Japanese outpatients with diabetes, hypertension or dyslipidemia. <i>Diabetology International</i> , 2015 , 6, 268-274	2.3	15
251	Fasting glucose and HbA1c levels as risk factors for the development of hypertension in Japanese individuals: Toranomon hospital health management center study 16 (TOPICS 16). <i>Journal of Human Hypertension</i> , 2015 , 29, 254-9	2.6	19
250	Pioglitazone treatment and cardiovascular event and death in subjects with type 2 diabetes without established cardiovascular disease (JDDM 36). <i>Diabetes Research and Clinical Practice</i> , 2015 , 109, 485-92	7.4	12
249	A case of hypoglycemia attributable to atypical antipsychotic drugs. <i>Diabetology International</i> , 2015 , 6, 341-346	2.3	2
248	Risk of the development of Type 2 diabetes in relation to overall obesity, abdominal obesity and the clustering of metabolic abnormalities in Japanese individuals: does metabolically healthy overweight really exist? The Niigata Wellness Study. <i>Diabetic Medicine</i> , 2015 , 32, 665-72	3.5	20
247	Potential impact of joint association of alanine aminotransferase and gamma-glutamyltransferase on insulin resistance in Japan: The Toranomon Hospital Health Management Center Study 19 (TOPICS 19). <i>Hepatology Research</i> , 2015 , 45, 247-58	5.1	
246	Skeletal muscle-specific HMG-CoA reductase knockout mice exhibit rhabdomyolysis: A model for statin-induced myopathy. <i>Biochemical and Biophysical Research Communications</i> , 2015 , 466, 536-40	3.4	49
245	Prevalence of resistant hypertension and associated factors in Japanese subjects with type 2 diabetes. <i>Diabetes Research and Clinical Practice</i> , 2015 , 110, 18-25	7.4	7
244	The Combination of Elevated Triglycerides and Abnormal Fasting Glucose Increases Risk of Cerebral Infarction in Patients With Mild to Moderate Hypercholesterolemia: A Post Hoc Analysis of the MEGA Study. <i>Journal of Cardiovascular Pharmacology and Therapeutics</i> , 2015 , 20, 169-73	2.6	3
243	The factors that affect exercise therapy for patients with type 2 diabetes in Japan: a nationwide survey. <i>Diabetology International</i> , 2015 , 6, 19-25	2.3	11
242	Trajectory of body mass index before the development of type 2 diabetes in Japanese men: Toranomon Hospital Health Management Center Study 15. <i>Journal of Diabetes Investigation</i> , 2015 , 6, 289-94	3.9	15
241	Impact of population aging on trends in diabetes prevalence: A meta-regression analysis of 160,000 Japanese adults. <i>Journal of Diabetes Investigation</i> , 2015 , 6, 533-42	3.9	75
240	Late onset post-transfusion hepatitis E developing during chemotherapy for acute promyelocytic leukemia. <i>Internal Medicine</i> , 2015 , 54, 657-61	1.1	17
239	Type 1 diabetes mellitus and isolated adrenocorticotropin deficiency manifested by parkinsonism: a case report and literature review. <i>Internal Medicine</i> , 2015 , 54, 2629-35	1.1	6

238	Simple self-reported behavioral or psychological characteristics as risk factors for future type 2 diabetes in Japanese individuals: Toranomon Hospital Health Management Center Study 14. <i>Journal of Diabetes Investigation</i> , 2015 , 6, 236-41	3.9	2
237	Fatal tracheal aspergillosis during rituximab combined chemotherapy for diffuse large B-cell lymphoma that developed after lung transplantation. <i>Transplant Infectious Disease</i> , 2015 , 17, 872-5	2.7	0
236	Circulating malondialdehyde-modified LDL-related variables and coronary artery stenosis in asymptomatic patients with type 2 diabetes. <i>Journal of Diabetes Research</i> , 2015 , 2015, 507245	3.9	5
235	The association of level of reduction of WilmsTumor gene 1 mRNA transcript in bone marrow and outcome in acute myeloid leukemia patients. <i>Leukemia Research</i> , 2015 , 39, 667-71	2.7	5
234	Reply to: Relationship between leptin and blood pressure in patients with multiple system atrophy. <i>Journal of the Neurological Sciences</i> , 2015 , 348, 285	3.2	
233	Absence of Elovl6 attenuates steatohepatitis but promotes gallstone formation in a lithogenic diet-fed Ldlr(-/-) mouse model. <i>Scientific Reports</i> , 2015 , 5, 17604	4.9	15
232	Clinical Significance of MYC, BCL2 and BCL6 Rearrangement and Protein Expression in GCB and Non-GCB Type Diffuse Large B-Cell Lymphoma. <i>Blood</i> , 2015 , 126, 1445-1445	2.2	
231	The Sil Index Is a Useful Prognostic Indicator for Diffuse Large B-Cell Lymphoma. <i>Blood</i> , 2015 , 126, 1512-1512	2.2	
230	Comparison of different aspects of BMI history to identify undiagnosed diabetes in Japanese men and women: Toranomon Hospital Health Management Center Study 12 (TOPICS 12). <i>Diabetic Medicine</i> , 2014 , 31, 1378-86	3.5	6
229	Food groups and weight gain in Japanese men. <i>Clinical Obesity</i> , 2014 , 4, 157-64	3.6	0
228	Dietary sodium intake and incidence of diabetes complications in Japanese patients with type 2 diabetes: analysis of the Japan Diabetes Complications Study (JDACS). <i>Journal of Clinical Endocrinology and Metabolism</i> , 2014 , 99, 3635-43	5.6	55
227	Role of body mass index history in predicting risk of the development of hypertension in Japanese individuals: Toranomon Hospital Health Management Center Study 18 (TOPICS 18). <i>Hypertension</i> , 2014 , 64, 247-52	8.5	13
226	Fasting mediated increase in p-BAD(ser155) and p-AKT(ser473) in the prefrontal cortex of mice. <i>Neuroscience Letters</i> , 2014 , 579, 134-9	3.3	2
225	High risk of failing eradication of Helicobacter pylori in patients with diabetes: a meta-analysis. <i>Diabetes Research and Clinical Practice</i> , 2014 , 106, 81-7	7.4	28
224	Metabolically healthy obesity, presence or absence of fatty liver, and risk of type 2 diabetes in Japanese individuals: Toranomon Hospital Health Management Center Study 20 (TOPICS 20). <i>Journal of Clinical Endocrinology and Metabolism</i> , 2014 , 99, 2952-60	5.6	60
223	Ablation of Elovl6 protects pancreatic islets from high-fat diet-induced impairment of insulin secretion. <i>Biochemical and Biophysical Research Communications</i> , 2014 , 450, 318-23	3.4	13
222	Weight loss maintenance for 2 years after a 6-month randomised controlled trial comparing education-only and group-based support in Japanese adults. <i>Obesity Facts</i> , 2014 , 7, 376-87	5.1	13
221	Efficacy of habitual exercise for improving lipid profiles depends on the PPAR α genotype in Japanese males. <i>Journal of Nutritional Science and Vitaminology</i> , 2014 , 60, 66-70	1.1	

220	Early diagnosis of hepatic intravascular lymphoma: a case report and literature review. <i>Internal Medicine</i> , 2014 , 53, 587-93	1.1	9
219	Manifestations of fulminant CD8 T-cell post-transplant lymphoproliferative disorder following the administration of rituximab for lymphadenopathy with a high level of Epstein-Barr Virus (EBV) replication after allogeneic hematopoietic stem cell transplantation. <i>Internal Medicine</i> , 2014 , 53, 2115-9	1.1	2
218	Stability and changes in metabolically healthy overweight or obesity and risk of future diabetes: Niigata wellness study. <i>Obesity</i> , 2014 , 22, 2420-5	8	30
217	Combination effect of hypertension and diabetes mellitus on urinary protein excretion. <i>Journal of Hypertension</i> , 2014 , 32, 2278	1.9	
216	Hypertension increases urinary excretion of immunoglobulin G, ceruloplasmin and transferrin in normoalbuminuric patients with type 2 diabetes mellitus. <i>Journal of Hypertension</i> , 2014 , 32, 432-8	1.9	15
215	A possible link between BDNF and mTOR in control of food intake. <i>Frontiers in Psychology</i> , 2014 , 5, 1093	3.4	34
214	Reversible brain atrophy and cognitive impairment in an adolescent Japanese patient with primary adrenal Cushing β syndrome. <i>Neuropsychiatric Disease and Treatment</i> , 2014 , 10, 1763-7	3.1	2
213	Comparison of the Framingham risk score, UK Prospective Diabetes Study (UKPDS) Risk Engine, Japanese Atherosclerosis Longitudinal Study-Existing Cohorts Combine (JALS-ECC) and maximum carotid intima-media thickness for predicting coronary artery stenosis in patients with	4	23
212	Utility of the triglyceride level for predicting incident diabetes mellitus according to the fasting status and body mass index category: the Ibaraki Prefectural Health Study. <i>Journal of Atherosclerosis and Thrombosis</i> , 2014 , 21, 1152-69	4	13
211	Cohort profile: The Japan diabetes complications study: a long-term follow-up of a randomised lifestyle intervention study of type 2 diabetes. <i>International Journal of Epidemiology</i> , 2014 , 43, 1054-62	7.8	34
210	Impact on short-term glycaemic control of initiating diabetes care versus leaving diabetes untreated among individuals with newly screening-detected diabetes in Japan. <i>Journal of Epidemiology and Community Health</i> , 2014 , 68, 1189-95	5.1	5
209	Response to comment on Heianza et al. Effect of postmenopausal status and age at menopause on type 2 diabetes and prediabetes in Japanese individuals: Toranomon Hospital Health Management Center Study 17 (TOPICS 17). <i>Diabetes Care</i> 2013;36:4007-4014. <i>Diabetes Care</i> , 2014 , 37, e165-6	14.6	
208	Body mass index and mortality among Japanese patients with type 2 diabetes: pooled analysis of the Japan diabetes complications study and the Japanese elderly diabetes intervention trial. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2014 , 99, E2692-6	5.6	25
207	Dietary intake in Japanese patients with type 2 diabetes: Analysis from Japan Diabetes Complications Study. <i>Journal of Diabetes Investigation</i> , 2014 , 5, 176-87	3.9	28
206	Commentary on the United Kingdom Prospective Diabetes Study outcomes model 2: Need for long-term follow up and quality of life data in Asian patients. <i>Journal of Diabetes Investigation</i> , 2014 , 5, 281-3	3.9	1
205	Importance of high-density lipoprotein cholesterol levels in elderly diabetic individuals with type IIb dyslipidemia: A 2-year survey of cardiovascular events. <i>Geriatrics and Gerontology International</i> , 2014 , 14, 806-10	2.9	2
204	Efficacy of HMG-CoA reductase inhibitors in the prevention of cerebrovascular attack in 1016 patients older than 75 years among 4014 type 2 diabetic individuals. <i>International Journal of Cardiology</i> , 2014 , 177, 860-6	3.2	1
203	The circulating level of leptin and blood pressure in patients with multiple system atrophy. <i>Journal of the Neurological Sciences</i> , 2014 , 347, 349-51	3.2	4

202	Role of sleep duration as a risk factor for Type 2 diabetes among adults of different ages in Japan: the Niigata Wellness Study. <i>Diabetic Medicine</i> , 2014 , 31, 1363-7	3.5	21
201	Hepatic CREB3L3 controls whole-body energy homeostasis and improves obesity and diabetes. <i>Endocrinology</i> , 2014 , 155, 4706-19	4.8	34
200	Quantitative relationship between body weight gain in adulthood and incident type 2 diabetes: a meta-analysis. <i>Obesity Reviews</i> , 2014 , 15, 202-14	10.6	69
199	Association of Helicobacter pylori infection with glycemic control in patients with diabetes: a meta-analysis. <i>Journal of Diabetes Research</i> , 2014 , 2014, 250620	3.9	15
198	High cholesterol intake is associated with elevated risk of type 2 diabetes mellitus - a meta-analysis. <i>Clinical Nutrition</i> , 2014 , 33, 946-50	5.9	18
197	Second trimester postload glucose level as an important predictor of low birth weight infants: Tanaka Women's Clinic Study. <i>Diabetes Research and Clinical Practice</i> , 2014 , 105, e16-9	7.4	1
196	Meta-analysis of the quantitative relation between pulse pressure and mean arterial pressure and cardiovascular risk in patients with diabetes mellitus. <i>American Journal of Cardiology</i> , 2014 , 113, 1058-65 ³		27
195	Log Reduction Levels of WT1 mRNA Expression in BM after Chemotherapies Are Predictive Markers of Good Prognosis in AML Patients Achieved CR after Induction Therapy. <i>Blood</i> , 2014 , 124, 2334-2334	2.2	
194	Metabolic predictors of ischemic heart disease and cerebrovascular attack in elderly diabetic individuals: difference in risk by age. <i>Cardiovascular Diabetology</i> , 2013 , 12, 10	8.7	7
193	Association between physical activity and risk of all-cause mortality and cardiovascular disease in patients with diabetes: a meta-analysis. <i>Diabetes Care</i> , 2013 , 36, 471-9	14.6	114
192	Predicting macro- and microvascular complications in type 2 diabetes: the Japan Diabetes Complications Study/the Japanese Elderly Diabetes Intervention Trial risk engine. <i>Diabetes Care</i> , 2013 , 36, 1193-9	14.6	70
191	Use of high-normal levels of haemoglobin A(1C) and fasting plasma glucose for diabetes screening and for prediction: a meta-analysis. <i>Diabetes/Metabolism Research and Reviews</i> , 2013 , 29, 680-92	7.5	17
190	Importance of high-density lipoprotein cholesterol control during pravastatin treatment in hypercholesterolemic Japanese with type 2 diabetes mellitus: a post hoc analysis of MEGA study. <i>Diabetes Research and Clinical Practice</i> , 2013 , 100, e31-3	7.4	3
189	Diabetic retinopathy and microalbuminuria can predict macroalbuminuria and renal function decline in Japanese type 2 diabetic patients: Japan Diabetes Complications Study. <i>Diabetes Care</i> , 2013 , 36, 2803-9	14.6	35
188	The hyperglycemia stimulated myocardial endoplasmic reticulum (ER) stress contributes to diabetic cardiomyopathy in the transgenic non-obese type 2 diabetic rats: a differential role of unfolded protein response (UPR) signaling proteins. <i>International Journal of Biochemistry and Cell Biology</i> , 2013 , 45, 438-47	5.6	70
187	TFE3 inhibits myoblast differentiation in C2C12 cells via down-regulating gene expression of myogenin. <i>Biochemical and Biophysical Research Communications</i> , 2013 , 430, 664-9	3.4	9
186	Risk of cardiovascular diseases is increased even with mild diabetic retinopathy: the Japan Diabetes Complications Study. <i>Ophthalmology</i> , 2013 , 120, 574-582	7.3	56
185	Low BMI at age 20 years predicts gestational diabetes independent of BMI in early pregnancy in Japan: Tanaka Women's Clinic Study. <i>Diabetic Medicine</i> , 2013 , 30, 70-3	3.5	15

184	Correction to: Comparison of Education-Only versus Group-Based Intervention in Promoting Weight Loss: A Randomised Controlled Trial. <i>Obesity Facts</i> , 2013 , 6, 89-90	5.1	78
183	Leisure-time physical activity is a significant predictor of stroke and total mortality in Japanese patients with type 2 diabetes: analysis from the Japan Diabetes Complications Study (JDACS). <i>Diabetologia</i> , 2013 , 56, 1021-30	10.3	36
182	Molecular understanding of curcumin in diabetic nephropathy. <i>Drug Discovery Today</i> , 2013 , 18, 756-63	8.8	35
181	TFE3 controls lipid metabolism in adipose tissue of male mice by suppressing lipolysis and thermogenesis. <i>Endocrinology</i> , 2013 , 154, 3577-88	4.8	26
180	Development of a screening score for undiagnosed diabetes and its application in estimating absolute risk of future type 2 diabetes in Japan: Toranomon Hospital Health Management Center Study 10 (TOPICS 10). <i>Journal of Clinical Endocrinology and Metabolism</i> , 2013 , 98, 1051-60	5.6	19
179	Role of alcohol drinking pattern in type 2 diabetes in Japanese men: the Toranomon Hospital Health Management Center Study 11 (TOPICS 11). <i>American Journal of Clinical Nutrition</i> , 2013 , 97, 561-87	7	27
178	Quality and accuracy of Internet information concerning a healthy diet. <i>International Journal of Food Sciences and Nutrition</i> , 2013 , 64, 1007-13	3.7	12
177	Fruit intake and incident diabetic retinopathy with type 2 diabetes. <i>Epidemiology</i> , 2013 , 24, 204-11	3.1	50
176	Intakes of dietary fiber, vegetables, and fruits and incidence of cardiovascular disease in Japanese patients with type 2 diabetes. <i>Diabetes Care</i> , 2013 , 36, 3916-22	14.6	28
175	Efficacy and safety in sitagliptin therapy for diabetes complicated by non-alcoholic fatty liver disease. <i>Hepatology Research</i> , 2013 , 43, 1163-8	5.1	24
174	Two authors reply. <i>American Journal of Epidemiology</i> , 2013 , 177, 863	3.8	
173	Direct effect of dasatinib on proliferation and cytotoxicity of natural killer cells in in vitro study. <i>Hematological Oncology</i> , 2013 , 31, 156-63	1.3	14
172	Association of living alone with the presence of undiagnosed diabetes in Japanese men: the role of modifiable risk factors for diabetes: Toranomon Hospital Health Management Center Study 13 (TOPICS 13). <i>Diabetic Medicine</i> , 2013 , 30, 1355-9	3.5	3
171	Diabetes and risk of hearing impairment in adults: a meta-analysis. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2013 , 98, 51-8	5.6	141
170	Effect of postmenopausal status and age at menopause on type 2 diabetes and prediabetes in Japanese individuals: Toranomon Hospital Health Management Center Study 17 (TOPICS 17). <i>Diabetes Care</i> , 2013 , 36, 4007-14	14.6	64
169	Association between remission of macroalbuminuria and preservation of renal function in patients with type 2 diabetes with overt proteinuria. <i>Diabetes Care</i> , 2013 , 36, 3227-33	14.6	36
168	Genetic variants of the fatty acid desaturase gene cluster are associated with plasma LDL cholesterol levels in Japanese males. <i>Journal of Nutritional Science and Vitaminology</i> , 2013 , 59, 325-35	1.1	15
167	Body weight change and type 2 diabetes. <i>Epidemiology</i> , 2013 , 24, 778-9	3.1	7

166	Carotid artery plaque and LDL-to-HDL cholesterol ratio predict atherosclerotic status in coronary arteries in asymptomatic patients with type 2 diabetes mellitus. <i>Journal of Atherosclerosis and Thrombosis</i> , 2013 , 20, 452-64	4	29
165	Early Responses At 3 Months and 12 Months After Starting Imatinib As Predictive Factors For The Achievement Of Deep MR In Japanese CML Patients. <i>Blood</i> , 2013 , 122, 2744-2744	2.2	
164	Nutrition education in Japanese medical schools: a follow-up survey. <i>Asia Pacific Journal of Clinical Nutrition</i> , 2013 , 22, 144-9	1	11
163	Long-term multiple risk factor interventions in Japanese elderly diabetic patients: the Japanese Elderly Diabetes Intervention Trial--study design, baseline characteristics and effects of intervention. <i>Geriatrics and Gerontology International</i> , 2012 , 12 Suppl 1, 7-17	2.9	57
162	Non-high-density lipoprotein cholesterol: an important predictor of stroke and diabetes-related mortality in Japanese elderly diabetic patients. <i>Geriatrics and Gerontology International</i> , 2012 , 12 Suppl 1, 18-28	2.9	28
161	Prevalence and risk factors for diabetic maculopathy, and its relationship to diabetic retinopathy in elderly Japanese patients with type 2 diabetes mellitus. <i>Geriatrics and Gerontology International</i> , 2012 , 12 Suppl 1, 134-40	2.9	8
160	Long-term risk factors for diabetic retinopathy and diabetic maculopathy in elderly Japanese patients with type 2 diabetes mellitus. <i>Geriatrics and Gerontology International</i> , 2012 , 12 Suppl 1, 141-4	2.9	1
159	High normal HbA(1c) levels were associated with impaired insulin secretion without escalating insulin resistance in Japanese individuals: the Toranomon Hospital Health Management Center Study 8 (TOPICS 8). <i>Diabetic Medicine</i> , 2012 , 29, 1285-90	3.5	22
158	Screening for pre-diabetes to predict future diabetes using various cut-off points for HbA(1c) and impaired fasting glucose: the Toranomon Hospital Health Management Center Study 4 (TOPICS 4). <i>Diabetic Medicine</i> , 2012 , 29, e279-85	3.5	49
157	Elovl6 promotes nonalcoholic steatohepatitis. <i>Hepatology</i> , 2012 , 56, 2199-208	11.2	120
156	Self-reported rate of eating is significantly associated with body mass index in Japanese patients with type 2 diabetes. Japan Diabetes Clinical Data Management Study Group (JDDM26). <i>Appetite</i> , 2012 , 59, 252-5	4.5	15
155	Impact of introducing HbA1c into the diagnostic criteria on prevalence and cardiovascular risk profiles of individuals with newly diagnosed diabetes in Japan: the Toranomon Hospital Health Management Center Study 2 (TOPICS 2). <i>Diabetes Research and Clinical Practice</i> , 2012 , 95, 283-90	7.4	9
154	Longitudinal trajectories of HbA1c and fasting plasma glucose levels during the development of type 2 diabetes: the Toranomon Hospital Health Management Center Study 7 (TOPICS 7). <i>Diabetes Care</i> , 2012 , 35, 1050-2	14.6	32
153	Cholesterol sulfate induces expression of the skin barrier protein filaggrin in normal human epidermal keratinocytes through induction of ROR β . <i>Biochemical and Biophysical Research Communications</i> , 2012 , 428, 99-104	3.4	23
152	Enhancement of antigen presenting ability in the leukemic plasmacytoid dendritic cell line (PMDC05) by lentiviral vector-mediated transduction of CD80 gene. <i>Leukemia Research</i> , 2012 , 36, 1541-6	2.7	3
151	Low lung function and risk of type 2 diabetes in Japanese men: the Toranomon Hospital Health Management Center Study 9 (TOPICS 9). <i>Mayo Clinic Proceedings</i> , 2012 , 87, 853-61	6.4	25
150	Development of a new scoring system for predicting the 5-year incidence of type 2 diabetes in Japan: the Toranomon Hospital Health Management Center Study 6 (TOPICS 6). <i>Diabetologia</i> , 2012 , 55, 3213-23	10.3	33
149	Fasting insulin levels and metabolic risk factors in type 2 diabetic patients at the first visit in Japan: a 10-year, nationwide, observational study (JDDM 28). <i>Diabetes Care</i> , 2012 , 35, 1853-7	14.6	18

148	Management of type IIb dyslipidemia. <i>Journal of Atherosclerosis and Thrombosis</i> , 2012 , 19, 105-114	4	12
147	Quantification of BCR-ABL mRNA in plasma/serum of patients with chronic myelogenous leukemia. <i>International Journal of Medical Sciences</i> , 2012 , 9, 901-8	3.7	7
146	Diagnosis and management of type I and type V hyperlipoproteinemia. <i>Journal of Atherosclerosis and Thrombosis</i> , 2012 , 19, 1-12	4	65
145	Fasting and post-challenge glucose as quantitative cardiovascular risk factors: a meta-analysis. <i>Journal of Atherosclerosis and Thrombosis</i> , 2012 , 19, 385-96	4	17
144	Present situation of exercise therapy for patients with diabetes mellitus in Japan: a nationwide survey. <i>Diabetology International</i> , 2012 , 3, 86-91	2.3	5
143	Chronic kidney disease categories and renal-cardiovascular outcomes in type 2 diabetes without prevalent cardiovascular disease: a prospective cohort study (JDDM25). <i>Diabetologia</i> , 2012 , 55, 1911-8	10.3	33
142	HbA(1c) variability and the development of microalbuminuria in type 2 diabetes: Tsukuba Kawai Diabetes Registry 2. <i>Diabetologia</i> , 2012 , 55, 2128-31	10.3	75
141	Impact of psychological stress caused by the Great East Japan Earthquake on glycemic control in patients with diabetes. <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2012 , 120, 560-3	2.3	32
140	TFE3 regulates muscle metabolic gene expression, increases glycogen stores, and enhances insulin sensitivity in mice. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2012 , 302, E896-902	6	26
139	Comparison of various lipid variables as predictors of coronary heart disease in Japanese men and women with type 2 diabetes: subanalysis of the Japan Diabetes Complications Study. <i>Diabetes Care</i> , 2012 , 35, 1150-7	14.6	38
138	Comparisons of the strength of associations with future type 2 diabetes risk among anthropometric obesity indicators, including waist-to-height ratio: a meta-analysis. <i>American Journal of Epidemiology</i> , 2012 , 176, 959-69	3.8	150
137	Effect of Web-based lifestyle modification on weight control: a meta-analysis. <i>International Journal of Obesity</i> , 2012 , 36, 675-85	5.5	98
136	Quality of Internet information related to the Mediterranean diet. <i>Public Health Nutrition</i> , 2012 , 15, 885-93	3.3	25
135	Background to discuss guidelines for control of plasma HDL-cholesterol in Japan. <i>Journal of Atherosclerosis and Thrombosis</i> , 2012 , 19, 207-12	4	3
134	Alcohol consumption and risk of atrial fibrillation: a meta-analysis. <i>Journal of the American College of Cardiology</i> , 2011 , 57, 427-36	15.1	190
133	Contribution of glimepiride to basal-prandial insulin therapy in patients with type 2 diabetes. <i>Diabetes Research and Clinical Practice</i> , 2011 , 91, 148-53	7.4	7
132	Contribution of first trimester fasting plasma insulin levels to the incidence of glucose intolerance in later pregnancy: Tanaka women's clinic study. <i>Diabetes Research and Clinical Practice</i> , 2011 , 92, 293-8	7.4	10
131	Self-reported fast eating is a potent predictor of development of impaired glucose tolerance in Japanese men and women: Tsukuba Medical Center Study. <i>Diabetes Research and Clinical Practice</i> , 2011 , 94, e72-4	7.4	7

130	Accumulation of cardiovascular risks in Japanese women with abnormal glucose and mild to moderate hypercholesterolemia. <i>International Journal of Cardiology</i> , 2011 , 152, 254-6	3.2	
129	Comparison of education-only versus group-based intervention in promoting weight loss: a randomised controlled trial. <i>Obesity Facts</i> , 2011 , 4, 222-8	5.1	13
128	HbA1c 5.7-6.4% and impaired fasting plasma glucose for diagnosis of prediabetes and risk of progression to diabetes in Japan (TOPICS 3): a longitudinal cohort study. <i>Lancet, The</i> , 2011 , 378, 147-55	4.0	167
127	Flow-mediated dilation is associated with microalbuminuria independent of cardiovascular risk factors in type 2 diabetes - interrelations with arterial thickness and stiffness. <i>Journal of Atherosclerosis and Thrombosis</i> , 2011 , 18, 744-52	4	24
126	Low incidence of cardiovascular events in Japanese patients with Type 2 diabetes in primary care settings: a prospective cohort study (JDDM 20). <i>Diabetic Medicine</i> , 2011 , 28, 1221-8	3.5	32
125	Skipping breakfast and prevalence of overweight and obesity in Asian and Pacific regions: a meta-analysis. <i>Preventive Medicine</i> , 2011 , 53, 260-7	4.3	155
124	Low transition rate from normo- and low microalbuminuria to proteinuria in Japanese type 2 diabetic individuals: the Japan Diabetes Complications Study (JDCCS). <i>Diabetologia</i> , 2011 , 54, 1025-31	10.3	48
123	Low serum potassium levels and risk of type 2 diabetes: the Toranomon Hospital Health Management Center Study 1 (TOPICS 1). <i>Diabetologia</i> , 2011 , 54, 762-6	10.3	30
122	Incidence and progression of diabetic retinopathy in Japanese adults with type 2 diabetes: 8 year follow-up study of the Japan Diabetes Complications Study (JDCCS). <i>Diabetologia</i> , 2011 , 54, 2288-94	10.3	72
121	Relationship between hemoglobin A1c and cardiovascular disease in mild-to-moderate hypercholesterolemic Japanese individuals: subanalysis of a large-scale randomized controlled trial. <i>Cardiovascular Diabetology</i> , 2011 , 10, 58	8.7	14
120	Age, gender, insulin and blood glucose control status alter the risk of ischemic heart disease and stroke among elderly diabetic patients. <i>Cardiovascular Diabetology</i> , 2011 , 10, 86	8.7	10
119	Dose of 3-methylcholanthrene enhances vitamin C accumulation and mRNA expression of its transporter in the liver of ODS rats and in HepG2 cells. <i>Journal of Biochemical and Molecular Toxicology</i> , 2011 , 25, 369-76	3.4	
118	The level of orally ingested vitamin C affected the expression of vitamin C transporters and vitamin C accumulation in the livers of ODS rats. <i>Bioscience, Biotechnology and Biochemistry</i> , 2011 , 75, 2394-7	2.1	1
117	Factors associated with inappropriate weight loss attempts by early adolescent girls in Japan. <i>Eating and Weight Disorders</i> , 2011 , 16, e157-63	3.6	2
116	Risks for glomerular filtration rate decline in association with progression of albuminuria in type 2 diabetes. <i>Nephrology Dialysis Transplantation</i> , 2011 , 26, 2924-30	4.3	30
115	Serum level of triglycerides is a potent risk factor comparable to LDL cholesterol for coronary heart disease in Japanese patients with type 2 diabetes: subanalysis of the Japan Diabetes Complications Study (JDCCS). <i>Journal of Clinical Endocrinology and Metabolism</i> , 2011 , 96, 3448-56	5.6	135
114	Macrophage Elovl6 deficiency ameliorates foam cell formation and reduces atherosclerosis in low-density lipoprotein receptor-deficient mice. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2011 , 31, 1973-9	9.4	20
113	Sterol regulatory element-binding protein-1 determines plasma remnant lipoproteins and accelerates atherosclerosis in low-density lipoprotein receptor-deficient mice. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2011 , 31, 1788-95	9.4	33

112	Inhibition of ubiquitin ligase F-box and WD repeat domain-containing 7 (Fbw7) causes hepatosteatosis through Krüppel-like factor 5 (KLF5)/peroxisome proliferator-activated receptor α (PPAR α) pathway but not SREBP-1c protein in mice. <i>Journal of Biological Chemistry</i> , 2011 , 286, 40835-46	5.4	22
111	Evidence for the Diagnosis of Metabolic Syndrome in Japan. <i>The Japanese Journal of Nutrition and Dietetics</i> , 2011 , 69, 205-213	0.2	2
110	A Hierarchical Regression Model for Dietary Data Adjusting for Covariates Measurement Error by Regression Calibration: An Application to a Large Prospective Study for Diabetic Complications. <i>Japanese Journal of Biometrics</i> , 2010 , 31, 49-62	0	
109	Effect of endurance exercise training on oxidative stress in spontaneously hypertensive rats (SHR) after emergence of hypertension. <i>Clinical and Experimental Hypertension</i> , 2010 , 32, 407-15	2.2	27
108	Are serum cholesterol levels associated with silent brain infarcts? The Seiryu Clinic Study. <i>Atherosclerosis</i> , 2010 , 210, 674-7	3.1	9
107	Oxygenomics in environmental stress. <i>Redox Report</i> , 2010 , 15, 98-114	5.9	18
106	Effects of magnesium on postprandial serum lipid responses in healthy human subjects. <i>British Journal of Nutrition</i> , 2010 , 103, 469-72	3.6	36
105	Body Image: International Comparisons. <i>Nutrition Today</i> , 2010 , 45, 113-117	1.6	1
104	Long-term lifestyle intervention lowers the incidence of stroke in Japanese patients with type 2 diabetes: a nationwide multicentre randomised controlled trial (the Japan Diabetes Complications Study). <i>Diabetologia</i> , 2010 , 53, 419-28	10.3	101
103	Astaxanthin suppresses scavenger receptor expression and matrix metalloproteinase activity in macrophages. <i>European Journal of Nutrition</i> , 2010 , 49, 119-26	5.2	69
102	Influence of fat and carbohydrate proportions on the metabolic profile in patients with type 2 diabetes: a meta-analysis. <i>Diabetes Care</i> , 2009 , 32, 959-65	14.6	119
101	Low HDL cholesterol is associated with the risk of stroke in elderly diabetic individuals: changes in the risk for atherosclerotic diseases at various ages. <i>Diabetes Care</i> , 2009 , 32, 1221-3	14.6	23
100	Cardiorespiratory fitness as a quantitative predictor of all-cause mortality and cardiovascular events in healthy men and women: a meta-analysis. <i>JAMA - Journal of the American Medical Association</i> , 2009 , 301, 2024-35	27.4	1725
99	Determinants of decline in glomerular filtration rate in nonproteinuric subjects with or without diabetes and hypertension. <i>Clinical Journal of the American Society of Nephrology: CJASN</i> , 2009 , 4, 1432-40	6.9	59
98	Physical fitness during adolescence and adult mortality. <i>Epidemiology</i> , 2009 , 20, 463-4	3.1	23
97	Thinness in Japanese young women. <i>Epidemiology</i> , 2009 , 20, 464-5	3.1	21
96	Waist circumference as a cardiovascular and metabolic risk in Japanese patients with type 2 diabetes. <i>Obesity</i> , 2009 , 17, 585-92	8	15
95	Association between serum uric acid and development of type 2 diabetes. <i>Diabetes Care</i> , 2009 , 32, 1737-42	14.6	325

94	No relationship between body mass index during adolescence and all-cause mortality in Japanese women--a 56.5-year observational study. <i>Annals of Epidemiology</i> , 2009 , 19, 590-1	6.4	
93	Prevalence of albuminuria and renal insufficiency and associated clinical factors in type 2 diabetes: the Japan Diabetes Clinical Data Management study (JDDM15). <i>Nephrology Dialysis Transplantation</i> , 2009 , 24, 1212-9	4.3	134
92	Components of metabolic syndrome and their combinations as predictors of cardiovascular disease in Japanese patients with type 2 diabetes. Implications for improved definition. Analysis from Japan Diabetes Complications Study (JDCCS). <i>Journal of Atherosclerosis and Thrombosis</i> , 2009 , 16, 380-7	4	25
91	Reduced GFR and microalbuminuria are independently associated with prevalent cardiovascular disease in Type 2 diabetes: JDDM study 16. <i>Diabetic Medicine</i> , 2008 , 25, 1426-32	3.5	26
90	Familial predisposition to cardiovascular risk and disease contributes to cardiovascular risk and disease interacting with other cardiovascular risk factors in diabetes: implication for common soil (JDDM 14). <i>Atherosclerosis</i> , 2008 , 201, 332-8	3.1	10
89	Palmitate impairs and eicosapentaenoate restores insulin secretion through regulation of SREBP-1c in pancreatic islets. <i>Diabetes</i> , 2008 , 57, 2382-92	0.9	72
88	Importance of lipid levels in elderly diabetic individuals: baseline characteristics and 1-year survey of cardiovascular events. <i>Circulation Journal</i> , 2008 , 72, 218-25	2.9	8
87	Effect of weight reduction on concentration of plasma total homocysteine in obese Japanese men. <i>Obesity Research and Clinical Practice</i> , 2007 , 1, I-II	5.4	3
86	Crucial role of a long-chain fatty acid elongase, Elovl6, in obesity-induced insulin resistance. <i>Nature Medicine</i> , 2007 , 13, 1193-202	50.5	380
85	Risk imparted by various parameters of smoking in Japanese men with type 2 diabetes on their development of microalbuminuria: analysis from the Tsukuba Kawai Diabetes Registry. <i>Diabetes Care</i> , 2007 , 30, 1286-8	14.6	9
84	The regulation of adiponectin receptors expression by acute exercise in mice. <i>Experimental and Clinical Endocrinology and Diabetes</i> , 2007 , 115, 417-22	2.3	18
83	Effect of aerobic exercise training on serum levels of high-density lipoprotein cholesterol: a meta-analysis. <i>Archives of Internal Medicine</i> , 2007 , 167, 999-1008		379
82	Protein kinase A suppresses sterol regulatory element-binding protein-1C expression via phosphorylation of liver X receptor in the liver. <i>Journal of Biological Chemistry</i> , 2007 , 282, 11687-95	5.4	79
81	Even low-intensity and low-volume exercise training may improve insulin resistance in the elderly. <i>Internal Medicine</i> , 2007 , 46, 1071-7	1.1	31
80	Influence of adiponectin gene polymorphism SNP276 (G/T) on adiponectin in response to exercise training. <i>Endocrine Journal</i> , 2007 , 54, 879-86	2.9	14
79	Involvement of glomerular SREBP-1c in diabetic nephropathy. <i>Biochemical and Biophysical Research Communications</i> , 2007 , 364, 502-8	3.4	47
78	Cross-sectional association between BMI, glycemic control and energy intake in Japanese patients with type 2 diabetes. Analysis from the Japan Diabetes Complications Study. <i>Diabetes Research and Clinical Practice</i> , 2007 , 77 Suppl 1, S23-9	7.4	18
77	EFFECT OF EXERCISE TRAINING ON SERUM HIGH-SENSITIVITY C-REACTIVE PROTEIN CONCENTRATION IN HEALTHY MIDDLE-AGED AND ELDERLY SUBJECTS. <i>Japanese Journal of Physical Fitness and Sports Medicine</i> , 2007 , 56, 179-190	0.1	

76	Antioxidants and an inhibitor of advanced glycation ameliorate death of retinal microvascular cells in diabetic retinopathy. <i>Diabetes/Metabolism Research and Reviews</i> , 2006 , 22, 38-45	7.5	40
75	The effect of exercise training on adiponectin receptor expression in KKAy obese/diabetic mice. <i>Journal of Endocrinology</i> , 2006 , 189, 643-53	4.7	44
74	Granuphilin is activated by SREBP-1c and involved in impaired insulin secretion in diabetic mice. <i>Cell Metabolism</i> , 2006 , 4, 143-54	24.6	55
73	Outcome of one-year of specialist care of patients with type 2 diabetes: a multi-center prospective survey (JDDM 2). <i>Internal Medicine</i> , 2006 , 45, 589-97	1.1	20
72	TFE3 transcriptionally activates hepatic IRS-2, participates in insulin signaling and ameliorates diabetes. <i>Nature Medicine</i> , 2006 , 12, 107-13	50.5	124
71	The new worldwide definition of metabolic syndrome is not a better diagnostic predictor of cardiovascular disease in Japanese diabetic patients than the existing definitions: additional analysis from the Japan Diabetes Complications Study. <i>Diabetes Care</i> , 2006 , 29, 145-7	14.6	11
70	Transgenic mice overexpressing SREBP-1a under the control of the PEPCK promoter exhibit insulin resistance, but not diabetes. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2005 , 1740, 427-33	6.9	18
69	Plasma chloride concentration as a new diagnostic indicator of insulin insufficiency. <i>Diabetes Research and Clinical Practice</i> , 2005 , 67, 137-43	7.4	0
68	Lipid synthetic transcription factor SREBP-1a activates p21WAF1/CIP1, a universal cyclin-dependent kinase inhibitor. <i>Molecular and Cellular Biology</i> , 2005 , 25, 8938-47	4.8	50
67	Is the diagnosis of metabolic syndrome useful for predicting cardiovascular disease in asian diabetic patients? Analysis from the Japan Diabetes Complications Study. <i>Diabetes Care</i> , 2005 , 28, 1463-71	14.6	113
66	Sterol regulatory element-binding proteins activate insulin gene promoter directly and indirectly through synergy with BETA2/E47. <i>Journal of Biological Chemistry</i> , 2005 , 280, 34577-89	5.4	23
65	Transgenic mice overexpressing nuclear SREBP-1c in pancreatic beta-cells. <i>Diabetes</i> , 2005 , 54, 492-9	0.9	73
64	Vascular risk factors and diabetic neuropathy. <i>New England Journal of Medicine</i> , 2005 , 352, 1925-7; author reply 1925-7	59.2	10
63	Insulin-independent induction of sterol regulatory element-binding protein-1c expression in the livers of streptozotocin-treated mice. <i>Diabetes</i> , 2004 , 53, 560-9	0.9	151
62	Type 2 diabetes prevalence in Asian subjects. <i>Diabetes Care</i> , 2004 , 27, 1251-2; author reply 1252	14.6	35
61	SREBP-1 interacts with hepatocyte nuclear factor-4 alpha and interferes with PGC-1 recruitment to suppress hepatic gluconeogenic genes. <i>Journal of Biological Chemistry</i> , 2004 , 279, 12027-35	5.4	120
60	SREBPs suppress IRS-2-mediated insulin signalling in the liver. <i>Nature Cell Biology</i> , 2004 , 6, 351-7	23.4	260
59	Statins downregulate ATP-binding-cassette transporter A1 gene expression in macrophages. <i>Biochemical and Biophysical Research Communications</i> , 2004 , 316, 790-4	3.4	51

58	WGEF is a novel RhoGEF expressed in intestine, liver, heart, and kidney. <i>Biochemical and Biophysical Research Communications</i> , 2004 , 324, 1053-8	3.4	18
57	Requirement for hypertension and hyperlipidemia medication in U.S. and Japanese patients with diabetes. <i>American Journal of Medicine</i> , 2004 , 117, 711-2	2.4	3
56	Energy intake and obesity in Japanese patients with type 2 diabetes. <i>Lancet, The</i> , 2004 , 363, 248-9	4.0	47
55	Alcohol use and diabetes mellitus. <i>Annals of Internal Medicine</i> , 2004 , 141, 408-9; author reply 409	8	6
54	p57Kip2 regulates actin dynamics by binding and translocating LIM-kinase 1 to the nucleus. <i>Journal of Biological Chemistry</i> , 2003 , 278, 52919-23	5.4	88
53	Cross-talk between peroxisome proliferator-activated receptor (PPAR) alpha and liver X receptor (LXR) in nutritional regulation of fatty acid metabolism. I. PPARs suppress sterol regulatory element binding protein-1c promoter through inhibition of LXR signaling. <i>Molecular Endocrinology</i> , 2003 , 17, 4210-21		214
52	Effect of thiazolidinediones and metformin on LDL oxidation and aortic endothelium relaxation in diabetic GK rats. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2003 , 284, E1125-30	6	50
51	Eicosapentaenoic acid suppresses basal and insulin-stimulated endothelin-1 production in human endothelial cells. <i>Hypertension Research</i> , 2003 , 26, 655-61	4.7	22
50	Mouse MafA, homologue of zebrafish somite Maf 1, contributes to the specific transcriptional activity through the insulin promoter. <i>Biochemical and Biophysical Research Communications</i> , 2003 , 312, 831-42	3.4	57
49	Obesity and type 2 diabetes in Japanese patients. <i>Lancet, The</i> , 2003 , 361, 85	4.0	145
48	Physiological changes in circulating mannose levels in normal, glucose-intolerant, and diabetic subjects. <i>Metabolism: Clinical and Experimental</i> , 2003 , 52, 1019-27	12.7	35
47	Cross-talk between peroxisome proliferator-activated receptor (PPAR) alpha and liver X receptor (LXR) in nutritional regulation of fatty acid metabolism. II. LXRs suppress lipid degradation gene promoters through inhibition of PPAR signaling. <i>Molecular Endocrinology</i> , 2003 , 17, 1255-67		140
46	Low-density lipoprotein receptor-related protein 5 (LRP5) is essential for normal cholesterol metabolism and glucose-induced insulin secretion. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2003 , 100, 229-34	11.5	338
45	Genetic and physiological analysis of branched-chain alcohols and isoamyl acetate production in <i>Saccharomyces cerevisiae</i> . <i>Applied Microbiology and Biotechnology</i> , 2002 , 59, 501-8	5.7	64
44	The long-term effects of self-management education for patients with type 2 diabetes on glycemic control: response to Norris et al. <i>Diabetes Care</i> , 2002 , 25, 2115-6	14.6	10
43	Effects of lifestyle modifications on patients with type 2 diabetes: the Japan Diabetes Complications Study (JDACS) study design, baseline analysis and three year-interim report. <i>Hormone and Metabolic Research</i> , 2002 , 34, 509-15	3.1	85
42	Acetyl-coenzyme A synthetase is a lipogenic enzyme controlled by SREBP-1 and energy status. <i>American Journal of Physiology - Endocrinology and Metabolism</i> , 2002 , 282, E222-30	6	67
41	Insulin inhibits apoptosis of macrophage cell line, THP-1 cells, via phosphatidylinositol-3-kinase-dependent pathway. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2002 , 22, 380-6	9.4	48

40	Polyunsaturated fatty acids suppress sterol regulatory element-binding protein 1c promoter activity by inhibition of liver X receptor (LXR) binding to LXR response elements. <i>Journal of Biological Chemistry</i> , 2002 , 277, 1705-11	5.4	299
39	Glucose uptake and adenoviral mediated GLUT1 infection decrease hypoxia-induced HIF-1alpha levels in cardiac myocytes. <i>Journal of Molecular and Cellular Cardiology</i> , 2002 , 34, 1063-73	5.8	30
38	Acute gliclazide administration enhances glucose and ketone body utilization in the perfused hind limb of normal and streptozotocin-diabetic rats. <i>Life Sciences</i> , 2002 , 71, 647-54	6.8	4
37	Vascular endothelial growth factor gene expression in a retinal pigmented cell is up-regulated by glucose deprivation through 3PUTR. <i>Life Sciences</i> , 2002 , 71, 1607-14	6.8	17
36	HMG-CoA reductase inhibitor decreases small dense low-density lipoprotein and remnant-like particle cholesterol in patients with type-2 diabetes. <i>Life Sciences</i> , 2002 , 71, 2403-12	6.8	36
35	Cloning and characterization of a mammalian fatty acyl-CoA elongase as a lipogenic enzyme regulated by SREBPs. <i>Journal of Lipid Research</i> , 2002 , 43, 911-920	6.3	142
34	Dual regulation of mouse Delta(5)- and Delta(6)-desaturase gene expression by SREBP-1 and PPARalpha. <i>Journal of Lipid Research</i> , 2002 , 43, 107-14	6.3	206
33	Cloning and characterization of a mammalian fatty acyl-CoA elongase as a lipogenic enzyme regulated by SREBPs. <i>Journal of Lipid Research</i> , 2002 , 43, 911-20	6.3	123
32	Insulin up-regulates tumor necrosis factor-alpha production in macrophages through an extracellular-regulated kinase-dependent pathway. <i>Journal of Biological Chemistry</i> , 2001 , 276, 32531-7	5.4	44
31	Severe hypercholesterolemia, hypertriglyceridemia, and atherosclerosis in mice lacking both leptin and the low density lipoprotein receptor. <i>Journal of Biological Chemistry</i> , 2001 , 276, 37402-8	5.4	166
30	Neutralization of vascular endothelial growth factor prevents collagen-induced arthritis and ameliorates established disease in mice. <i>Biochemical and Biophysical Research Communications</i> , 2001 , 281, 562-8	3.4	104
29	Sterol regulatory element-binding proteins induce an entire pathway of cholesterol synthesis. <i>Biochemical and Biophysical Research Communications</i> , 2001 , 286, 176-83	3.4	164
28	Disease model: hyperinsulinemia and insulin resistance. Part A-targeted disruption of insulin signaling or glucose transport. <i>Trends in Molecular Medicine</i> , 2001 , 7, 320-2	11.5	24
27	Disease model: hyperinsulinemia and insulin resistance. Part B--polygenic and other animal models. <i>Trends in Molecular Medicine</i> , 2001 , 7, 373-6	11.5	7
26	Elevated levels of vascular endothelial growth factor in the sera of patients with rheumatoid arthritis correlation with disease activity. <i>Life Sciences</i> , 2001 , 69, 1861-9	6.8	47
25	Ibuprofen-related hypoglycemia in a patient receiving sulfonylurea. <i>Annals of Internal Medicine</i> , 2001 , 134, 344	8	16
24	Refractive changes in diabetic patients during intensive glycaemic control. <i>British Journal of Ophthalmology</i> , 2000 , 84, 1097-102	5.5	64
23	Enhancement of glucose transport by vascular endothelial growth factor in retinal endothelial cells. <i>Investigative Ophthalmology and Visual Science</i> , 2000 , 41, 1876-84		57

22	Increased mutant frequency and altered mutation spectrum of the lacI transgene in Wilson disease rats with hepatitis. <i>Cancer Research</i> , 2000 , 60, 5080-6	10.1	4
21	Isolation and characterization of the ATF2 gene encoding alcohol acetyltransferase II in the bottom fermenting yeast <i>Saccharomyces pastorianus</i> . <i>Yeast</i> , 1999 , 15, 409-17	3.4	37
20	Effects of intraocular or systemic administration of neutralizing antibody against vascular endothelial growth factor on the murine experimental model of retinopathy. <i>Life Sciences</i> , 1999 , 65, 2573-80	6.8	28
19	Increase in serum vascular endothelial growth factor levels during altitude training. <i>Acta Physiologica Scandinavica</i> , 1998 , 162, 455-9		48
18	Hypoxia and endothelin-1 induce VEGF production in human vascular smooth muscle cells. <i>Life Sciences</i> , 1998 , 63, 477-84	6.8	48
17	Acute effects of thromboxane dual blocker (KDI-792) on different portions of lower limb blood flow--a study using Doppler ultrasonography and laser Doppler flowmetry in type 2 diabetic patients. <i>Prostaglandins</i> , 1997 , 53, 395-409		2
16	Ocular vascular endothelial growth factor levels in diabetic rats are elevated before observable retinal proliferative changes. <i>Diabetologia</i> , 1997 , 40, 726-30	10.3	115
15	Primary adrenal lymphoma presenting as Addisonian crisis. Pitfalls in the diagnosis of bilateral adrenal swelling. <i>Hormone and Metabolic Research</i> , 1996 , 28, 116	3.1	11
14	Effects of high glucose concentration and a thromboxane synthase inhibitor on the production of thromboxane A2 and prostaglandin I2 and E2 by retinal endothelial cells. <i>Life Sciences</i> , 1996 , 58, 239-43	6.8	11
13	Progesterone induces vascular endothelial growth factor on retinal pigment epithelial cells in culture. <i>Life Sciences</i> , 1996 , 59, 21-5	6.8	24
12	Vascular endothelial growth factor is induced by long-term high glucose concentration and up-regulated by acute glucose deprivation in cultured bovine retinal pigmented epithelial cells. <i>Biochemical and Biophysical Research Communications</i> , 1996 , 221, 193-8	3.4	96
11	Acute effect of beraprost sodium on lower limb circulation in patients with non-insulin-dependent diabetes mellitus-evaluation by color Doppler ultrasonography and laser cutaneous blood flowmetry. <i>Prostaglandins</i> , 1996 , 52, 375-84		5
10	Predicting long-term glycemic control of post-educational type II diabetic patients by evaluating serum 1,5-anhydroglucitol levels. <i>Diabetes Research and Clinical Practice</i> , 1996 , 34, 83-8	7.4	5
9	Vascular endothelial growth factor level in aqueous humor of diabetic patients with rubeotic glaucoma is markedly elevated. <i>Diabetes Care</i> , 1996 , 19, 1306-7	14.6	42
8	Long-term effects of eicosapentaenoic acid on diabetic peripheral neuropathy and serum lipids in patients with type II diabetes mellitus. <i>Journal of Diabetes and Its Complications</i> , 1996 , 10, 280-7	3.2	46
7	Radioiodinated metaiodobenzylguanidine scintigraphy for pheochromocytoma. A false-positive case of adrenocortical adenoma and literature review. <i>Hormone Research</i> , 1996 , 46, 138-42		22
6	Transient nephrogenic diabetes insipidus accompanied by possible psychogenic polydipsia. <i>Hormone Research</i> , 1995 , 44, 193-6		1
5	Efficacy of Ibudilast on lower limb circulation of diabetic patients with minimally impaired baseline flow: a study using color Doppler ultrasonography and laser Doppler flowmetry. <i>Angiology</i> , 1995 , 46, 699-703	2.1	5

4	The effect of nilvadipine on bloodflow in the dorsal pedis artery in type 2 diabetic patients--a study using duplex Doppler ultrasonography. <i>Postgraduate Medical Journal</i> , 1995 , 71, 613-6	2	2
3	Restoration of myo-inositol uptake by eicosapentaenoic acid in human skin fibroblasts cultured in high-glucose medium. <i>Life Sciences</i> , 1995 , 57, PL71-4	6.8	2
2	Diabetes care in emergency settings. <i>Diabetes Care</i> , 1995 , 18, 1310-1	14.6	5
1	Syndrome of inappropriate secretion of antidiuretic hormone (SIADH) and Gerhardt syndrome associated with Shy-Drager syndrome. <i>Internal Medicine</i> , 1994 , 33, 773-8	1.1	12