

Edward L Lecluyse

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/3424885/edward-l-lecluyse-publications-by-year.pdf>

Version: 2024-04-24

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

83

papers

8,851

citations

48

h-index

83

g-index

83

ext. papers

9,480

ext. citations

4.4

avg, IF

5.7

L-index

#	Paper	IF	Citations
83	Hepatocyte size fractionation allows dissection of human liver zonation. <i>Journal of Cellular Physiology</i> , 2021 , 236, 5885-5894	7	0
82	Reprogramming of Human Hepatic Non-Parenchymal Cells: Step-by-Step Protocol. <i>Current Protocols in Stem Cell Biology</i> , 2020 , 53, e112	2.8	
81	Identifying qualitative differences in PPAR signaling networks in human and rat hepatocytes and their significance for next generation chemical risk assessment methods. <i>Toxicology in Vitro</i> , 2020 , 64, 104463	3.6	7
80	Long-Term Engineered Cultures of Primary Mouse Hepatocytes for Strain and Species Comparison Studies During Drug Development. <i>Gene Expression</i> , 2019 , 19, 199-214	3.4	4
79	Advancing nonclinical innovation and safety in pharmaceutical testing. <i>Drug Discovery Today</i> , 2019 , 24, 624-628	8.8	3
78	Liver Structure and Microanatomy 2018 ,		
77	Co-culture of Hepatocytes and Kupffer Cells as an In Vitro Model of Inflammation and Drug-Induced Hepatotoxicity. <i>Journal of Pharmaceutical Sciences</i> , 2016 , 105, 950-964	3.9	58
76	Nuclear Receptor-Mediated Gene Expression Changes in a Human Hepatic Micropatterned Coculture Model After Treatment with Hepatotoxic Compounds. <i>Applied in Vitro Toxicology</i> , 2016 , 2, 8-16	1.3	
75	The Promise of New Technologies to Reduce, Refine, or Replace Animal Use while Reducing Risks of Drug Induced Liver Injury in Pharmaceutical Development. <i>ILAR Journal</i> , 2016 , 57, 186-211	1.7	30
74	Editor's Highlight: Modeling Compound-Induced Fibrogenesis In Vitro Using Three-Dimensional Bioprinted Human Liver Tissues. <i>Toxicological Sciences</i> , 2016 , 154, 354-367	4.4	97
73	Differential Effects of Trovafloxacin on TNF- α and IL-6 Profiles in a Rat Hepatocyte-Kupffer Cell Coculture System. <i>Applied in Vitro Toxicology</i> , 2015 , 1, 45-54	1.3	11
72	A map of the PPAR transcription regulatory network for primary human hepatocytes. <i>Chemico-Biological Interactions</i> , 2014 , 209, 14-24	5	79
71	A micropatterned hepatocyte coculture model for assessment of liver toxicity using high-content imaging analysis. <i>Assay and Drug Development Technologies</i> , 2014 , 12, 16-27	2.1	37
70	Development of 3D dynamic flow model of human liver and its application to prediction of metabolic clearance of 7-ethoxycoumarin. <i>Tissue Engineering - Part C: Methods</i> , 2014 , 20, 641-51	2.9	13
69	Liver biomarker and in vitro assessment confirm the hepatic origin of aminotransferase elevations lacking histopathological correlate in beagle dogs treated with GABA _A receptor antagonist NP260. <i>Toxicology and Applied Pharmacology</i> , 2014 , 277, 131-7	4.6	20
68	Bioactivation and toxicity of acetaminophen in a rat hepatocyte micropatterned coculture system. <i>Journal of Biochemical and Molecular Toxicology</i> , 2013 , 27, 471-8	3.4	8
67	Recent advances in 2D and 3D in vitro systems using primary hepatocytes, alternative hepatocyte sources and non-parenchymal liver cells and their use in investigating mechanisms of hepatotoxicity, cell signaling and ADME. <i>Archives of Toxicology</i> , 2013 , 87, 1315-530	5.8	837

66	Early identification of clinically relevant drug interactions with the human bile salt export pump (BSEP/ABCB11). <i>Toxicological Sciences</i> , 2013 , 136, 328-43	4.4	113
65	Knockout of the aryl hydrocarbon receptor results in distinct hepatic and renal phenotypes in rats and mice. <i>Toxicology and Applied Pharmacology</i> , 2013 , 272, 503-18	4.6	56
64	models for liver toxicity testing. <i>Toxicology Research</i> , 2013 , 2, 23-39	2.6	304
63	Long-term stability of primary rat hepatocytes in micropatterned cocultures. <i>Journal of Biochemical and Molecular Toxicology</i> , 2013 , 27, 204-12	3.4	50
62	Organotypic liver culture models: meeting current challenges in toxicity testing. <i>Critical Reviews in Toxicology</i> , 2012 , 42, 501-48	5.7	252
61	A comprehensive evaluation of metabolic activity and intrinsic clearance in suspensions and monolayer cultures of cryopreserved primary human hepatocytes. <i>Journal of Pharmaceutical Sciences</i> , 2012 , 101, 3989-4002	3.9	62
60	Cross-species comparisons of transcriptomic alterations in human and rat primary hepatocytes exposed to 2,3,7,8-tetrachlorodibenzo-p-dioxin. <i>Toxicological Sciences</i> , 2012 , 127, 199-215	4.4	58
59	Human and rat primary hepatocyte CYP1A1 and 1A2 induction with 2,3,7,8-tetrachlorodibenzo-p-dioxin, 2,3,7,8-tetrachlorodibenzofuran, and 2,3,4,7,8-pentachlorodibenzofuran. <i>Toxicological Sciences</i> , 2010 , 118, 224-35	4.4	38
58	Xenobiotic-metabolizing enzyme and transporter gene expression in primary cultures of human hepatocytes modulated by ToxCast chemicals. <i>Journal of Toxicology and Environmental Health - Part B: Critical Reviews</i> , 2010 , 13, 329-46	8.6	47
57	Incorporating human dosimetry and exposure into high-throughput in vitro toxicity screening. <i>Toxicological Sciences</i> , 2010 , 117, 348-58	4.4	189
56	Isolation and culture of primary hepatocytes from resected human liver tissue. <i>Methods in Molecular Biology</i> , 2010 , 640, 57-82	1.4	99
55	Evaluation of an in vitro toxicogenetic mouse model for hepatotoxicity. <i>Toxicology and Applied Pharmacology</i> , 2010 , 249, 208-16	4.6	21
54	Regulation of cytochrome P450 2C9 expression in primary cultures of human hepatocytes. <i>Journal of Biochemical and Molecular Toxicology</i> , 2009 , 23, 43-58	3.4	35
53	Primary human hepatocyte culture for HCV study. <i>Methods in Molecular Biology</i> , 2009 , 510, 373-82	1.4	12
52	Response to Comments on Anti-Influenza Prodrug Oseltamivir Is Activated by Carboxylesterase Human Carboxylesterase 1, and the Activation Is Inhibited by Antiplatelet Agent Clopidogrel <i>Journal of Pharmacology and Experimental Therapeutics</i> , 2007 , 322, 424-425	4.7	1
51	Relative activation of human pregnane X receptor versus constitutive androstane receptor defines distinct classes of CYP2B6 and CYP3A4 inducers. <i>Journal of Pharmacology and Experimental Therapeutics</i> , 2007 , 320, 72-80	4.7	251
50	Temporal kinetics and concentration-response relationships for induction of CYP1A, CYP2B, and CYP3A in primary cultures of beagle dog hepatocytes. <i>Journal of Biochemical and Molecular Toxicology</i> , 2006 , 20, 69-78	3.4	14
49	Cloning, tissue expression, and regulation of beagle dog CYP4A genes. <i>Toxicological Sciences</i> , 2006 , 92, 356-67	4.4	11

48	Anti-influenza prodrug oseltamivir is activated by carboxylesterase human carboxylesterase 1, and the activation is inhibited by antiplatelet agent clopidogrel. <i>Journal of Pharmacology and Experimental Therapeutics</i> , 2006 , 319, 1477-84	4-7	223
47	Differential regulation of hepatic CYP2B6 and CYP3A4 genes by constitutive androstane receptor but not pregnane X receptor. <i>Journal of Pharmacology and Experimental Therapeutics</i> , 2006 , 317, 1200-9	4-7	154
46	Antiplatelet agents aspirin and clopidogrel are hydrolyzed by distinct carboxylesterases, and clopidogrel is transesterificated in the presence of ethyl alcohol. <i>Journal of Pharmacology and Experimental Therapeutics</i> , 2006 , 319, 1467-76	4-7	199
45	Differential UGT1A1 induction by chrysin in primary human hepatocytes and HepG2 Cells. <i>Journal of Pharmacology and Experimental Therapeutics</i> , 2005 , 315, 1256-64	4-7	40
44	Isolation and culture of primary human hepatocytes. <i>Methods in Molecular Biology</i> , 2005 , 290, 207-29	1-4	106
43	Modulation of UDP-glucuronosyltransferase 1A1 in primary human hepatocytes by prototypical inducers. <i>Journal of Biochemical and Molecular Toxicology</i> , 2005 , 19, 96-108	3-4	32
42	Human CYP2C8 is transcriptionally regulated by the nuclear receptors constitutive androstane receptor, pregnane X receptor, glucocorticoid receptor, and hepatic nuclear factor 4alpha. <i>Molecular Pharmacology</i> , 2005 , 68, 747-57	4-3	168
41	Kupffer cell-mediated IL-2 suppression of CYP3A activity in human hepatocytes. <i>Drug Metabolism and Disposition</i> , 2004 , 32, 359-63	4	86
40	Effects of avasimibe on cytochrome P450 2C9 expression in vitro and in vivo. <i>Drug Metabolism and Disposition</i> , 2004 , 32, 1370-6	4	25
39	The human sulfotransferase SULT1A1 gene is regulated in a synergistic manner by Sp1 and GA binding protein. <i>Molecular Pharmacology</i> , 2004 , 66, 1690-701	4-3	46
38	Human constitutive androstane receptor mediates induction of CYP2B6 gene expression by phenytoin. <i>Journal of Biological Chemistry</i> , 2004 , 279, 29295-301	5-4	115
37	Interindividual variation in the metabolism of arsenic in cultured primary human hepatocytes. <i>Toxicology and Applied Pharmacology</i> , 2004 , 201, 166-77	4-6	73
36	P-glycoprotein expression, localization, and function in sandwich-cultured primary rat and human hepatocytes: relevance to the hepatobiliary disposition of a model opioid peptide. <i>Pharmaceutical Research</i> , 2004 , 21, 1294-302	4-5	126
35	Regulation of CYP2B6 in primary human hepatocytes by prototypical inducers. <i>Drug Metabolism and Disposition</i> , 2004 , 32, 348-58	4	166
34	Glucocorticoid receptor enhancement of pregnane X receptor-mediated CYP2B6 regulation in primary human hepatocytes. <i>Drug Metabolism and Disposition</i> , 2003 , 31, 620-30	4	82
33	Role of orphan nuclear receptors in the regulation of drug-metabolising enzymes. <i>Clinical Pharmacokinetics</i> , 2003 , 42, 1331-57	6.2	212
32	Selenium compounds modulate the activity of recombinant rat AsIII-methyltransferase and the methylation of arsenite by rat and human hepatocytes. <i>Chemical Research in Toxicology</i> , 2003 , 16, 261-5	4	74
31	Comparative effects of thiazolidinediones on in vitro P450 enzyme induction and inhibition. <i>Drug Metabolism and Disposition</i> , 2003 , 31, 439-46	4	104

30	Avasimibe induces CYP3A4 and multiple drug resistance protein 1 gene expression through activation of the pregnane X receptor. <i>Journal of Pharmacology and Experimental Therapeutics</i> , 2003 , 306, 1027-34	4.7	58
29	Effects of prototypical microsomal enzyme inducers on cytochrome P450 expression in cultured human hepatocytes. <i>Drug Metabolism and Disposition</i> , 2003 , 31, 421-31	4	295
28	CYP3A induction by N-hydroxyformamide tumor necrosis factor-alpha converting enzyme/matrix metalloproteinase inhibitors use of a pregnane X receptor activation assay and primary hepatocyte culture for assessing induction potential in humans. <i>Drug Metabolism and Disposition</i> , 2003 , 31, 870-7	4	9
27	A novel distal enhancer module regulated by pregnane X receptor/constitutive androstane receptor is essential for the maximal induction of CYP2B6 gene expression. <i>Journal of Biological Chemistry</i> , 2003 , 278, 14146-52	5.4	176
26	Regulation of human CYP2C9 by the constitutive androstane receptor: discovery of a new distal binding site. <i>Molecular Pharmacology</i> , 2002 , 62, 737-46	4.3	134
25	The effect of cyclophosphamide with and without dexamethasone on cytochrome P450 3A4 and 2B6 in human hepatocytes. <i>Drug Metabolism and Disposition</i> , 2002 , 30, 814-22	4	78
24	Regulation of cell morphology and cytochrome P450 expression in human hepatocytes by extracellular matrix and cell-cell interactions. <i>Cell and Tissue Research</i> , 2001 , 306, 85-99	4.2	165
23	Optimization of culture conditions for determining hepatobiliary disposition of taurocholate in sandwich-cultured rat hepatocytes. <i>In Vitro Cellular and Developmental Biology - Animal</i> , 2001 , 37, 380-5	2.6	34
22	Pregnane X receptor: molecular basis for species differences in CYP3A induction by xenobiotics. <i>Chemico-Biological Interactions</i> , 2001 , 134, 283-9	5	164
21	Human hepatocyte culture systems for the in vitro evaluation of cytochrome P450 expression and regulation. <i>European Journal of Pharmaceutical Sciences</i> , 2001 , 13, 343-68	5.1	403
20	OPTIMIZATION OF CULTURE CONDITIONS FOR DETERMINING HEPATOBILIARY DISPOSITION OF TAUROCHOLATE IN SANDWICH-CULTURED RAT HEPATOCYTES. <i>In Vitro Cellular and Developmental Biology - Animal</i> , 2001 , 37, 380	2.6	14
19	Application of cDNA microarray to the study of arsenic-induced liver diseases in the population of Guizhou, China. <i>Toxicological Sciences</i> , 2001 , 59, 185-92	4.4	87
18	Expression and regulation of cytochrome P450 enzymes in primary cultures of human hepatocytes. <i>Journal of Biochemical and Molecular Toxicology</i> , 2000 , 14, 177-88	3.4	161
17	In vivo and in vitro induction of human cytochrome P4503A4 by dexamethasone. <i>Clinical Pharmacology and Therapeutics</i> , 2000 , 68, 356-66	6.1	110
16	Comparative toxicity of trivalent and pentavalent inorganic and methylated arsenicals in rat and human cells. <i>Archives of Toxicology</i> , 2000 , 74, 289-99	5.8	787
15	The pregnane X receptor: a promiscuous xenobiotic receptor that has diverged during evolution. <i>Molecular Endocrinology</i> , 2000 , 14, 27-39		565
14	Biliary excretion in primary rat hepatocytes cultured in a collagen-sandwich configuration. <i>American Journal of Physiology - Renal Physiology</i> , 1999 , 277, G12-21	5.1	73
13	Dodecylphosphocholine-mediated enhancement of paracellular permeability and cytotoxicity in Caco-2 cell monolayers. <i>Journal of Pharmaceutical Sciences</i> , 1999 , 88, 1161-8	3.9	77

12	Metabolism of arsenic in primary cultures of human and rat hepatocytes. <i>Chemical Research in Toxicology</i> , 1999 , 12, 560-5	4	118
11	Partial maintenance of taurocholate uptake by adult rat hepatocytes cultured in a collagen sandwich configuration. <i>Pharmaceutical Research</i> , 1998 , 15, 1533-9	4.5	69
10	In vitro models for selection of development candidates. Permeability studies to define mechanisms of absorption enhancement. <i>Advanced Drug Delivery Reviews</i> , 1997 , 23, 163-183	18.5	41
9	Sulfotransferase gene expression in primary cultures of rat hepatocytes. <i>Biochemical Pharmacology</i> , 1996 , 52, 1621-30	6	34
8	Strategies for restoration and maintenance of normal hepatic structure and function in long-term cultures of rat hepatocytes. <i>Advanced Drug Delivery Reviews</i> , 1996 , 22, 133-186	18.5	162
7	Cultured rat hepatocytes. <i>Pharmaceutical Biotechnology</i> , 1996 , 8, 121-59		84
6	Enhanced bioavailability of cefoxitin using palmitoylcarnitine. II. Use of directly compressed tablet formulations in the rat and dog. <i>Pharmaceutical Research</i> , 1993 , 10, 1516-20	4.5	13
5	Simultaneous in vitro measurement of intestinal tissue permeability and transepithelial electrical resistance (TEER) using Sweetana-Grass diffusion cells. <i>Pharmaceutical Research</i> , 1992 , 9, 316-9	4.5	24
4	Enhanced bioavailability of cefoxitin using palmitoyl L-carnitine. I. Enhancer activity in different intestinal regions. <i>Pharmaceutical Research</i> , 1992 , 9, 191-4	4.5	16
3	Relationship between drug absorption enhancing activity and membrane perturbing effects of acylcarnitines. <i>Pharmaceutical Research</i> , 1991 , 8, 84-7	4.5	50
2	Species Differences in Receptor-Mediated Gene Regulation71-98		1
1	Cytochrome P450 Induction265-314		1