

Humberto Milani

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/3428609/humberto-milani-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

68
papers

1,180
citations

19
h-index

30
g-index

69
ext. papers

1,358
ext. citations

3.5
avg, IF

4.32
L-index

#	Paper	IF	Citations
68	Positive effects of roflumilast on behavior, neuroinflammation, and white matter injury in mice with global cerebral ischemia. <i>Behavioural Pharmacology</i> , 2021 , 32, 459-471	2.4	1
67	Roflumilast protects against spatial memory impairments and exerts anti-inflammatory effects after transient global cerebral ischemia. <i>European Journal of Neuroscience</i> , 2021 , 53, 1171-1188	3.5	4
66	Differential contribution of CB1, CB2, 5-HT1A, and PPAR- α receptors to cannabidiol effects on ischemia-induced emotional and cognitive impairments. <i>European Journal of Neuroscience</i> , 2021 , 53, 1738-1751	3.5	5
65	Cannabidiol Confers Neuroprotection in Rats in a Model of Transient Global Cerebral Ischemia: Impact of Hippocampal Synaptic Neuroplasticity. <i>Molecular Neurobiology</i> , 2021 , 58, 5338-5355	6.2	1
64	Activation of 5-HT postsynaptic receptors by NLX-101 results in functional recovery and an increase in neuroplasticity in mice with brain ischemia. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2020 , 99, 109832	5.5	19
63	Glycemic homeostasis and hepatic metabolism are modified in rats with global cerebral ischemia. <i>Biochimica Et Biophysica Acta - Molecular Basis of Disease</i> , 2020 , 1866, 165934	6.9	4
62	Anxiolytic-like and proneurogenic effects of Trichilia catigua ethyl-acetate fraction in mice with cerebral ischemia. <i>Revista Brasileira De Farmacognosia</i> , 2019 , 29, 613-620	2	1
61	Postischemic fish oil treatment restores dendritic integrity and synaptic proteins levels after transient, global cerebral ischemia in rats. <i>Journal of Chemical Neuroanatomy</i> , 2019 , 101, 101683	3.2	8
60	DHA-Enriched Formulations as a Promising Strategy for the Treatment of Hypoxic/Ischemic Brain Injury 2019 , 391-401		
59	Longitudinal modeling using log-gamma mixed model: case of memory deterioration after chronic cerebral hypoperfusion associated with diabetes in rats. <i>Acta Scientiarum - Technology</i> , 2019 , 41, 35789	0.5	
58	Longitudinal modeling using log-gamma mixed model: case of memory deterioration after chronic cerebral hypoperfusion associated with diabetes in rats. <i>Acta Scientiarum - Technology</i> , 2019 , 41, 35789	0.5	1
57	Effects of Cannabidiol on Diabetes Outcomes and Chronic Cerebral Hypoperfusion Comorbidities in Middle-Aged Rats. <i>Neurotoxicity Research</i> , 2019 , 35, 463-474	4.3	10
56	Ethyl-acetate fraction of Trichilia catigua protects against oxidative stress and neuroinflammation after cerebral ischemia/reperfusion. <i>Journal of Ethnopharmacology</i> , 2018 , 221, 109-118	5	16
55	Pioglitazone reduces mortality, prevents depressive-like behavior, and impacts hippocampal neurogenesis in the 6-OHDA model of Parkinson's disease in rats. <i>Experimental Neurology</i> , 2018 , 300, 188-200	5.7	28
54	Ethyl-acetate fraction of Trichilia catigua restores long-term retrograde memory and reduces oxidative stress and inflammation after global cerebral ischemia in rats. <i>Behavioural Brain Research</i> , 2018 , 337, 173-182	3.4	15
53	Roflumilast promotes memory recovery and attenuates white matter injury in aged rats subjected to chronic cerebral hypoperfusion. <i>Neuropharmacology</i> , 2018 , 138, 360-370	5.5	24
52	Cognitive, neurohistological and mortality outcomes following the four-vessel occlusion/internal carotid artery model of chronic cerebral hypoperfusion: The impact of diabetes and aging. <i>Behavioural Brain Research</i> , 2018 , 339, 169-178	3.4	5

51	Postischemic fish oil treatment confers task-dependent memory recovery. <i>Physiology and Behavior</i> , 2017 , 177, 196-207	3.5	10
50	Depletion of 5 hydroxy-triptamine (5-HT) affects the antidepressant-like effect of neuronal nitric oxide synthase inhibitor in mice. <i>Neuroscience Letters</i> , 2017 , 656, 131-137	3.3	9
49	The phosphodiesterase type 2 inhibitor BAY 60-7550 reverses functional impairments induced by brain ischemia by decreasing hippocampal neurodegeneration and enhancing hippocampal neuronal plasticity. <i>European Journal of Neuroscience</i> , 2017 , 45, 510-520	3.5	15
48	Myricitrin induces antidepressant-like effects and facilitates adult neurogenesis in mice. <i>Behavioural Brain Research</i> , 2017 , 316, 59-65	3.4	19
47	Cannabidiol reduces neuroinflammation and promotes neuroplasticity and functional recovery after brain ischemia. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2017 , 75, 94-105	5.5	69
46	Robust and enduring atorvastatin-mediated memory recovery following the 4-vessel occlusion/internal carotid artery model of chronic cerebral hypoperfusion in middle-aged rats. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2016 , 65, 179-87	5.5	5
45	Influence of single and repeated cannabidiol administration on emotional behavior and markers of cell proliferation and neurogenesis in non-stressed mice. <i>Progress in Neuro-Psychopharmacology and Biological Psychiatry</i> , 2016 , 64, 27-34	5.5	78
44	Mixed models in cerebral ischemia study. <i>Acta Scientiarum - Technology</i> , 2016 , 38, 345	0.5	
43	Postischemic fish oil treatment restores long-term retrograde memory and dendritic density: An analysis of the time window of efficacy. <i>Behavioural Brain Research</i> , 2016 , 311, 425-439	3.4	11
42	Rolipram improves cognition, reduces anxiety- and despair-like behaviors and impacts hippocampal neuroplasticity after transient global cerebral ischemia. <i>Neuroscience</i> , 2016 , 326, 69-83	3.9	45
41	Trichilia catigua ethyl-acetate fraction protects against cognitive impairments and hippocampal cell death induced by bilateral common carotid occlusion in mice. <i>Journal of Ethnopharmacology</i> , 2015 , 172, 232-7	5	14
40	Dopamine in the nucleus accumbens core, but not shell, increases during signaled food reward and decreases during delayed extinction. <i>Neurobiology of Learning and Memory</i> , 2015 , 123, 125-39	3.1	9
39	4-hydroxy-3-methoxy-acetophenone-mediated long-lasting memory recovery, hippocampal neuroprotection, and reduction of glial cell activation after transient global cerebral ischemia in rats. <i>Journal of Neuroscience Research</i> , 2015 , 93, 1240-9	4.4	6
38	Cilostazol but not sildenafil prevents memory impairment after chronic cerebral hypoperfusion in middle-aged rats. <i>Behavioural Brain Research</i> , 2015 , 283, 61-8	3.4	15
37	Fish oil prevents oxidative stress and exerts sustained anti-amnesic effect after global cerebral ischemia. <i>CNS and Neurological Disorders - Drug Targets</i> , 2015 , 14, 400-10	2.6	12
36	Phosphodiesterase Inhibition as a Therapeutic Target for Brain Ischemia. <i>CNS and Neurological Disorders - Drug Targets</i> , 2015 , 14, 1012-23	2.6	4
35	Protective effects of cannabidiol against hippocampal cell death and cognitive impairment induced by bilateral common carotid artery occlusion in mice. <i>Neurotoxicity Research</i> , 2014 , 26, 307-16	4.3	60
34	P.6.a.017 Chronic ethanol exposure increases inducible nitric oxide synthase expression in the dorsolateral periaqueductal gray matter of rats. <i>European Neuropsychopharmacology</i> , 2014 , 24, S662-S663 ^{1,2}		

33	Fish oil provides a sustained anti-amnesic effect after acute, transient forebrain ischemia but not after chronic cerebral hypoperfusion in middle-aged rats. <i>Behavioural Brain Research</i> , 2014 , 265, 101-10	3.4	12
32	Effects of nitric oxide synthase inhibition in the dorsolateral periaqueductal gray matter on ethanol withdrawal-induced anxiety-like behavior in rats. <i>Psychopharmacology</i> , 2013 , 228, 487-98	4.7	20
31	Cognitive impairment and persistent anxiety-related responses following bilateral common carotid artery occlusion in mice. <i>Behavioural Brain Research</i> , 2013 , 249, 28-37	3.4	39
30	Neurohistological and behavioral changes following the four-vessel occlusion/internal carotid artery model of chronic cerebral hypoperfusion: comparison between normotensive and spontaneously hypertensive rats. <i>Behavioural Brain Research</i> , 2013 , 252, 214-21	3.4	24
29	Fish oil provides robust and sustained memory recovery after cerebral ischemia: influence of treatment regimen. <i>Physiology and Behavior</i> , 2013 , 119, 61-71	3.5	18
28	Sildenafil provides sustained neuroprotection in the absence of learning recovery following the 4-vessel occlusion/internal carotid artery model of chronic cerebral hypoperfusion in middle-aged rats. <i>Brain Research Bulletin</i> , 2013 , 90, 58-65	3.9	12
27	Eag1, Eag2, and SK3 potassium channel expression in the rat hippocampus after global transient brain ischemia. <i>Journal of Neuroscience Research</i> , 2012 , 90, 632-40	4.4	8
26	Subchronic administration of <i>Trichilia catigua</i> ethyl-acetate fraction promotes antidepressant-like effects and increases hippocampal cell proliferation in mice. <i>Journal of Ethnopharmacology</i> , 2012 , 143, 179-84	5	23
25	Time-course of neurodegeneration and memory impairment following the 4-vessel occlusion/internal carotid artery model of chronic cerebral hypoperfusion in middle-aged rats. <i>Behavioural Brain Research</i> , 2012 , 229, 340-8	3.4	18
24	Ethanol withdrawal activates nitric oxide-producing neurons in anxiety-related brain areas. <i>Alcohol</i> , 2011 , 45, 641-52	2.7	24
23	Acute, post-ischemic sensorimotor deficits correlate positively with infarct size but fail to predict its occurrence and magnitude after middle cerebral artery occlusion in rats. <i>Behavioural Brain Research</i> , 2011 , 216, 29-35	3.4	9
22	Middle-aged, but not young, rats develop cognitive impairment and cortical neurodegeneration following the four-vessel occlusion/internal carotid artery model of chronic cerebral hypoperfusion. <i>European Journal of Neuroscience</i> , 2011 , 34, 1131-40	3.5	17
21	Imipramine enhances cell proliferation and decreases neurodegeneration in the hippocampus after transient global cerebral ischemia in rats. <i>Neuroscience Letters</i> , 2010 , 470, 43-8	3.3	16
20	Sildenafil prevents mortality and reduces hippocampal damage after permanent, stepwise, 4-vessel occlusion in rats. <i>Brain Research Bulletin</i> , 2010 , 81, 631-40	3.9	11
19	Vitamin E improves learning performance and changes the expression of nitric oxide-producing neurons in the brains of diabetic rats. <i>Behavioural Brain Research</i> , 2010 , 210, 38-45	3.4	33
18	The cognitive and histopathological effects of chronic 4-vessel occlusion in rats depend on the set of vessels occluded and the age of the animals. <i>Behavioural Brain Research</i> , 2009 , 197, 378-87	3.4	17
17	Long-term treatment with fish oil prevents memory impairments but not hippocampal damage in rats subjected to transient, global cerebral ischemia. <i>Nutrition Research</i> , 2008 , 28, 798-808	4	45
16	Tacrolimus (FK506) reduces hippocampal damage but fails to prevent learning and memory deficits after transient, global cerebral ischemia in rats. <i>Pharmacology Biochemistry and Behavior</i> , 2007 , 88, 28-38	3.9	16

15	Sustained neuroprotection and facilitation of behavioral recovery by the Ginkgo biloba extract, EGb 761, after transient forebrain ischemia in rats. <i>Behavioural Brain Research</i> , 2006 , 174, 70-7	3.4	29
14	The Ginkgo biloba extract, EGb 761, fails to reduce brain infarct size in rats after transient, middle cerebral artery occlusion in conditions of unprevented, ischemia-induced fever. <i>Phytotherapy Research</i> , 2006 , 20, 438-43	6.7	9
13	Permanent, 3-stage, 4-vessel occlusion as a model of chronic and progressive brain hypoperfusion in rats: a neurohistological and behavioral analysis. <i>Behavioural Brain Research</i> , 2005 , 160, 312-22	3.4	23
12	Effect of tacrolimus (FK506) on ischemia-induced brain damage and memory dysfunction in rats. <i>Pharmacology Biochemistry and Behavior</i> , 2004 , 77, 607-15	3.9	12
11	A novel version of the 8-arm radial maze: effects of cerebral ischemia on learning and memory. <i>Journal of Neuroscience Methods</i> , 2004 , 132, 9-18	3	26
10	Validation of a simple and inexpensive method for the quantitation of infarct in the rat brain. <i>Brazilian Journal of Medical and Biological Research</i> , 2004 , 37, 511-21	2.8	9
9	Tacrolimus (FK506) reduces ischemia-induced hippocampal damage in rats: a 7- and 30-day study. <i>Brazilian Journal of Medical and Biological Research</i> , 2003 , 36, 495-502	2.8	19
8	Magnesium chloride alone or in combination with diazepam fails to prevent hippocampal damage following transient forebrain ischemia. <i>Brazilian Journal of Medical and Biological Research</i> , 1999 , 32, 1285-93	2.8	14
7	Loss of CA1 cells following global ischaemia correlates with spatial deficits in the circular platform task. <i>Journal of Neuroscience Methods</i> , 1998 , 80, 19-27	3	12
6	Interaction between recovery from behavioral asymmetries induced by hemivibrissotomy in the rat and the effects of apomorphine and amphetamine.. <i>Behavioral Neuroscience</i> , 1990 , 104, 470-476	2.1	16
5	Interaction between recovery from behavioral asymmetries induced by hemivibrissotomy in the rat and the effects of apomorphine and amphetamine. <i>Behavioral Neuroscience</i> , 1990 , 104, 470-6	2.1	5
4	Analysis of recovery from behavioral asymmetries induced by unilateral removal of vibrissae in the rat.. <i>Behavioral Neuroscience</i> , 1989 , 103, 1067-1074	2.1	50
3	Analysis of recovery from behavioral asymmetries induced by unilateral removal of vibrissae in the rat. <i>Behavioral Neuroscience</i> , 1989 , 103, 1067-74	2.1	12
2	GABA-benzodiazepine modulation of aversion in the medial hypothalamus of the rat. <i>Pharmacology Biochemistry and Behavior</i> , 1987 , 28, 21-7	3.9	56
1	Role of GABA in the anti-aversive action of anxiolytics. <i>Advances in Biochemical Psychopharmacology</i> , 1986 , 42, 79-86		3