

# Masahiro Osakabe

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3471005/publications.pdf>

Version: 2024-02-01

115  
papers

2,559  
citations

201674

27  
h-index

265206

42  
g-index

117  
all docs

117  
docs citations

117  
times ranked

1140  
citing authors

#	ARTICLE	IF	CITATIONS
1	Combination of target site mutation and associated CYPs confers high-level resistance to pyridaben in <i>Tetranychus urticae</i> . <i>Pesticide Biochemistry and Physiology</i> , 2022, 181, 105000.	3.6	12
2	freqPCR: Estimation of population allele frequency using qPCR $\hat{p}$ Cq measures from bulk samples. <i>Molecular Ecology Resources</i> , 2022, 22, 1380-1393.	4.8	3
3	A mutation in chitin synthase I associated with etoxazole resistance in the citrus red mite <i>Panonychus citri</i> (Acari: Tetranychidae) and its uneven geographical distribution in Japan. <i>Pest Management Science</i> , 2022, 78, 4028-4036.	3.4	3
4	Mechanisms underlying the impact and interaction of temperature and UV-B on the hatching of spider mite and phytoseiid mite eggs. <i>Pest Management Science</i> , 2022, 78, 4314-4323.	3.4	1
5	Strawberry pollen as a source of UV-B protection ingredients for the phytoseiid mite <i>Neoseiulus californicus</i> (Acari: Phytoseiidae). <i>Pest Management Science</i> , 2021, 77, 851-859.	3.4	9
6	Biological impact of ultraviolet-B radiation on spider mites and its application in integrated pest management. <i>Applied Entomology and Zoology</i> , 2021, 56, 139-155.	1.2	15
7	Up-regulation of pathogenesis-related genes in strawberry leaves treated with powdery mildew-suppressing ultraviolet irradiation. <i>Plant Pathology</i> , 2021, 70, 1378-1387.	2.4	8
8	Co-occurrence of subunit B and C mutations in respiratory complex II confers high resistance levels to pyflubumide and cyenopyrafen in the two-spotted spider mite <i>Tetranychus urticae</i> (Acari: Tetranychidae). <i>Pest Management Science</i> , 2021, 77, 5149-5157.	3.4	14
9	Molecular monitoring of <i>Neoseiulus californicus</i> released from sheltered slow-release sachets for spider mite control in a Japanese pear greenhouse. <i>Experimental and Applied Acarology</i> , 2020, 80, 203-214.	1.6	3
10	QTL mapping using microsatellite linkage reveals target-site mutations associated with high levels of resistance against three mitochondrial complex II inhibitors in <i>Tetranychus urticae</i> . <i>Insect Biochemistry and Molecular Biology</i> , 2020, 123, 103410.	2.7	36
11	Effects of Growth Phase and Ultraviolet-B Pretreatment in Perilla Leaves on the Two-Spotted Spider Mite. <i>Environmental Entomology</i> , 2020, 49, 886-894.	1.4	2
12	Dose-Response and Temperature Dependence of the Mortality of Spider Mite and Predatory Mite Eggs Caused by Daily Nighttime Ultraviolet-B Irradiation. <i>Photochemistry and Photobiology</i> , 2020, 96, 877-882.	2.5	6
13	Diagnostic prediction of acaricide resistance gene frequency using quantitative real-time PCR with resistance allele-specific primers in the two-spotted spider mite <i>Tetranychus urticae</i> population (Acari: Tetranychidae). <i>Journal of Economic Entomology</i> , 2020, 53, 1078-1084.	1.7	1
14	Convergent evolution of cytochrome P450s underlies independent origins of keto-carotenoid pigmentation in animals. <i>Proceedings of the Royal Society B: Biological Sciences</i> , 2019, 286, 20191039.	2.6	28
15	Mechanism of acequinocyl resistance and cross-resistance to bifenazate in the two-spotted spider mite, <i>Tetranychus urticae</i> (Acari: Tetranychidae). <i>Applied Entomology and Zoology</i> , 2019, 54, 421-427.	1.2	10
16	Effects of Low Temperature on Spider Mite Control by Intermittent Ultraviolet-B Irradiation for Practical Use in Greenhouse Strawberries. <i>Environmental Entomology</i> , 2018, 47, 140-147.	1.4	20
17	A Pollen Diet Confers Ultraviolet-B Resistance in Phytoseiid Mites by Providing Antioxidants. <i>Frontiers in Ecology and Evolution</i> , 2018, 6, .	2.2	11
18	Combination of restriction endonuclease digestion with the $\hat{p}$ Ct method in real-time PCR to monitor etoxazole resistance allele frequency in the two-spotted spider mite. <i>Pesticide Biochemistry and Physiology</i> , 2017, 139, 1-8.	3.6	36

#	ARTICLE	IF	CITATIONS
19	A mutation in the PSST homologue of complex I (NADH:ubiquinone oxidoreductase) from <i>Tetranychus urticae</i> is associated with resistance to METI acaricides. <i>Insect Biochemistry and Molecular Biology</i> , 2017, 80, 79-90.	2.7	82
20	Photo-enzymatic repair of UVB-induced DNA damage in the two-spotted spider mite <i>Tetranychus urticae</i> . <i>Experimental and Applied Acarology</i> , 2017, 71, 15-34.	1.6	27
21	Developmental Phase-Specific Mortality After Ultraviolet-B Radiation Exposure in the Two-Spotted Spider Mite. <i>Environmental Entomology</i> , 2017, 46, 1448-1455.	1.4	14
22	Disruption of a horizontally transferred phytoene desaturase abolishes carotenoid accumulation and diapause in <i>Tetranychus urticae</i> . <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2017, 114, E5871-E5880.	7.1	114
23	Antioxidant Protection by Astaxanthin in the Citrus Red Mite (Acari: Tetranychidae). <i>Environmental Entomology</i> , 2017, 46, 1143-1150.	1.4	22
24	Protocols for the delivery of small molecules to the two-spotted spider mite, <i>Tetranychus urticae</i> . <i>PLoS ONE</i> , 2017, 12, e0180658.	2.5	40
25	Low level of polyandry constrains phenotypic plasticity of male body size in mites. <i>PLoS ONE</i> , 2017, 12, e0188924.	2.5	7
26	Difference between Nighttime and Daytime UV-B Irradiation with Respect to the Extent of Damage to Perilla Leaves. <i>Horticulture Journal</i> , 2017, 86, 349-356.	0.8	5
27	Construction of a Spraying System to Replace the Rotary Distributing Sprayer. <i>Japanese Journal of Applied Entomology and Zoology</i> , 2017, 61, 192-194.	0.1	6
28	Ultimate Drivers and Proximate Correlates of Polyandry in Predatory Mites. <i>PLoS ONE</i> , 2016, 11, e0154355.	2.5	9
29	Phytoseiid mites under environmental stress. <i>Biological Control</i> , 2016, 96, 120-134.	3.0	75
30	Imaginal Feeding for Progression of Diapause Phenotype in the Two-Spotted Spider Mite (Acari: Tetranychidae). <i>Journal of Economic Entomology</i> , 2016, 49, 1075-1083.	1.4	37
31	Physical Control of Spider Mites Using Ultraviolet-B With Light Reflection Sheets in Greenhouse Strawberries. <i>Journal of Economic Entomology</i> , 2016, 109, 1758-1765.	1.8	42
32	Molecular analysis of cyenopyrafen resistance in the two-spotted spider mite <i>Tetranychus urticae</i> . <i>Pest Management Science</i> , 2016, 72, 103-112.	3.4	60
33	The effects of prestarvation diet on starvation tolerance of the predatory mite <i>Noseiulus californicus</i> (Acari: Phytoseiidae). <i>Physiological Entomology</i> , 2015, 40, 296-303.	1.5	4
34	Joint Effect of Solar UVB and Heat Stress on the Seasonal Change of Egg Hatching Success in the Herbivorous False Spider Mite (Acari: Tenuipalpidae). <i>Environmental Entomology</i> , 2015, 44, 1605-1613.	1.4	7
35	Suspension of Egg Hatching Caused by High Humidity and Submergence in Spider Mites. <i>Environmental Entomology</i> , 2015, 44, 1210-1219.	1.4	13
36	Effects of combination between web density and size of spider mite on predation by a generalist and a specialist phytoseiid mite. <i>Experimental and Applied Acarology</i> , 2015, 66, 219-225.	1.6	5

#	ARTICLE	IF	CITATIONS
37	Spectrum-Specific Uv Egg Damage and Dispersal Responses in the Phytoseiid Predatory Mite <i>Neoseiulus californicus</i> (Acari: Phytoseiidae). <i>Environmental Entomology</i> , 2014, 43, 787-794.	1.4	31
38	Factors affecting photoreactivation in UVB-irradiated herbivorous spider mite ( <i>Tetranychus urticae</i> ). <i>Experimental and Applied Acarology</i> , 2014, 63, 253-265.	1.6	36
39	Cross-resistance between cyenopyrafen and pyridaben in the twospotted spider mite <i>Tetranychus urticae</i> (Acari: Tetranychidae). <i>Pest Management Science</i> , 2014, 70, 1090-1096.	3.4	36
40	The Bunsen-Roscoe reciprocity law in ultraviolet-B-induced mortality of the two-spotted spider mite <i>Tetranychus urticae</i> . <i>Journal of Insect Physiology</i> , 2013, 59, 241-247.	2.0	44
41	Stellate hairs on leaves of a deciduous shrub <i>Viburnum erosum</i> var. <i>punctatum</i> (Adoxaceae) effectively protect <i>Brevipalpus obovatus</i> (Acari: Tenuipalpidae) eggs from the predator <i>Phytoseius nipponicus</i> (Acari: Phytoseiidae). <i>Experimental and Applied Acarology</i> , 2013, 60, 299-311.	1.6	7
42	PCR-RFLP Analysis for Identification of <i>Tetranychus</i> Spider Mite Species (Acari: Tetranychidae). <i>Journal of Insect Physiology</i> , 2013, 59, 241-247.	1.8	19
43	Geotaxis and leaf-surface preferences mitigate negative effects of a predatory mite on an herbivorous mite. <i>Experimental and Applied Acarology</i> , 2013, 59, 409-420.	1.6	4
44	Tolerance to Solar Ultraviolet-B Radiation in the Citrus Red Mite, An Upper Surface User of Host Plant Leaves. <i>Photochemistry and Photobiology</i> , 2013, 89, 424-431.	2.5	43
45	Effects of UV-B radiation on the reproduction of <i>Tetranychus ludeni</i> and the growth of melon plants. <i>Proceedings of the Kansai Plant Protection Society</i> , 2013, 55, 37-41.	0.1	6
46	DNA preparation method in eggs, immature stages, and diapausing females of <i>Tetranychus</i> spider mites (Acari: Tetranychidae) for diagnostic PCR-RFLP. <i>Applied Entomology and Zoology</i> , 2012, 47, 295-300.	1.2	2
47	Vulnerability and behavioral response to ultraviolet radiation in the components of a foliar mite predator system. <i>Die Naturwissenschaften</i> , 2012, 99, 1031-1038.	1.6	46
48	A comparison of the effects of gravity and the nutritional advantage of leaf surfaces on fecundity in the two-spotted spider mite (Acari: Tetranychidae). <i>Journal of the Acarological Society of Japan</i> , 2012, 21, 1-6.	0.2	5
49	Seasonal changes in the deleterious effects of solar ultraviolet-B radiation on eggs of the twospotted spider mite, <i>Tetranychus urticae</i> (Acari: Tetranychidae). <i>Applied Entomology and Zoology</i> , 2012, 47, 67-73.	1.2	46
50	Kanzawa spider mites acquire enemy-free space on a detrimental host plant, oleander. <i>Entomologia Experimentalis Et Applicata</i> , 2011, 138, 212-222.	1.4	14
51	Do plant mites commonly prefer the underside of leaves?. <i>Experimental and Applied Acarology</i> , 2011, 55, 25-38.	1.6	52
52	The predatory mite <i>Neoseiulus womersleyi</i> (Acari: Phytoseiidae) follows extracts of trails left by the two-spotted spider mite <i>Tetranychus urticae</i> (Acari: Tetranychidae). <i>Experimental and Applied Acarology</i> , 2010, 52, 111-118.	1.6	13
53	Spectrum-specific Damage and Solar Ultraviolet Radiation Avoidance in the Two-spotted Spider Mite. <i>Photochemistry and Photobiology</i> , 2010, 86, 925-932.	2.5	77
54	Acaricide resistance in spider mites. <i>Journal of Pesticide Sciences</i> , 2009, 34, 207-214.	1.4	2

#	ARTICLE	IF	CITATIONS
55	Development and Evaluation of Microsatellite Markers in <i>Tetranychus truncatus</i> Ehara (Acari: Tetranychidae). <i>Journal of Economic Entomology</i> , 2009, 42, 107-111.	0.2	3
56	Evolutionary Aspects of Acaricide-Resistance Development in Spider Mites. <i>Psyche: Journal of Entomology</i> , 2009, 2009, 1-11.	0.9	27
57	The fine-scale genetic structure of the two-spotted spider mite in a commercial greenhouse. <i>Experimental and Applied Acarology</i> , 2009, 47, 99-109.	1.6	26
58	Evidence of a high level of gene flow among apple trees in <i>Tetranychus urticae</i> . <i>Experimental and Applied Acarology</i> , 2009, 49, 281-290.	1.6	23
59	Do spider mite-infested plants and spider mite trails attract predatory mites?. <i>Ecological Research</i> , 2009, 24, 1173-1178.	1.5	21
60	Deleterious Effects of UV-B Radiation on Herbivorous Spider Mites: They Can Avoid It by Remaining on Lower Leaf Surfaces. <i>Environmental Entomology</i> , 2009, 38, 920-929.	1.4	102
61	Effects of wall structure and light intensity on the settlement of a predatory mite, <i>Euseius sojaensis</i> (Ehara) (Acari: Phytoseiidae). <i>Applied Entomology and Zoology</i> , 2009, 44, 81-84.	1.2	9
62	Pre-winter copulation enhances overwintering success of <i>Orius</i> females (Heteroptera: Anthracoridae). <i>Applied Entomology and Zoology</i> , 2009, 44, 47-52.	1.2	15
63	Aerodynamic advantages of upside down take-off for aerial dispersal in <i>Tetranychus</i> spider mites. <i>Experimental and Applied Acarology</i> , 2008, 44, 165-183.	1.6	28
64	Development, long-term survival, and the maintenance of fertility in <i>Neoseiulus californicus</i> (Acari: Tetranychidae). <i>Journal of Economic Entomology</i> , 2008, 41, 107-111.	1.6	28
65	Restriction Fragment Length Polymorphism Catalog for Molecular Identification of Japanese <i>Tetranychus</i> Spider Mites (Acari: Tetranychidae). <i>Journal of Economic Entomology</i> , 2008, 41, 1167-1175.	1.8	18
66	Restriction Fragment Length Polymorphism Catalog for Molecular Identification of Japanese <i>Tetranychus</i> Spider Mites (Acari: Tetranychidae). <i>Journal of Economic Entomology</i> , 2008, 41, 1167-1175.	1.8	23
67	Linkage Between One of the Polygenic Hexythiazox Resistance Genes and an Etoxazole Resistance Gene in the Twospotted Spider Mite (Acari: Tetranychidae). <i>Journal of Economic Entomology</i> , 2008, 41, 1704-1710.	1.8	21
68	Identification of spider mites (Acari: Tetranychidae) by DNA sequences: A case study in northern Vietnam. <i>International Journal of Acarology</i> , 2007, 33, 53-60.	0.7	38
69	Isolation and characterization of microsatellite loci in the two-spotted spider mite, <i>Tetranychus urticae</i> (Acari: Tetranychidae). <i>Molecular Ecology Notes</i> , 2007, 7, 290-292.	1.7	24
70	Host race formation in the Acari. <i>Experimental and Applied Acarology</i> , 2007, 42, 225-238.	1.6	89
71	Host plants utilized during the immature development of <i>Tetranychus kanzawai</i> (Acari: Tetranychidae) determine the preference of the adult females for the plants. <i>Journal of the Acarological Society of Japan</i> , 2007, 16, 121-127.	0.2	2
72	Effect of cytochrome P450 inhibitor, piperonyl butoxide, on survival of <i>Panonychus citri</i> (McGregor) (Acari: Tetranychidae) on citrus leaves. <i>Applied Entomology and Zoology</i> , 2006, 41, 487-491.	1.2	2

#	ARTICLE	IF	CITATIONS
73	Web-Mediated Interspecific Competition Among Spider Mites. <i>Journal of Economic Entomology</i> , 2006, 99, 678-684.	1.8	20
74	Amensalism via webs causes unidirectional shifts of dominance in spider mite communities. <i>Oecologia</i> , 2006, 150, 496-505.	2.0	30
75	Web-Mediated Interspecific Competition Among Spider Mites. <i>Journal of Economic Entomology</i> , 2006, 99, 678-684.	1.8	24
76	Why does the predatory mite <i>Neoseiulus womersleyi</i> Schicha (Acari: Phytoseiidae) prefer spider mite eggs to adults?. <i>Applied Entomology and Zoology</i> , 2005, 40, 675-678.	1.2	23
77	Prey preference of the predatory mite <i>Neoseiulus womersleyi</i> Schicha is determined by spider mite webs. <i>Journal of Applied Entomology</i> , 2005, 129, 336-339.	1.8	18
78	Significance of habitat type for the genetic population structure of <i>Panonychus citri</i> (Acari: Tetranychidae). <i>Journal of Applied Entomology and Zoology</i> , 2005, 40, 542-547.	1.6	25
79	Mating Strategies of <i>Tetranychus kanzawai</i> (Acari: Tetranychidae) in Relation to Mating Status of Females. <i>Annals of the Entomological Society of America</i> , 2005, 98, 625-628.	2.5	16
80	Spider mites assess predation risk by using the odor of injured conspecifics. <i>Journal of Chemical Ecology</i> , 2003, 29, 2609-2613.	1.8	21
81	Development of genetic differentiation and postzygotic isolation in experimental metapopulations of spider mites. <i>Experimental and Applied Acarology</i> , 2003, 31, 161-176.	1.6	6
82	Discrimination of four Japanese <i>Tetranychus</i> species (Acari: Tetranychidae) using PCR-RFLP of the inter-transcribed spacer region of nuclear ribosomal DNA.. <i>Applied Entomology and Zoology</i> , 2002, 37, 399-407.	1.2	25
83	Influence of Trap and Barrier Crops on Occurrence of and Damage by Stink Bugs and Lepidopterous Pod Borers in Soybean Fields.. <i>Japanese Journal of Applied Entomology and Zoology</i> , 2002, 46, 233-241.	0.1	20
84	Genetic Basis of Resistances to Chlorfenapyr and Etoxazole in the Two-Spotted Spider Mite (Acari: Tetranychidae). <i>Journal of Applied Entomology and Zoology</i> , 2002, 37, 542-547.	1.8	65
85	Which predatory mite can control both a dominant mite pest, <i>Tetranychus urticae</i> , and a latent mite pest, <i>Eotetranychus asiaticus</i> , on strawberry?. <i>Experimental and Applied Acarology</i> , 2002, 26, 219-230.	1.6	8
86	Phylogenetic analysis of green and red forms of the two-spotted spider mite, <i>Tetranychus urticae</i> Koch (Acari: Tetranychidae), in Japan, based on mitochondrial cytochrome oxidase subunit I sequences.. <i>Applied Entomology and Zoology</i> , 2001, 36, 459-464.	1.2	50
87	Occurrence and hybridization of two parasitoid wasps, <i>Torymus sinensis</i> Kamijo and <i>T. beneficus</i> Yasumatsu et Kamijo (Hymenoptera: Torymidae) in the Oki islands.. <i>Applied Entomology and Zoology</i> , 2000, 35, 151-154.	1.2	16
88	Molecular cloning and characterization of a microsatellite locus found in an RAPD marker of a spider mite, <i>Panonychus citri</i> (Acari: Tetranychidae). <i>Experimental and Applied Acarology</i> , 2000, 24, 385-395.	1.6	15
89	Interspecific diversity of mitochondrial COI sequences in Japanese <i>Panonychus</i> species (Acari: Tetranychidae). <i>Journal of Applied Entomology and Zoology</i> , 2000, 35, 413-420.	1.6	42
90	Laboratory experiments on a change in genetic structure with an increase of population density in the citrus red mite population, <i>Panonychus citri</i> (McGregor) (Acari : Tetranychidae). <i>Applied Entomology and Zoology</i> , 1999, 34, 413-420.	1.2	6

#	ARTICLE	IF	CITATIONS
91	Differences in ecological and genetic traits between <i>Panonychus citri</i> (Acari: Tetranychidae) populations infesting citrus and <i>Osmant hus</i> . <i>Experimental and Applied Acarology</i> , 1997, 21, 365-378.	1.6	6
92	Use of Malic Enzyme to Detect Hybrids between <i>Torymus sinensis</i> and <i>T. beneficus</i> (Hymenoptera:) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50 Hybridization.. <i>Japanese Journal of Applied Entomology and Zoology</i> , 1996, 40, 205-208.	0.1	16
93	Differences in esterase isozymes between <i>Panonychus citri</i> populations infesting citrus and <i>Osmanthus</i> . <i>Experimental and Applied Acarology</i> , 1996, 20, 113-119.	1.6	19
94	Genetic distinctness between two forms of <i>Tetranychus urticae</i> Koch (Acari: Tetranychidae) detected by electrophoresis. <i>Experimental and Applied Acarology</i> , 1996, 20, 683-693.	1.6	18
95	Host Range Segregation and Reproductive Incompatibility among <i>Panonychus citri</i> Populations Infesting <i>Osmanthus</i> Trees and Other Host Plants. <i>Applied Entomology and Zoology</i> , 1996, 31, 397-406.	1.2	14
96	Relation between Banding Patterns of Malic Enzyme by Electrophoresis and a Morphological Character in Exotic and Native <i>Torymus</i> Species. <i>Applied Entomology and Zoology</i> , 1995, 30, 37-41.	1.2	8
97	RFLP analysis of ribosomal DNA in sibling species of spider mite, genus <i>Panonychus</i> (Acari:) Tj ETQq1 1 0.784314 rgBT /Overlock 10 Tf 50	2.0	17
98	Divergence of the Northern and Southwestern Populations of <i>Panonychus mori</i> YOKOYAMA(Acari:Tetranychidae)in Japan in Host Range and Reproductive Compatibility. <i>Applied Entomology and Zoology</i> , 1993, 28, 189-197.	1.2	18
99	A Method for Preparing Permanent Specimens of Mites with Canada Balsam. <i>Applied Entomology and Zoology</i> , 1993, 28, 593-597.	1.2	12
100	Protein Differences Detected by Two-Dimensional Electrophoresis among Local Populations of <i>Panonychus citri</i> (MCGREGOR)(Acari:Tetranychidae)in Japan. <i>Applied Entomology and Zoology</i> , 1993, 28, 497-502.	1.2	10
101	A New Fixation Method for Preparing Mite Specimens for Optical and SEM Microscopic Observations. <i>Applied Entomology and Zoology</i> , 1992, 27, 427-436.	1.2	26
102	Predator fauna on windbreaks around citrus groves.. <i>Japanese Journal of Applied Entomology and Zoology</i> , 1991, 35, 49-56.	0.1	16
103	Genetic Analysis of Esterases in the Citrus Red Mite, <i>Panonychus citri</i> (MCGREGOR)(Acari:Tetranychidae). <i>Applied Entomology and Zoology</i> , 1991, 26, 307-312.	1.2	12
104	Development of the European Red Mite, <i>Panonychus ulmi</i> (KOCH)(Acari : Tetranychidae), on Mulberry and Some Other Plants. <i>Applied Entomology and Zoology</i> , 1990, 25, 326-328.	1.2	4
105	Relationships between Food Substances and Developmental Success in <i>Amblyseius sojaensis</i> EHARA (Acarina : Phytoseiidae). <i>Applied Entomology and Zoology</i> , 1988, 23, 45-51.	1.2	33
106	Identification of pesticide resistant phytoseiid mite populations in citrus orchards, and on grapevines in glasshouses and vinyl-houses (Acarina:Phytoseiidae).. <i>Japanese Journal of Applied Entomology and Zoology</i> , 1987, 31, 398-403.	0.1	9
107	Hibernation of the Kanzawa spider mite, <i>Tetranychus kanzawai</i> Kishida (Acarina: Tetranychidae) and its predators in grapevine glasshouses.. <i>Japanese Journal of Applied Entomology and Zoology</i> , 1987, 31, 23-27.	0.1	2
108	Effect of <i>Amblyseius sojaensis</i> EHARA (Acarina : Phytoseiidae) as a Predator of <i>Panonychus citri</i> (McGREGOR) and <i>Tetranychus kanzawai</i> KISHIDA : Acarina : Tetranychidae. <i>Applied Entomology and Zoology</i> , 1987, 22, 594-599.	1.2	30

#	ARTICLE	IF	CITATIONS
109	Esterase Activities and Developmental Success of the Citrus Red Mite, <i>Panonychus citri</i> (MCGREGOR) (Acarina : Tetranychidae), on Several Plants. <i>Applied Entomology and Zoology</i> , 1987, 22, 35-44.	1.2	21
110	Difference of Esterase Isozymes between Non-diapausing and Diapausing Strains of the Citrus Red Mite, <i>Panonychus citri</i> (MCGREGOR) : Acarina : Tetranychidae. <i>Applied Entomology and Zoology</i> , 1987, 22, 577-584.	1.2	17
111	Reproductive rates in the citrus red mite, <i>Panonychus citri</i> (McGregor), under high temperature conditions with daily fluctuations.. <i>Japanese Journal of Applied Entomology and Zoology</i> , 1986, 30, 21-26.	0.1	1
112	Feeding, Reproduction and Development of <i>Amblyseius sojaensis</i> EHARA (Acarina : Phytoseiidae) on Two Species of Spider Mites and on Tea Pollen. <i>Applied Entomology and Zoology</i> , 1986, 21, 322-327.	1.2	21
113	Properties of non-specific esterases of the citrus red mite, <i>Panonychus citri</i> (McGregor), which hydrolyze .ALPHA.- and .BETA.-naphthyl acetate.. <i>Japanese Journal of Applied Entomology and Zoology</i> , 1985, 29, 50-54.	0.1	4
114	Esterase zymogram of the citrus red mite, <i>Panonychus citri</i> (McGregor), on citrus and pear. <i>Japanese Journal of Applied Entomology and Zoology</i> , 1984, 28, 1-4.	0.1	13
115	ã,³ã,-ãfãfã,-éjžãç™è,²ã,¹ãf†ãf¹/4ã,ã·ã¹/4ã,é†ã·ãé-Çãj,. <i>Japanese Journal of Applied Entomology and Zoology</i> , 1982, 26, 294-299		