

Jung Ko

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3476224/publications.pdf>

Version: 2024-02-01

8
papers

61
citations

1684188
5
h-index

1588992
8
g-index

8
all docs

8
docs citations

8
times ranked

68
citing authors

#	ARTICLE	IF	CITATIONS
1	Diet containing dehulled adlay ameliorates hepatic steatosis, inflammation and insulin resistance in rats with non-alcoholic fatty liver disease. <i>British Journal of Nutrition</i> , 2022, 128, 369-376.	2.3	5
2	Consumption of Dehulled Adlay Improved Lipid Metabolism and Inflammation in Overweight and Obese Individuals after a 6-Week Single-Arm Pilot Study. <i>Nutrients</i> , 2022, 14, 2250.	4.1	5
3	Dehulled Adlay Consumption Modulates Blood Pressure in Spontaneously Hypertensive Rats and Overweight and Obese Young Adults. <i>Nutrients</i> , 2021, 13, 2305.	4.1	5
4	Crude extract of <i>Camellia oleifera</i> pomace ameliorates the progression of non-alcoholic fatty liver disease via decreasing fat accumulation, insulin resistance and inflammation. <i>British Journal of Nutrition</i> , 2020, 123, 508-515.	2.3	11
5	<i>Hylocereus polyrhizus</i> Peel Extract Retards Alcoholic Liver Disease Progression by Modulating Oxidative Stress and Inflammatory Responses in C57BL/6 Mice. <i>Nutrients</i> , 2020, 12, 3884.	4.1	8
6	<i>Camellia Oleifera</i> Seed Extract Mildly Ameliorates Carbon Tetrachloride-Induced Hepatotoxicity in Rats by Suppressing Inflammation. <i>Journal of Food Science</i> , 2019, 84, 1586-1591.	3.1	11
7	<i>Camellia oleifera</i> seed extract attenuated abdominal and hepatic fat accumulation in rats fed a high-fat diet. <i>Applied Physiology, Nutrition and Metabolism</i> , 2019, 44, 320-325.	1.9	7
8	Renoprotective Effects of Antroquinonol in Rats with Nitro-L-Arginine Methyl Ester-Induced Hypertension. <i>Nutrients</i> , 2018, 10, 1521.	4.1	9