

Sandra P Crispim

List of Publications by Year in Descending Order

Source: <https://exaly.com/author-pdf/3476555/sandra-p-crispim-publications-by-year.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

33
papers

753
citations

13
h-index

27
g-index

42
ext. papers

930
ext. citations

3.9
avg, IF

3.31
L-index

#	Paper	IF	Citations
33	Vitamin A deficiency and associated risk factors in children aged 12-59 months living in poorest municipalities in the South Region of Brazil.. <i>Public Health Nutrition</i> , 2022 , 1-26	3.3	
32	Development of a mobile application to assess Brazilian schoolchildren's diet: CADE - food consumption at home and at school.. <i>Journal of Nutritional Science</i> , 2022 , 11, e27	2.7	
31	Individual and contextual predictors of children's hemoglobin levels from Southern Brazilian municipalities in social vulnerability. <i>Cadernos De Saude Publica</i> , 2021 , 36, e00166619	3.2	1
30	Comparing Methods from the National Cancer Institute vs Multiple Source Method for Estimating Usual Intake of Nutrients in the Hispanic Community Health Study/Study of Latino Youth. <i>Journal of the Academy of Nutrition and Dietetics</i> , 2021 , 121, 59-73.e16	3.9	1
29	Methodological aspects of the assessment of dietary intake in the Brazilian National Survey on Child Nutrition (ENANI-2019): a population-based household survey. <i>Cadernos De Saude Publica</i> , 2021 , 37, e00301420	3.2	1
28	Health, lifestyle and sociodemographic characteristics are associated with Brazilian dietary patterns: Brazilian National Health Survey. <i>PLoS ONE</i> , 2021 , 16, e0247078	3.7	3
27	Assessment of bias and associated factors for food portion quantification with photos in Brazil. <i>Measurement Food</i> , 2021 , 100007		
26	Perspectives from individuals with low education and interviewers using the GloboDiet 24 h recall: a qualitative study. <i>Journal of Nutritional Science</i> , 2020 , 9, e13	2.7	2
25	Method for the Development of WISH, a Globally Applicable Index for Healthy Diets from Sustainable Food Systems. <i>Nutrients</i> , 2020 , 13,	6.7	2
24	Subjects' Perception in Quantifying Printed and Digital Photos of Food Portions. <i>Nutrients</i> , 2019 , 11,	6.7	6
23	Excess Body Weight, Snack Limits and Dental Caries in Brazilian Preschoolers: A Population-Based Study. <i>Pesquisa Brasileira Em Odontopediatria E Clinica Integrada</i> , 2019 , 19, 1-9	0.7	4
22	Comparison of meal patterns across five European countries using standardized 24-h recall (GloboDiet) data from the EFCOVAL project. <i>European Journal of Nutrition</i> , 2018 , 57, 1045-1057	5.2	14
21	Identification of Requirements for Computer-Supported Matching of Food Consumption Data with Food Composition Data. <i>Nutrients</i> , 2018 , 10,	6.7	7
20	Adapting the standardised computer- and interview-based 24 h dietary recall method (GloboDiet) for dietary monitoring in Latin America. <i>Public Health Nutrition</i> , 2017 , 20, 2847-2858	3.3	13
19	Comparison of different approaches to calculate nutrient intakes based upon 24-h recall data derived from a multicenter study in European adolescents. <i>European Journal of Nutrition</i> , 2016 , 55, 537-545	5.2	25
18	Association of Overweight with Food Portion Size among Adults of So Paulo - Brazil. <i>PLoS ONE</i> , 2016 , 11, e0164127	3.7	6
17	Comparison of the ISU, NCI, MSM, and SPADE Methods for Estimating Usual Intake: A Simulation Study of Nutrients Consumed Daily. <i>Nutrients</i> , 2016 , 8, 166	6.7	29

16	Reporting accuracy of population dietary sodium intake using duplicate 24 h dietary recalls and a salt questionnaire. <i>British Journal of Nutrition</i> , 2015 , 113, 488-97	3.6	21
15	Comparison of two food record-based dietary assessment methods for a pan-European food consumption survey among infants, toddlers, and children using data quality indicators. <i>European Journal of Nutrition</i> , 2015 , 54, 437-45	5.2	13
14	VI Latin American Workshop on Leadership in Nutrition, Cuba 2012: the first harvest. <i>Nutrition</i> , 2014 , 30, 369-70	4.8	
13	Quality assurance of the international computerised 24 h dietary recall method (EPIC-Soft). <i>British Journal of Nutrition</i> , 2014 , 111, 506-15	3.6	25
12	Evaluation of food and nutrient intake assessment using concentration biomarkers in European adolescents from the Healthy Lifestyle in Europe by Nutrition in Adolescence study. <i>British Journal of Nutrition</i> , 2013 , 109, 736-47	3.6	27
11	Bias in protein and potassium intake collected with 24-h recalls (EPIC-Soft) is rather comparable across European populations. <i>European Journal of Nutrition</i> , 2012 , 51, 997-1010	5.2	20
10	Review and evaluation of innovative technologies for measuring diet in nutritional epidemiology. <i>International Journal of Epidemiology</i> , 2012 , 41, 1187-203	7.8	254
9	Predicting urinary creatinine excretion and its usefulness to identify incomplete 24 h urine collections. <i>British Journal of Nutrition</i> , 2012 , 108, 1118-25	3.6	24
8	Design aspects of 24 h recall assessments may affect the estimates of protein and potassium intake in dietary surveys. <i>Public Health Nutrition</i> , 2012 , 15, 1196-200	3.3	6
7	Biomarker-based evaluation of two 24-h recalls for comparing usual fish, fruit and vegetable intakes across European centers in the EFCOVAL Study. <i>European Journal of Clinical Nutrition</i> , 2011 , 65 Suppl 1, S38-47	5.2	40
6	Two non-consecutive 24 h recalls using EPIC-Soft software are sufficiently valid for comparing protein and potassium intake between five European centres--results from the European Food Consumption Validation (EFCOVAL) study. <i>British Journal of Nutrition</i> , 2011 , 105, 447-58	3.6	63
5	Inventory of experiences from national/regional dietary monitoring surveys using EPIC-Soft. <i>European Journal of Clinical Nutrition</i> , 2011 , 65 Suppl 1, S16-28	5.2	13
4	Respondents' evaluation of the 24-h dietary recall method (EPIC-Soft) in the EFCOVAL Project. <i>European Journal of Clinical Nutrition</i> , 2011 , 65 Suppl 1, S29-37	5.2	15
3	Dietary exposure to flavouring substances: from screening methods to detailed assessments using food consumption data collected with EPIC-Soft software. <i>Food Additives and Contaminants - Part A Chemistry, Analysis, Control, Exposure and Risk Assessment</i> , 2010 , 27, 433-46	3.2	12
2	Validade relativa de um questionário de frequência alimentar para utilização em adultos. <i>Revista De Nutricao</i> , 2009 , 22, 81-95	1.8	7
1	The influence of education in the validation process of a food frequency questionnaire for adults in Viçosa, Minas Gerais, Brazil. <i>European Journal of Clinical Nutrition</i> , 2006 , 60, 1311-6	5.2	5