
Esi Domi

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/347685/publications.pdf Version: 2024-02-01



FSI DOMI

#	Article	IF	CITATIONS
1	Genetic deletion or pharmacological blockade of nociceptin/orphanin FQ receptors in the ventral tegmental area attenuates nicotineâ€motivated behaviour. British Journal of Pharmacology, 2022, 179, 2647-2658.	2.7	5
2	A Role for Neuropeptide S in Alcohol and Cocaine Seeking. Pharmaceuticals, 2022, 15, 800.	1.7	3
3	Downregulation of Synaptotagmin 1 in the Prelimbic Cortex Drives Alcohol-Associated Behaviors in Rats. Biological Psychiatry, 2021, 89, 398-406.	0.7	14
4	Dysregulation of the histone demethylase KDM6B in alcohol dependence is associated with epigenetic regulation of inflammatory signaling pathways. Addiction Biology, 2021, 26, e12816.	1.4	28
5	Targeting the Opioid Receptors: A Promising Therapeutic Avenue for Treatment in "Heavy Drinking Smokers― Alcohol and Alcoholism, 2021, 56, 127-138.	0.9	6
6	Stressâ€induced escalation of alcohol selfâ€administration, anxietyâ€like behavior, and elevated amygdala Avp expression in a susceptible subpopulation of rats. Addiction Biology, 2021, 26, e13009.	1.4	12
7	Neurobiology of alcohol seeking behavior. Journal of Neurochemistry, 2021, 157, 1585-1614.	2.1	29
8	A neural substrate of compulsive alcohol use. Science Advances, 2021, 7, .	4.7	46
9	Further evidence for the involvement of the PPARÎ ³ system on alcohol intake and sensitivity in rodents. Psychopharmacology, 2020, 237, 2983-2992.	1.5	6
10	Nicotine increases alcohol selfâ€administration in male rats via a μâ€opioid mechanism within the mesolimbic pathway. British Journal of Pharmacology, 2020, 177, 4516-4531.	2.7	9
11	Activation of PPARÎ ³ Attenuates the Expression of Physical and Affective Nicotine Withdrawal Symptoms through Mechanisms Involving Amygdala and Hippocampus Neurotransmission. Journal of Neuroscience, 2019, 39, 9864-9875.	1.7	26
12	Sub-dimensions of Alcohol Use Disorder in Alcohol Preferring and Non-preferring Rats, a Comparative Study. Frontiers in Behavioral Neuroscience, 2019, 13, 3.	1.0	19
13	Oxytocin Reduces Alcohol Cue-Reactivity in Alcohol-Dependent Rats and Humans. Neuropsychopharmacology, 2018, 43, 1235-1246.	2.8	85
14	A molecular mechanism for choosing alcohol over an alternative reward. Science, 2018, 360, 1321-1326.	6.0	169
15	Protection against alcohol-induced neuronal and cognitive damage by the PPARÎ ³ receptor agonist pioglitazone. Brain, Behavior, and Immunity, 2017, 64, 320-329.	2.0	37
16	Genetic Deletion of Neuronal PPARÎ ³ Enhances the Emotional Response to Acute Stress and Exacerbates Anxiety: An Effect Reversed by Rescue of Amygdala PPARÎ ³ Function. Journal of Neuroscience, 2016, 36, 12611-12623.	1.7	48
17	Biological profile and bioavailability of imidazoline compounds on morphine tolerance modulation. European Journal of Pharmacology, 2015, 769, 219-224.	1.7	2
18	Investigation of allyphenyline efficacy in the treatment of alcohol withdrawal symptoms. European Journal of Pharmacology, 2015, 760, 122-128.	1.7	6

4	#	Article	IF	CITATIONS
	19	Polymorphism in the corticotropin-releasing factor receptor 1 (CRF1-R) gene plays a role in shaping the high anxious phenotype of Marchigian Sardinian alcohol-preferring (msP) rats. Psychopharmacology, 2015, 232, 1083-1093.	1.5	25
4	20	Role of a Genetic Polymorphism in the Corticotropin-Releasing Factor Receptor 1 Gene in Alcohol Drinking and Seeking Behaviors of Marchigian Sardinian Alcohol-Preferring Rats. Frontiers in Psychiatry, 2013, 4, 23.	1.3	42