
Xingchen Zhao

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/3478522/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Advanced glycation endâ€products suppress autophagic flux in podocytes by activating mammalian target of rapamycin and inhibiting nuclear translocation of transcription factor EB. Journal of Pathology, 2018, 245, 235-248.	4.5	71
2	Receptor activator of NF-κB mediates podocyte injury in diabetic nephropathy. Kidney International, 2021, 100, 377-390.	5.2	27
3	Protective effect of NSA on intestinal epithelial cells in a necroptosis model. Oncotarget, 2017, 8, 86726-86735.	1.8	27
4	RIP3 impedes transcription factor EB to suppress autophagic degradation in septic acute kidney injury. Cell Death and Disease, 2021, 12, 593.	6.3	20
5	RIPK3 mediates renal tubular epithelial cell apoptosis in endotoxin‑induced acute kidney injury. Molecular Medicine Reports, 2019, 20, 1613-1620.	2.4	17
6	Extracellular Vesicles From High Glucose-Treated Podocytes Induce Apoptosis of Proximal Tubular Epithelial Cells. Frontiers in Physiology, 2020, 11, 579296.	2.8	13
7	Lipopolysaccharide-induced podocyte injury is mediated by suppression of autophagy. Molecular Medicine Reports, 2016, 14, 811-818.	2.4	11
8	Peripheral kynurenine/tryptophan ratio is not a reliable marker of systemic indoleamine 2,3-dioxygenase: A lesson drawn from patients on hemodialysis. Oncotarget, 2017, 8, 25261-25269.	1.8	11
9	Septin 7 mediates high glucose-induced podocyte apoptosis. Biochemical and Biophysical Research Communications, 2018, 506, 522-528.	2.1	9
10	WWC1 promotes podocyte survival via stabilizing slit diaphragm protein dendrin. Molecular Medicine Reports, 2017, 16, 8685-8690.	2.4	8
11	Amino acid starvation promotes podocyte autophagy through mammalian target of rapamycin inhibition and transcription factor EB activation. Molecular Medicine Reports, 2018, 18, 4342-4348.	2.4	6