
Joanne S Haviland

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/348580/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Can patient decision aids reduce decisional conflict in a de-escalation of breast radiotherapy clinical trial? The PRIMETIME Study Within a Trial implemented using a cluster stepped-wedge trial design. Trials, 2021, 22, 397.	0.7	4
2	When the World Throws You a Curve Ball: Lessons Learned in Breast Cancer Management. American Society of Clinical Oncology Educational Book / ASCO American Society of Clinical Oncology Meeting, 2021, 41, e79-e89.	1.8	3
3	Impact of Hypofractionated Radiotherapy on Patient-reported Outcomes in Prostate Cancer: Results up to 5Âyr in the CHHiP trial (CRUK/06/016). European Urology Oncology, 2021, 4, 980-992.	2.6	14
4	Ten-Year Results of FAST: A Randomized Controlled Trial of 5-Fraction Whole-Breast Radiotherapy for Early Breast Cancer. Journal of Clinical Oncology, 2020, 38, 3261-3272.	0.8	175
5	Targeted Intraoperative Radiotherapy for Early Breast Cancer. JAMA Oncology, 2020, 6, 1636.	3.4	8
6	Internal mammary node irradiation in breast cancer: does benefit outweigh risk?. Lancet Oncology, The, 2020, 21, 1541-1543.	5.1	2
7	Breast cancer radiation therapy – Authors' reply. Lancet, The, 2020, 396, 1559-1560.	6.3	0
8	Hypofractionated breast radiotherapy for 1 week versus 3 weeks (FAST-Forward): 5-year efficacy and late normal tissue effects results from a multicentre, non-inferiority, randomised, phase 3 trial. Lancet, The, 2020, 395, 1613-1626.	6.3	603
9	Abstract P6-12-04: Does the addition of a patient decision aid video reduce decisional conflict in patients considering a de-escalation of radiotherapy trial? Results from the PRIMETIME study. , 2020, , .		2
10	Abstract P6-12-10: Method development for a â€~Study within a Trial' using a cluster stepped wedge trial design: Evaluating a patient decision aid video on patients' decisional conflict in the PRIMETIME study. , 2020, , .		0
11	Smoking, alcohol consumption, diet and physical activity following stoma formation surgery, stoma-related concerns, and desire for lifestyle advice: a United Kingdom survey. BMC Public Health, 2019, 19, 574.	1.2	15
12	Is breast seroma after tumour resection associated with patient-reported breast appearance change following radiotherapy? Results from the IMPORT HIGH (CRUK/06/003) trial. Radiotherapy and Oncology, 2019, 136, 190-196.	0.3	8
13	Can patient-reported outcomes be used instead of clinician-reported outcomes and photographs as primary endpoints of late normal tissue effects in breast radiotherapy trials? Results from the IMPORT LOW trial. Radiotherapy and Oncology, 2019, 134, 220-230.	0.3	17
14	Patient-Reported Outcomes Over 5 Years After Whole- or Partial-Breast Radiotherapy: Longitudinal Analysis of the IMPORT LOW (CRUK/06/003) Phase III Randomized Controlled Trial. Journal of Clinical Oncology, 2019, 37, 305-317.	0.8	58
15	Women's Free-text Comments on their Quality of Life: An Exploratory Analysis from the UK Standardisation of Breast Radiotherapy (START) Trials for Early Breast Cancer. Clinical Oncology, 2018, 30, 433-441.	0.6	7
16	Targeted radiotherapy for early breast cancer – Authors' reply. Lancet, The, 2018, 391, 27-28.	6.3	1
17	Ki67 Is an Independent Predictor of Recurrence in the Largest Randomized Trial of 3 Radiation Fractionation Schedules in Localized Prostate Cancer. International Journal of Radiation Oncology Biology Physics, 2018, 101, 309-315.	0.4	19
18	Late normal tissue effects in the arm and shoulder following lymphatic radiotherapy: Results from the UK START (Standardisation of Breast Radiotherapy) trials. Radiotherapy and Oncology, 2018, 126, 155-162.	0.3	72

JOANNE S HAVILAND

#	Article	IF	CITATIONS
19	FAST Phase III RCT of Radiotherapy Hypofractionation for Treatment of Early Breast Cancer: 10-Year Results (CRUKE/04/015). International Journal of Radiation Oncology Biology Physics, 2018, 102, 1603-1604.	0.4	40
20	OC-0156: Concordance of patient reported outcomes with clinician and photographic assessments in IMPORT LOW. Radiotherapy and Oncology, 2018, 127, S79.	0.3	1
21	Assessment of Ki67 in relation to radiotherapy (RT) fractionation and prognosis in the CHHiP (CRUK/06/016) trial Journal of Clinical Oncology, 2018, 36, 98-98.	0.8	0
22	SP-0314: Partial breast radiotherapy after breast conservation: 5 year outcomes from the IMPORT LOW (CRUK/06/003) phase III trial. Radiotherapy and Oncology, 2017, 123, S162-S163.	0.3	1
23	Partial-breast radiotherapy after breast conservation surgery for patients with early breast cancer (UK IMPORT LOW trial): 5-year results from a multicentre, randomised, controlled, phase 3, non-inferiority trial. Lancet, The, 2017, 390, 1048-1060.	6.3	448
24	Social support following diagnosis and treatment for colorectal cancer and associations with healthâ€related quality of life: Results from the UK ColoREctal Wellbeing (CREW) cohort study. Psycho-Oncology, 2017, 26, 2276-2284.	1.0	53
25	Colorectal cancer patient's self-efficacy for managing illness-related problems in the first 2Âyears after diagnosis, results from the ColoREctal Well-being (CREW) study. Journal of Cancer Survivorship, 2017, 11, 634-642.	1.5	34
26	Prolongation of overall treatment time as a cause of treatment failure in early breast cancer: An analysis of the UK START (Standardisation of Breast Radiotherapy) trials of radiotherapy fractionation. Radiotherapy and Oncology, 2016, 121, 420-423.	0.3	36
27	A multicentre study of the evidence for customized margins in photon breast boost radiotherapy. British Journal of Radiology, 2016, 89, 20150603.	1.0	6
28	Does breast composition influence late adverse effects in breast radiotherapy?. Breast, 2016, 26, 25-30.	0.9	2
29	Pre-Surgery Depression and Confidence to Manage Problems Predict Recovery Trajectories of Health and Wellbeing in the First Two Years following Colorectal Cancer: Results from the CREW Cohort Study. PLoS ONE, 2016, 11, e0155434.	1.1	80
30	The UK HeartSpare Study (Stage IB): Randomised comparison of a voluntary breath-hold technique and prone radiotherapy after breast conserving surgery. Radiotherapy and Oncology, 2015, 114, 66-72.	0.3	72
31	In Regard to Vaidya etÂal. International Journal of Radiation Oncology Biology Physics, 2015, 92, 954-955.	0.4	4
32	Late effects of breast radiotherapy. Lancet, The, 2014, 383, 1383-1384.	6.3	1
33	Haste makes waste: Are the data regarding TARGIT-A IORT ready for prime time?. Breast Cancer Research and Treatment, 2014, 147, 221-222.	1.1	2
34	Cytokine levels as biomarkers of radiation fibrosis in patients treated with breast radiotherapy. Radiation Oncology, 2014, 9, 103.	1.2	11
35	Radiotherapy for breast cancer, the TARGIT-A trial. Lancet, The, 2014, 383, 1716-1717.	6.3	32
36	A genome wide association study (GWAS) providing evidence of an association between common genetic variants and late radiotherapy toxicity. Radiotherapy and Oncology, 2014, 111, 178-185.	0.3	128

JOANNE S HAVILAND

#	Article	IF	CITATIONS
37	Voluntary Breath-hold Technique for Reducing Heart Dose in Left Breast Radiotherapy. Journal of Visualized Experiments, 2014, , .	0.2	27
38	The UK Standardisation of Breast Radiotherapy (START) trials of radiotherapy hypofractionation for treatment of early breast cancer: 10-year follow-up results of two randomised controlled trials. Lancet Oncology, The, 2013, 14, 1086-1094.	5.1	1,141
39	The UK HeartSpare Study: Randomised evaluation of voluntary deep-inspiratory breath-hold in women undergoing breast radiotherapy. Radiotherapy and Oncology, 2013, 108, 242-247.	0.3	163
40	Conducting radiogenomic research – Do not forget careful consideration of the clinical data. Radiotherapy and Oncology, 2012, 105, 337-340.	0.3	28
41	The impact of dose heterogeneity on late normal tissue complication risk after hypofractionated whole breast radiotherapy. Radiotherapy and Oncology, 2012, 104, 143-147.	0.3	19
42	Quality of Life and Experience of Care in Women With Metastatic Breast Cancer: A Cross-Sectional Survey. Journal of Pain and Symptom Management, 2012, 43, 747-758.	0.6	75
43	First results of the randomised UK FAST Trial of radiotherapy hypofractionation for treatment of early breast cancer (CRUKE/04/015). Radiotherapy and Oncology, 2011, 100, 93-100.	0.3	226
44	Corrigendum to "Randomised phase II trial of hyperbaric oxygen therapy in patients with chronic arm lymphoedema after radiotherapy for cancer―[Radiother Oncol 97 (2010) 101–107]. Radiotherapy and Oncology, 2011, 98, 285.	0.3	0
45	Large breast size as a risk factor for late adverse effects of breast radiotherapy: Is residual dose inhomogeneity, despite 3D treatment planning and delivery, the main explanation?. Radiotherapy and Oncology, 2011, 100, 236-240.	0.3	66
46	Ipsilateral Breast Tumor Relapse: Local Recurrence Versus New Primary Tumor and the Effect of Whole-Breast Radiotherapy on the Rate of New Primaries. International Journal of Radiation Oncology Biology Physics, 2011, 79, 19-25.	0.4	49
47	Hypofractionated Whole-Breast Radiotherapy for Women With Early Breast Cancer: Myths and Realities. International Journal of Radiation Oncology Biology Physics, 2011, 79, 1-9.	0.4	142
48	The course of anxiety and depression over 5 years of follow-up and risk factors in women with early breast cancer: Results from the UK Standardisation of Radiotherapy Trials (START). Breast, 2010, 19, 84-91.	0.9	86
49	Pushing the limits of hypofractionation for adjuvant whole breast radiotherapy. Breast, 2010, 19, 176-179.	0.9	15
50	Test of association between variant tgî²1 alleles and late adverse effects of breast radiotherapy. Radiotherapy and Oncology, 2010, 97, 15-18.	0.3	19
51	Randomised phase II trial of hyperbaric oxygen therapy in patients with chronic arm lymphoedema after radiotherapy for cancer. Radiotherapy and Oncology, 2010, 97, 101-107.	0.3	42
52	Comparison of patient-reported breast, arm, and shoulder symptoms and body image after radiotherapy for early breast cancer: 5-year follow-up in the randomised Standardisation of Breast Radiotherapy (START) trials. Lancet Oncology, The, 2010, 11, 231-240.	5.1	201
53	Quality of life after breast radiotherapy – Authors' reply. Lancet Oncology, The, 2010, 11, 612-613.	5.1	5
54	Hypofractionated Radiotherapy for Breast Cancer. New England Journal of Medicine, 2010, 362, 1843-1844.	13.9	34

#	Article	IF	CITATIONS
55	Hypofractionated Adjuvant Whole Breast Radiotherapy: Progress and Prospects. Acta Oncológica, 2010, 49, 1288-1292.	0.8	11
56	Corrigendum to "The impact of age and clinical factors on quality of life in early breast cancer: An analysis of 2208 women recruited to the UK START Trial (Standardisation of Breast Radiotherapy) Tj ETQq0 0 0 rg	gB ō,∕ Øverl	oc b 10 Tf 50
57	A Randomized Controlled Trial of Relaxation Training to Reduce Hot Flashes in Women with Primary Breast Cancer. Journal of Pain and Symptom Management, 2008, 35, 397-405.	0.6	51
58	The impact of age and clinical factors on quality of life in early breast cancer: An analysis of 2208 women recruited to the UK START Trial (Standardisation of Breast Radiotherapy Trial). Breast, 2007, 16, 241-251.	0.9	172
59	Effect of radiotherapy fraction size on tumour control in patients with early-stage breast cancer after local tumour excision: long-term results of a randomised trial. Lancet Oncology, The, 2006, 7, 467-471.	5.1	520

60	Dose and fractionation regimens for breast cancer – Authors' reply. Lancet Oncology, The, 2006, 7, 619-620.	5.1	0
----	--	-----	---