

Antonella Meloni

List of Publications by Citations

Source: <https://exaly.com/author-pdf/3489387/antonella-meloni-publications-by-citations.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

59
papers

1,494
citations

21
h-index

38
g-index

69
ext. papers

1,821
ext. citations

5
avg, IF

3.65
L-index

#	Paper	IF	Citations
59	A common mutation in Sardinian autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophy patients. <i>Human Genetics</i> , 1998 , 103, 428-34	6.3	192
58	Autoimmune polyendocrine syndrome type 1: an extensive longitudinal study in Sardinian patients. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2012 , 97, 1114-24	5.6	120
57	Cardiac and hepatic iron and ejection fraction in thalassemia major: multicentre prospective comparison of combined deferiprone and deferoxamine therapy against deferiprone or deferoxamine monotherapy. <i>Journal of Cardiovascular Magnetic Resonance</i> , 2013 , 15, 1	6.9	111
56	Deferasirox, deferiprone and desferrioxamine treatment in thalassemia major patients: cardiac iron and function comparison determined by quantitative magnetic resonance imaging. <i>Haematologica</i> , 2011 , 96, 41-7	6.6	105
55	Cardiac iron and cardiac disease in males and females with transfusion-dependent thalassemia major: a T2* magnetic resonance imaging study. <i>Haematologica</i> , 2011 , 96, 515-20	6.6	80
54	Cytochrome P450 1A2 is a hepatic autoantigen in autoimmune polyglandular syndrome type 1. <i>Journal of Clinical Endocrinology and Metabolism</i> , 1997 , 82, 1353-61	5.6	69
53	Multiparametric Cardiac Magnetic Resonance Survey in Children With Thalassemia Major: A Multicenter Study. <i>Circulation: Cardiovascular Imaging</i> , 2015 , 8, e003230	3.9	51
52	The use of appropriate calibration curves corrects for systematic differences in liver R2* values measured using different software packages. <i>British Journal of Haematology</i> , 2013 , 161, 888-91	4.5	49
51	Prevalence of autoimmune thyroiditis in children with celiac disease and effect of gluten withdrawal. <i>Journal of Pediatrics</i> , 2009 , 155, 51-5, 55.e1	3.6	48
50	Single region of interest versus multislice T2* MRI approach for the quantification of hepatic iron overload. <i>Journal of Magnetic Resonance Imaging</i> , 2011 , 33, 348-55	5.6	45
49	Prediction of cardiac complications for thalassemia major in the widespread cardiac magnetic resonance era: a prospective multicentre study by a multi-parametric approach. <i>European Heart Journal Cardiovascular Imaging</i> , 2018 , 19, 299-309	4.1	43
48	Myocardial iron overload in thalassaemia major. How early to check?. <i>British Journal of Haematology</i> , 2014 , 164, 579-85	4.5	39
47	Multislice multiecho T2* cardiac magnetic resonance for the detection of heterogeneous myocardial iron distribution in thalassaemia patients. <i>NMR in Biomedicine</i> , 2009 , 22, 707-15	4.4	39
46	Defective autoimmune regulator-dependent central tolerance to myelin protein zero is linked to autoimmune peripheral neuropathy. <i>Journal of Immunology</i> , 2012 , 188, 4906-12	5.3	39
45	Preferential patterns of myocardial iron overload by multislice multiecho T*2 CMR in thalassemia major patients. <i>Magnetic Resonance in Medicine</i> , 2010 , 64, 211-9	4.4	39
44	Cardiac complications and diabetes in thalassaemia major: a large historical multicentre study. <i>British Journal of Haematology</i> , 2013 , 163, 520-7	4.5	38
43	Different patterns of myocardial iron distribution by whole-heart T2* magnetic resonance as risk markers for heart complications in thalassemia major. <i>International Journal of Cardiology</i> , 2014 , 177, 1012-9	3.2	29

42	Influence of myocardial fibrosis and blood oxygenation on heart T2* values in thalassemia patients. <i>Journal of Magnetic Resonance Imaging</i> , 2009 , 29, 832-7	5.6	24
41	Chronic inflammatory demyelinating polyneuropathy as a possible novel component of autoimmune poly-endocrine-candidiasis-ectodermal dystrophy. <i>European Journal of Pediatrics</i> , 2009 , 168, 237-40	4.1	23
40	Improvement of heart iron with preserved patterns of iron store by CMR-guided chelation therapy. <i>European Heart Journal Cardiovascular Imaging</i> , 2015 , 16, 325-34	4.1	21
39	Regional and global pancreatic T*2 MRI for iron overload assessment in a large cohort of healthy subjects: normal values and correlation with age and gender. <i>Magnetic Resonance in Medicine</i> , 2011 , 65, 764-9	4.4	21
38	Safety of cardiovascular magnetic resonance gadolinium chelates contrast agents in patients with hemoglobinopathies. <i>Haematologica</i> , 2009 , 94, 1625-7	6.6	21
37	Pancreatic iron overload by T2* MRI in a large cohort of well treated thalassemia major patients: can it tell us heart iron distribution and function?. <i>American Journal of Hematology</i> , 2015 , 90, E189-90	7.1	19
36	Extramedullary hematopoiesis is associated with lower cardiac iron loading in chronically transfused thalassemia patients. <i>American Journal of Hematology</i> , 2015 , 90, 1008-12	7.1	19
35	Myocardial fibrosis by late gadolinium enhancement cardiac magnetic resonance and hepatitis C virus infection in thalassemia major patients. <i>Journal of Cardiovascular Medicine</i> , 2015 , 16, 689-95	1.9	17
34	Median eminence dopaminergic nerve terminals: a novel target in autoimmune polyendocrine syndrome?. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2005 , 90, 4108-11	5.6	17
33	MRI multicentre prospective survey in thalassaemia major patients treated with deferasirox versus deferiprone and desferrioxamine. <i>British Journal of Haematology</i> , 2018 , 183, 783-795	4.5	16
32	Comparison of biventricular dimensions and function between pediatric sickle-cell disease and thalassemia major patients without cardiac iron. <i>American Journal of Hematology</i> , 2013 , 88, 213-8	7.1	14
31	Fast generation of T2* maps in the entire range of clinical interest: application to thalassemia major patients. <i>Computers in Biology and Medicine</i> , 2015 , 56, 200-10	7	13
30	Gender differences in the development of cardiac complications: a multicentre study in a large cohort of thalassaemia major patients to optimize the timing of cardiac follow-up. <i>British Journal of Haematology</i> , 2018 , 180, 879-888	4.5	12
29	The Close Link of Pancreatic Iron With Glucose Metabolism and With Cardiac Complications in Thalassemia Major: A Large, Multicenter Observational Study. <i>Diabetes Care</i> , 2020 , 43, 2830-2839	14.6	11
28	Accurate estimate of pancreatic T2* values: how to deal with fat infiltration. <i>Abdominal Imaging</i> , 2015 , 40, 3129-36		10
27	Multicenter validation of the magnetic resonance T2* technique for quantification of pancreatic iron. <i>European Radiology</i> , 2019 , 29, 2246-2252	8	10
26	Cost-Utility Analysis of Three Iron Chelators Used in Monotherapy for the Treatment of Chronic Iron Overload in Thalassaemia Major Patients: An Italian Perspective. <i>Clinical Drug Investigation</i> , 2017 , 37, 453-464	3.2	9
25	CMR for myocardial iron overload quantification: calibration curve from the MIOT Network. <i>European Radiology</i> , 2020 , 30, 3217-3225	8	9

24	The impact of liver steatosis on the ability of serum ferritin levels to be predictive of liver iron concentration in non-transfusion-dependent thalassaemia patients. <i>British Journal of Haematology</i> , 2018 , 180, 721-726	4.5	9
23	Soluble form of transferrin receptor-1 level is associated with the age at first diagnosis and the risk of therapeutic intervention and iron overloading in patients with non-transfusion-dependent thalassemia. <i>Annals of Hematology</i> , 2017 , 96, 1541-1546	3	9
22	Novel neuronal and endocrine autoantibody targets in Autoimmune Polyendocrine Syndrome type 1. <i>Autoimmunity</i> , 2012 , 45, 485-94	3	7
21	Cardiac involvement by CMR in different genotypic groups of thalassemia major patients. <i>Blood Cells, Molecules, and Diseases</i> , 2019 , 77, 1-7	2.1	5
20	Longitudinal follow-up of patients with thalassaemia intermedia who started transfusion therapy in adulthood: a cohort study. <i>British Journal of Haematology</i> , 2020 , 191, 107-114	4.5	5
19	Biventricular Reference Values by Body Surface Area, Age, and Gender in a Large Cohort of Well-Treated Thalassemia Major Patients Without Heart Damage Using a Multiparametric CMR Approach. <i>Journal of Magnetic Resonance Imaging</i> , 2021 , 53, 61-70	5.6	5
18	Prospective CMR Survey in Children With Thalassemia Major: Insights From a National Network. <i>JACC: Cardiovascular Imaging</i> , 2020 , 13, 1284-1286	8.4	3
17	Survival and causes of death in 2,033 patients with non-transfusion-dependent β -thalassaemia. <i>Haematologica</i> , 2021 , 106, 2489-2492	6.6	3
16	Myocardial iron overload by cardiovascular magnetic resonance native segmental T1 mapping: a sensitive approach that correlates with cardiac complications. <i>Journal of Cardiovascular Magnetic Resonance</i> , 2021 , 23, 70	6.9	3
15	National networking in rare diseases and reduction of cardiac burden in thalassemia major.. <i>European Heart Journal</i> , 2021 ,	9.5	3
14	Myocardial T1 Values at 1.5 T: Normal Values for General Electric Scanners and Sex-Related Differences. <i>Journal of Magnetic Resonance Imaging</i> , 2021 , 54, 1486-1500	5.6	2
13	IGENDA protocol: gender differences in awareness, knowledge and perception of cardiovascular risk: an Italian multicenter study. <i>Journal of Cardiovascular Medicine</i> , 2019 , 20, 278-283	1.9	2
12	A complication risk score to evaluate clinical severity of thalassaemia syndromes. <i>British Journal of Haematology</i> , 2021 , 192, 626-633	4.5	2
11	Red blood cell consumption in a large cohort of patients with thalassaemia: a retrospective analysis of main predictors. <i>Annals of Hematology</i> , 2020 , 99, 1209-1215	3	1
10	Absence of T1 Hyperintensity in the Brain of High-risk Patients After Multiple Administrations of High-dose Gadobutrol for Cardiac Magnetic Resonance. <i>Clinical Neuroradiology</i> , 2021 , 31, 347-355	2.7	1
9	Myocardial T2 values at 1.5 T by a segmental approach with healthy aging and gender.. <i>European Radiology</i> , 2022 , 1	8	1
8	Genotypic groups as risk factors for cardiac magnetic resonance abnormalities and complications in thalassemia major: a large, multicentre study. <i>Blood Transfusion</i> , 2021 , 19, 168-176	3.6	1
7	The Link of Pancreatic Iron with Glucose Metabolism and Cardiac Iron in Thalassemia Intermedia: A Large, Multicenter Observational Study. <i>Journal of Clinical Medicine</i> , 2021 , 10,	5.1	1

6	Reduction of Total Brain and Cerebellum Volumes Associated With Neuronal Autoantibodies in Patients With APECED. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2019 , 104, 150-162	5.6	1
5	The effect of desferrioxamine chelation versus no therapy in patients with non transfusion-dependent thalassaemia: a multicenter prospective comparison from the MIOT network. <i>Annals of Hematology</i> , 2018 , 97, 1925-1932	3	1
4	Random Forest Clustering Identifies Three Subgroups of β -Thalassemia with Distinct Clinical Severity. <i>Thalassemia Reports</i> , 2022 , 12, 14-23	2	1
3	Risk of mortality from anemia and iron overload in nontransfusion-dependent β -thalassemia. <i>American Journal of Hematology</i> , 2021 ,	7.1	1
2	The use of hydroxyurea in the real life of MIOT network: an observational study.. <i>Expert Opinion on Drug Safety</i> , 2022 , 1-8	4.1	0
1	Pancreatic iron quantification with MR imaging: a practical guide. <i>Abdominal Radiology</i> ,	3	0