
## Rashna D Balsara

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/3492541/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Plasminogen activator inhibitor-1: the double-edged sword in apoptosis. Thrombosis and Haemostasis, 2008, 100, 1029-36.	3.4	39
2	Complement-mediated Opsonization of Invasive Group A Streptococcus pyogenes Strain AP53 Is Regulated by the Bacterial Two-component Cluster of Virulence Responder/Sensor (CovRS) System. Journal of Biological Chemistry, 2013, 288, 27494-27504.	3.4	24
3	Pharmacology of triheteromeric N-Methyl-d-Aspartate Receptors. Neuroscience Letters, 2016, 617, 240-246.	2.1	24
4	Rational design of syn-safencin, a novel linear antimicrobial peptide derived from the circular bacteriocin safencin AS-48. Journal of Antibiotics, 2018, 71, 592-600.	2.0	20
5	Conantokin-G Attenuates Detrimental Effects of NMDAR Hyperactivity in an Ischemic Rat Model of Stroke. PLoS ONE, 2015, 10, e0122840.	2.5	18
6	Synthetic Antimicrobial Peptide Tuning Permits Membrane Disruption and Interpeptide Synergy. ACS Pharmacology and Translational Science, 2020, 3, 418-424.	4.9	18
7	Contributions of different modules of the plasminogen-binding Streptococcus pyogenes M-protein that mediate its functional dimerization. Journal of Structural Biology, 2018, 204, 151-164.	2.8	14
8	A deficiency of the GluN2C subunit of the N-methyl-D-aspartate receptor is neuroprotective in a mouse model of ischemic stroke. Biochemical and Biophysical Research Communications, 2018, 495, 136-144.	2.1	13
9	In situ metabolite and lipid analysis of GluN2Dâ^'/â^' and wild-type mice after ischemic stroke using MALDI MSI. Analytical and Bioanalytical Chemistry, 2020, 412, 6275-6285.	3.7	11
10	Opposing action of conantokin-G on synaptically and extrasynaptically-activated NMDA receptors. Neuropharmacology, 2012, 62, 2227-2238.	4.1	10
11	Non-invasive Imaging and Analysis of Cerebral Ischemia in Living Rats Using Positron Emission Tomography with <sup>18</sup> F-FDG. Journal of Visualized Experiments, 2014, , .	0.3	9
12	Hydroxyproline-induced Helical Disruption in Conantokin Rl-B Affects Subunit-selective Antagonistic Activities toward Ion Channels of N-Methyl-d-aspartate Receptors. Journal of Biological Chemistry, 2015, 290, 18156-18172.	3.4	8
13	Antagonist Properties of Conus parius Peptides on N-Methyl-D-Aspartate Receptors and Their Effects on CREB Signaling. PLoS ONE, 2013, 8, e81405.	2.5	5
14	Discerning the Role of the Hydroxyproline Residue in the Structure of Conantokin Rl-B and Its Role in GluN2B Subunit-Selective Antagonistic Activity toward <i>N</i> -Methyl- <scp>d</scp> -Aspartate Receptors. Biochemistry, 2016, 55, 7112-7122.	2.5	3
15	Structural and Functional Properties of Nâ€Methylâ€Dâ€Aspartate Receptorâ€&pecific ConRlB. FASEB Journal, 2015, 29, 570.13.	0.5	0