

Yoshihiko Nakamura

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3495295/publications.pdf>

Version: 2024-02-01

37
papers

1,177
citations

430442

18
h-index

395343

33
g-index

38
all docs

38
docs citations

38
times ranked

1667
citing authors

#	ARTICLE	IF	CITATIONS
1	Early recognition of sepsis-induced coagulopathy using the C2PAC index: a ratio of soluble type C lectin-like receptor 2 (sCLEC-2) level and platelet count. <i>Platelets</i> , 2022, 33, 935-944.	1.1	18
2	Oral Health of COVID-19 Patients Using Extracorporeal Membrane Oxygenation: A Clinical Study of 19 Cases. <i>Journal of Clinical Medicine</i> , 2022, 11, 42.	1.0	4
3	The Association between the Oral-Gut Axis and the Outcomes of Severe COVID-19 Patients Receiving Extracorporeal Membrane Oxygenation: A Case-Control Study. <i>Journal of Clinical Medicine</i> , 2022, 11, 1167.	1.0	3
4	Effect of Whole Tissue Culture and Basic Fibroblast Growth Factor on Maintenance of Tie2 Molecule Expression in Human Nucleus Pulposus Cells. <i>International Journal of Molecular Sciences</i> , 2021, 22, 4723.	1.8	12
5	Optimization of Spheroid Colony Culture and Cryopreservation of Nucleus Pulposus Cells for the Development of Intervertebral Disc Regenerative Therapeutics. <i>Applied Sciences (Switzerland)</i> , 2021, 11, 3309.	1.3	7
6	Differences in coagulopathy and massive transfusion strategy based on trauma type. <i>American Journal of Emergency Medicine</i> , 2020, 38, 860-863.	0.7	1
7	Association of Self-Reported Medication Adherence with Potentially Inappropriate Medications in Elderly Patients: A Cross-Sectional Pilot Study. <i>International Journal of Environmental Research and Public Health</i> , 2020, 17, 5940.	1.2	4
8	Enhanced effect of recombinant human soluble thrombomodulin by ultrasound irradiation in acute liver failure. <i>Scientific Reports</i> , 2020, 10, 1742.	1.6	4
9	Inappropriate direct oral anticoagulant dosing in atrial fibrillation patients is associated with prescriptions for outpatients rather than inpatients: a single-center retrospective cohort study. <i>Journal of Pharmaceutical Health Care and Sciences</i> , 2020, 6, 2.	0.4	19
10	Antithrombin gamma attenuates macrophage/microglial activation and brain damage after transient focal cerebral ischemia in mice. <i>Life Sciences</i> , 2020, 252, 117665.	2.0	12
11	Characteristics of coagulation/fibrinolysis abnormalities in severe novel coronavirus disease 2019 (COVID-19) patients – Case series –. <i>Japanese Journal of Thrombosis and Hemostasis</i> , 2020, 31, 398-408.	0.1	2
12	Long-Term Treatment with Thrombomodulin Improves Functional Outcomes after Cerebral Ischemia Even if Administration is Delayed. <i>Thrombosis and Haemostasis</i> , 2019, 119, 467-478.	1.8	7
13	Comparison of accuracy of presepsin and procalcitonin concentrations in diagnosing sepsis in patients with and without acute kidney injury. <i>Clinica Chimica Acta</i> , 2019, 490, 200-206.	0.5	32
14	Soluble fibrin is a useful marker for predicting extracorporeal membrane oxygenation circuit exchange because of circuit clots. <i>Journal of Artificial Organs</i> , 2018, 21, 196-200.	0.4	11
15	Goreisan Prevents Brain Edema after Cerebral Ischemic Stroke by Inhibiting Aquaporin 4 Upregulation in Mice. <i>Journal of Stroke and Cerebrovascular Diseases</i> , 2018, 27, 758-763.	0.7	53
16	Association between medication adherence and illness perceptions in atrial fibrillation patients treated with direct oral anticoagulants: An observational cross-sectional pilot study. <i>PLoS ONE</i> , 2018, 13, e0204814.	1.1	43
17	Treatment of patients with sepsis in a closed intensive care unit is associated with improved survival: a nationwide observational study in Japan. <i>Journal of Intensive Care</i> , 2018, 6, 57.	1.3	17
18	Successful fishing for nucleus pulposus progenitor cells of the intervertebral disc across species. <i>JOR Spine</i> , 2018, 1, e1018.	1.5	44

#	ARTICLE	IF	CITATIONS
19	Fibrinogen level on admission is a predictor for massive transfusion in patients with severe blunt trauma: Analyses of a retrospective multicentre observational study. <i>Injury</i> , 2017, 48, 674-679.	0.7	28
20	Involvement of Charcot-Marie-Tooth disease gene mitofusin 2 expression in paclitaxel-induced mechanical allodynia in rats. <i>Neuroscience Letters</i> , 2017, 653, 337-340.	1.0	5
21	Potential survival benefit of polymyxin B hemoperfusion in patients with septic shock: a propensity-matched cohort study. <i>Critical Care</i> , 2017, 21, 134.	2.5	32
22	Adsorption of Nafamostat Mesilate on AN69ST Membranes: A Single-Center Retrospective and In Vitro Study. <i>Therapeutic Apheresis and Dialysis</i> , 2017, 21, 620-627.	0.4	9
23	Low-dose immunoglobulin G is not associated with mortality in patients with sepsis and septic shock. <i>Critical Care</i> , 2017, 21, 181.	2.5	19
24	Hyperfibrinolysis in severe isolated traumatic brain injury may occur without tissue hypoperfusion: a retrospective observational multicentre study. <i>Critical Care</i> , 2017, 21, 222.	2.5	29
25	Usefulness of plasminogen activator inhibitor-1 as a predictive marker of mortality in sepsis. <i>Journal of Intensive Care</i> , 2017, 5, 42.	1.3	42
26	Prognostic factors for candidaemia in intensive care unit patients: a retrospective analysis. <i>Singapore Medical Journal</i> , 2017, 58, 196-200.	0.3	7
27	Fibrinogen and base excess levels as predictive markers of the need for massive blood transfusion after blunt trauma. <i>Surgery Today</i> , 2016, 46, 774-779.	0.7	29
28	Characteristics, treatments, and outcomes of severe sepsis of 3195 ICU-treated adult patients throughout Japan during 2011-2013. <i>Journal of Intensive Care</i> , 2016, 4, 44.	1.3	29
29	High D-Dimer Levels Predict a Poor Outcome in Patients with Severe Trauma, Even with High Fibrinogen Levels on Arrival. <i>Shock</i> , 2016, 45, 308-314.	1.0	54
30	Recombinant human soluble thrombomodulin ameliorates cerebral ischemic injury through a high-mobility group box 1 inhibitory mechanism without hemorrhagic complications in mice. <i>Journal of the Neurological Sciences</i> , 2016, 362, 278-282.	0.3	18
31	Therapeutic Dose Monitoring for Linezolid in a Patient with MRSA Pneumonia with Bacteremia in Diabetes Insipidus. <i>Infectious Diseases and Therapy</i> , 2016, 5, 81-87.	1.8	1
32	Potential use of procalcitonin as biomarker for bacterial sepsis in patients with or without acute kidney injury. <i>Journal of Infection and Chemotherapy</i> , 2015, 21, 257-263.	0.8	42
33	Intravenous immunoglobulin improves sepsis-induced coagulopathy: A retrospective, single-center observational study. <i>Journal of Critical Care</i> , 2015, 30, 579-583.	1.0	15
34	Delayed treatment with ADAMTS13 ameliorates cerebral ischemic injury without hemorrhagic complication. <i>Brain Research</i> , 2015, 1624, 330-335.	1.1	21
35	Usefulness of presepsin in the diagnosis of sepsis in patients with or without acute kidney injury. <i>BMC Anesthesiology</i> , 2014, 14, 88.	0.7	75
36	Predicting the need for massive transfusion in trauma patients. <i>Journal of Trauma and Acute Care Surgery</i> , 2014, 76, 1243-1250.	1.1	72

#	ARTICLE	IF	CITATIONS
37	Exhaustion of nucleus pulposus progenitor cells with ageing and degeneration of the intervertebral disc. Nature Communications, 2012, 3, 1264.	5.8	357