

Andreas Hund

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3501021/publications.pdf>

Version: 2024-02-01

55
papers

3,046
citations

201385

27
h-index

174990

52
g-index

65
all docs

65
docs citations

65
times ranked

3331
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 1 | Plant phenotyping: from bean weighing to image analysis. <i>Plant Methods</i> , 2015, 11, 14. | 1.9 | 307 |
| 2 | Rooting depth and water use efficiency of tropical maize inbred lines, differing in drought tolerance. <i>Plant and Soil</i> , 2009, 318, 311-325. | 1.8 | 222 |
| 3 | The ETH field phenotyping platform FIP: a cable-suspended multi-sensor system. <i>Functional Plant Biology</i> , 2017, 44, 154. | 1.1 | 143 |
| 4 | Biochar amendment increases maize root surface areas and branching: a shovelomics study in Zambia. <i>Plant and Soil</i> , 2015, 395, 45-55. | 1.8 | 136 |
| 5 | Growth of axile and lateral roots of maize: I development of a phenotyping platform. <i>Plant and Soil</i> , 2009, 325, 335-349. | 1.8 | 135 |
| 6 | Remote, aerial phenotyping of maize traits with a mobile multi-sensor approach. <i>Plant Methods</i> , 2015, 11, 9. | 1.9 | 132 |
| 7 | QTL controlling root and shoot traits of maize seedlings under cold stress. <i>Theoretical and Applied Genetics</i> , 2004, 109, 618-629. | 1.8 | 130 |
| 8 | Global Wheat Head Detection (GWHD) Dataset: A Large and Diverse Dataset of High-Resolution RGB-Labelled Images to Develop and Benchmark Wheat Head Detection Methods. <i>Plant Phenomics</i> , 2020, 2020, 3521852. | 2.5 | 128 |
| 9 | Next generation shovelomics: set up a tent and REST. <i>Plant and Soil</i> , 2015, 388, 1-20. | 1.8 | 112 |
| 10 | A consensus map of QTLs controlling the root length of maize. <i>Plant and Soil</i> , 2011, 344, 143-158. | 1.8 | 98 |
| 11 | Rhizoslides: paper-based growth system for non-destructive, high throughput phenotyping of root development by means of image analysis. <i>Plant Methods</i> , 2014, 10, 13. | 1.9 | 95 |
| 12 | Ranking Quantitative Resistance to Septoria tritici Blotch in Elite Wheat Cultivars Using Automated Image Analysis. <i>Phytopathology</i> , 2018, 108, 568-581. | 1.1 | 88 |
| 13 | Mapping of QTLs for lateral and axile root growth of tropical maize. <i>Theoretical and Applied Genetics</i> , 2009, 119, 1413-1424. | 1.8 | 84 |
| 14 | Spectral Vegetation Indices to Track Senescence Dynamics in Diverse Wheat Germplasm. <i>Frontiers in Plant Science</i> , 2019, 10, 1749. | 1.7 | 80 |
| 15 | QTLs for the elongation of axile and lateral roots of maize in response to low water potential. <i>Theoretical and Applied Genetics</i> , 2010, 120, 621-631. | 1.8 | 71 |
| 16 | Cold tolerance of maize seedlings as determined by root morphology and photosynthetic traits. <i>European Journal of Agronomy</i> , 2008, 28, 178-185. | 1.9 | 65 |
| 17 | Hyperspectral Canopy Sensing of Wheat Septoria Tritici Blotch Disease. <i>Frontiers in Plant Science</i> , 2018, 9, 1195. | 1.7 | 61 |
| 18 | Root morphology and photosynthetic performance of maize inbred lines at low temperature. <i>European Journal of Agronomy</i> , 2007, 27, 52-61. | 1.9 | 60 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 19 | Global Wheat Head Detection 2021: An Improved Dataset for Benchmarking Wheat Head Detection Methods. <i>Plant Phenomics</i> , 2021, 2021, 9846158. | 2.5 | 60 |
| 20 | PhenoFly Planning Tool: flight planning for high-resolution optical remote sensing with unmanned areal systems. <i>Plant Methods</i> , 2018, 14, 116. | 1.9 | 49 |
| 21 | Image based phenotyping during winter: a powerful tool to assess wheat genetic variation in growth response to temperature. <i>Functional Plant Biology</i> , 2015, 42, 387. | 1.1 | 46 |
| 22 | Assessment of Multi-Image Unmanned Aerial Vehicle Based High-Throughput Field Phenotyping of Canopy Temperature. <i>Frontiers in Plant Science</i> , 2020, 11, 150. | 1.7 | 45 |
| 23 | An image analysis pipeline for automated classification of imaging light conditions and for quantification of wheat canopy cover time series in field phenotyping. <i>Plant Methods</i> , 2017, 13, 15. | 1.9 | 42 |
| 24 | Modern wheat semi-dwarfs root deep on demand: response of rooting depth to drought in a set of Swiss era wheats covering 100 years of breeding. <i>Euphytica</i> , 2019, 215, 1. | 0.6 | 38 |
| 25 | Precision Phenotyping Reveals Novel Loci for Quantitative Resistance to Septoria Tritici Blotch. <i>Plant Phenomics</i> , 2019, 2019, 3285904. | 2.5 | 37 |
| 26 | Collocations of QTLs for Seedling Traits and Yield Components of Tropical Maize under Water Stress Conditions. <i>Crop Science</i> , 2010, 50, 1385-1392. | 0.8 | 32 |
| 27 | Monitoring the dynamics of wheat stem elongation: genotypes differ at critical stages. <i>Euphytica</i> , 2017, 213, 1. | 0.6 | 32 |
| 28 | Cold Tolerance of the Photosynthetic Apparatus: Pleiotropic Relationship between Photosynthetic Performance and Specific Leaf Area of Maize Seedlings. <i>Molecular Breeding</i> , 2005, 16, 321-331. | 1.0 | 31 |
| 29 | Temperature response of wheat affects final height and the timing of stem elongation under field conditions. <i>Journal of Experimental Botany</i> , 2021, 72, 700-717. | 2.4 | 28 |
| 30 | Soil type determines how root and rhizosphere traits relate to phosphorus acquisition in field-grown maize genotypes. <i>Plant and Soil</i> , 2017, 412, 115-132. | 1.8 | 26 |
| 31 | In-Field Detection and Quantification of Septoria Tritici Blotch in Diverse Wheat Germplasm Using Spectral Temporal Features. <i>Frontiers in Plant Science</i> , 2019, 10, 1355. | 1.7 | 26 |
| 32 | Repeated Multiview Imaging for Estimating Seedling Tiller Counts of Wheat Genotypes Using Drones. <i>Plant Phenomics</i> , 2020, 2020, 3729715. | 2.5 | 26 |
| 33 | QTLs for early vigor of tropical maize. <i>Molecular Breeding</i> , 2010, 25, 91-103. | 1.0 | 25 |
| 34 | Early vertical distribution of roots and its association with drought tolerance in tropical maize. <i>Plant and Soil</i> , 2014, 377, 295-308. | 1.8 | 22 |
| 35 | RADIX: rhizoslide platform allowing high throughput digital image analysis of root system expansion. <i>Plant Methods</i> , 2016, 12, 40. | 1.9 | 22 |
| 36 | High-resolution quantification of root dynamics in split-nutrient rhizoslides reveals rapid and strong proliferation of maize roots in response to local high nitrogen. <i>Journal of Experimental Botany</i> , 2015, 66, 5507-5517. | 2.4 | 21 |

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 37 | High-throughput field phenotyping of soybean: Spotting an ideotype. <i>Remote Sensing of Environment</i> , 2022, 269, 112797. | 4.6 | 20 |
| 38 | Outdoor Plant Segmentation With Deep Learning for High-Throughput Field Phenotyping on a Diverse Wheat Dataset. <i>Frontiers in Plant Science</i> , 2021, 12, 774068. | 1.7 | 20 |
| 39 | Genetic diversity of Swiss maize (<i>Zea mays</i> L. ssp. <i>mays</i>) assessed with individuals and bulks on agarose gels. <i>Genetic Resources and Crop Evolution</i> , 2008, 55, 971-983. | 0.8 | 19 |
| 40 | Genetic structure and history of Swiss maize (<i>Zea mays</i> L. ssp. <i>mays</i>) landraces. <i>Genetic Resources and Crop Evolution</i> , 2010, 57, 71-84. | 0.8 | 19 |
| 41 | Opportunity costs for maize associated with localised application of sewage sludge derived fertilisers, as indicated by early root and phosphorus uptake responses. <i>Plant and Soil</i> , 2016, 406, 201-217. | 1.8 | 19 |
| 42 | Non-invasive field phenotyping of cereal development. <i>Burleigh Dodds Series in Agricultural Science</i> , 2019, , 249-292. | 0.1 | 19 |
| 43 | Temporal trends in canopy temperature and greenness are potential indicators of late-season drought avoidance and functional stay-green in wheat. <i>Field Crops Research</i> , 2021, 274, 108311. | 2.3 | 19 |
| 44 | Phenomics data processing: A plot-level model for repeated measurements to extract the timing of key stages and quantities at defined time points. <i>Field Crops Research</i> , 2021, 274, 108314. | 2.3 | 18 |
| 45 | Genetic variation in the gravitropic response of maize roots to low temperatures. <i>Plant Root</i> , 2010, 4, 22-30. | 0.3 | 16 |
| 46 | Can we improve heterosis for root growth of maize by selecting parental inbred lines with different temperature behaviour?. <i>Philosophical Transactions of the Royal Society B: Biological Sciences</i> , 2012, 367, 1580-1588. | 1.8 | 16 |
| 47 | Crop water use under Swiss pedoclimatic conditions – Evaluation of lysimeter data covering a seven-year period. <i>Field Crops Research</i> , 2017, 211, 48-65. | 2.3 | 16 |
| 48 | A soil-free root observation system for the study of root-microorganism interactions in maize. <i>Plant and Soil</i> , 2013, 367, 605-614. | 1.8 | 15 |
| 49 | Shovelomics root traits assessed on the EURoot maize panel are highly heritable across environments but show low genotype-by-nitrogen interaction. <i>Euphytica</i> , 2019, 215, 1. | 0.6 | 13 |
| 50 | The soil organic carbon stabilization potential of old and new wheat cultivars: a ^{13}C -CO $_2$ -labeling study. <i>Biogeosciences</i> , 2020, 17, 2971-2986. | 1.3 | 13 |
| 51 | A two-stage approach for the spatio-temporal analysis of high-throughput phenotyping data. <i>Scientific Reports</i> , 2022, 12, 3177. | 1.6 | 10 |
| 52 | Application of the <i>Prunus</i> spp. Cyanide Seed Defense System onto Wheat: Reduced Insect Feeding and Field Growth Tests. <i>Journal of Agricultural and Food Chemistry</i> , 2016, 64, 3501-3507. | 2.4 | 9 |
| 53 | Can Swiss wheat varieties escape future heat stress?. <i>European Journal of Agronomy</i> , 2021, 131, 126394. | 1.9 | 9 |
| 54 | Exploring genetic dependence of lipase activity to improve the quality of whole-grain wheat. <i>Journal of the Science of Food and Agriculture</i> , 2020, 100, 3120-3125. | 1.7 | 5 |

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 55 | Swiss maize landraces - Their diversity and genetic relationships. Acta Agronomica Hungarica: an International Multidisciplinary Journal in Agricultural Science, 2006, 54, 321-328. | 0.2 | 3 |