

Juergen Haas

List of Publications by Citations

Source: <https://exaly.com/author-pdf/3503711/juergen-haas-publications-by-citations.pdf>

Version: 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

13
papers

1,144
citations

12
h-index

13
g-index

13
ext. papers

1,242
ext. citations

4.3
avg, IF

3
L-index

#	Paper	IF	Citations
13	Reduced suppressive effect of CD4+CD25high regulatory T cells on the T cell immune response against myelin oligodendrocyte glycoprotein in patients with multiple sclerosis. <i>European Journal of Immunology</i> , 2005 , 35, 3343-52	6.1	319
12	Prevalence of newly generated naive regulatory T cells (Treg) is critical for Treg suppressive function and determines Treg dysfunction in multiple sclerosis. <i>Journal of Immunology</i> , 2007 , 179, 1322-30	5.3	190
11	In contrast to effector T cells, CD4+CD25+FoxP3+ regulatory T cells are highly susceptible to CD95 ligand- but not to TCR-mediated cell death. <i>Journal of Immunology</i> , 2005 , 175, 32-6	5.3	143
10	L-selectin is a possible biomarker for individual PML risk in natalizumab-treated MS patients. <i>Neurology</i> , 2013 , 81, 865-71	6.5	127
9	Glatiramer acetate improves regulatory T-cell function by expansion of naive CD4(+)CD25(+)FOXP3(+)CD31(+) T-cells in patients with multiple sclerosis. <i>Journal of Neuroimmunology</i> , 2009 , 216, 113-7	3.5	89
8	Interferon beta-induced restoration of regulatory T-cell function in multiple sclerosis is prompted by an increase in newly generated naive regulatory T cells. <i>Archives of Neurology</i> , 2008 , 65, 1434-9		75
7	Intracerebral human regulatory T cells: analysis of CD4+ CD25+ FOXP3+ T cells in brain lesions and cerebrospinal fluid of multiple sclerosis patients. <i>PLoS ONE</i> , 2011 , 6, e17988	3.7	68
6	Mutations in the COL5A1 coding sequence are not common in patients with spontaneous cervical artery dissections. <i>Stroke</i> , 1999 , 30, 1887-90	6.7	44
5	Specific recruitment of regulatory T cells into the CSF in lymphomatous and carcinomatous meningitis. <i>Blood</i> , 2008 , 111, 761-6	2.2	39
4	Similar sensitivity of regulatory T cells towards CD95L-mediated apoptosis in patients with multiple sclerosis and healthy individuals. <i>Journal of the Neurological Sciences</i> , 2006 , 251, 91-7	3.2	21
3	Single-cell PCR analysis of the immunoglobulin heavy-chain CDR3 region for the diagnosis of leptomeningeal involvement of B-cell malignancies using standard cerebrospinal fluid cytospins. <i>Journal of the Neurological Sciences</i> , 2004 , 219, 83-8	3.2	14
2	Rapid distinction of acute demyelinating disorders and central nervous system lymphoma by molecular analysis of cerebrospinal fluid cells. <i>Journal of Neurology</i> , 2001 , 248, 127-30	5.5	13
1	Identification by mRNA Differential Display of Two Up-regulated Genes as Candidate Mediators of AIDS Dementia. <i>Molecular Medicine</i> , 2001 , 7, 193-199	6.2	2