

# Iciar Martinez

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3505219/publications.pdf>

Version: 2024-02-01

57  
papers

1,704  
citations

257450

24  
h-index

289244

40  
g-index

58  
all docs

58  
docs citations

58  
times ranked

1546  
citing authors

#	ARTICLE	IF	CITATIONS
1	Antioxidant Activities and Selenogene Transcription in the European Sea Bass ( <i>Dicentrarchus labrax</i> ) Liver Depend, in a Non-linear Manner, on the Se/Hg Molar Ratio of the Feeds. <i>Biological Trace Element Research</i> , 2022, 200, 2365-2379.	3.5	1
2	Proteomic Strategies to Evaluate the Impact of Farming Conditions on Food Quality and Safety in Aquaculture Products. <i>Foods</i> , 2020, 9, 1050.	4.3	20
3	Protein Signatures to Trace Seafood Contamination and Processing. <i>Foods</i> , 2020, 9, 1751.	4.3	8
4	Reducing the Number of Individuals to Monitor Shoaling Fish Systems – Application of the Shannon Entropy to Construct a Biological Warning System Model. <i>Frontiers in Physiology</i> , 2018, 9, 493.	2.8	11
5	The Shannon Entropy Trend of a Fish System Estimated by a Machine Vision Approach Seems to Reflect the Molar Se:Hg Ratio of Its Feed. <i>Entropy</i> , 2018, 20, 90.	2.2	16
6	Estimation of Quality in Frozen Fish by Low Field NMR. , 2018, , 1901-1916.		0
7	Omega-3 Fatty Acid Content of Intact Muscle of Farmed Atlantic Salmon ( <i>Salmo salar</i> ) Examined by <sup>1</sup> H MAS NMR Spectroscopy. , 2018, , 1917-1925.		0
8	Shannon entropy of a european seabass ( <i>Dicentrarchus labrax</i> ) system in response to feed polluted with different concentrations of MeHg: a machine vision approach. , 2017, , .		0
9	Estimation of Quality in Frozen Fish by Low Field NMR. , 2017, , 1-16.		1
10	Shannon Entropy in a European Seabass ( <i>Dicentrarchus labrax</i> ) System during the Initial Recovery Period after a Short-Term Exposure to Methylmercury. <i>Entropy</i> , 2016, 18, 209.	2.2	14
11	Comparative study of muscle proteins in relation to the development of yake in three tropical tuna species yellowfin ( <i>Thunnus albacares</i> ), big eye ( <i>Thunnus obesus</i> ) and skipjack ( <i>Katsuwonus pelamis</i> ). <i>Food Chemistry</i> , 2016, 201, 284-291.	8.2	7
12	Evolution of Shannon entropy in a fish system (European seabass, <i>Dicentrarchus labrax</i> ) during the recuperation period after exposure to methylmercury. , 2015, , .		0
13	A paradigm shift in safe seafood production: From contaminant detection to fish monitoring – Application of biological warning systems to aquaculture. <i>Trends in Food Science and Technology</i> , 2015, 43, 104-113.	15.1	23
14	Low-Field Nuclear Magnetic Resonance of Proton ( <sup>1</sup> H LF NMR) Relaxometry for Monitoring the Time and Temperature History of Frozen Hake ( <i>Merluccius merluccius</i> L.) Muscle. <i>Food and Bioprocess Technology</i> , 2015, 8, 2137-2145.	4.7	30
15	The role of environmental biotechnology in exploring, exploiting, monitoring, preserving, protecting and decontaminating the marine environment. <i>New Biotechnology</i> , 2015, 32, 157-167.	4.4	48
16	Application of Entropy and Fractal Dimension Analyses to the Pattern Recognition of Contaminated Fish Responses in Aquaculture. <i>Entropy</i> , 2014, 16, 6133-6151.	2.2	52
17	Estimation of frozen storage time or temperature by kinetic modeling of the Kramer shear resistance and water holding capacity (WHC) of hake ( <i>Merluccius merluccius</i> , L.) muscle. <i>Journal of Food Engineering</i> , 2014, 120, 37-43.	5.2	36
18	Discrimination of contaminated fish responses by fractal dimension and entropy algorithms. , 2014, , .		0

#	ARTICLE	IF	CITATIONS
19	Development of a Real-Time PCR method for the identification of Atlantic mackerel ( <i>Scomber</i> ) Tj ETQq1 1 0.784314 rgBT /Overlock 10 T	8.2	21
20	Evaluation of Fish Quality and Safety by Proteomics Techniques. , 2013, , 161-180.		2
21	Evaluation of a Fast Method Based on the Presence of Two Restriction Sites in the Mitochondrial ND5 (mt ND5) Gene for the Identification of <i>Scomber</i> Species. Journal of Aquatic Food Product Technology, 2012, 21, 289-297.	1.4	1
22	Estimation of freezing storage time and quality changes in hake ( <i>Merluccius merluccius</i> , L.) by low field NMR. Food Chemistry, 2012, 135, 1626-1634.	8.2	135
23	Post-mortem degradation of myosin heavy chain in intact fish muscle: Effects of pH and enzyme inhibitors. Food Chemistry, 2011, 124, 1090-1095.	8.2	37
24	Protein expression and enzymatic activities in normal and soft textured Atlantic salmon ( <i>Salmo salar</i> ) muscle. Food Chemistry, 2011, 126, 140-148.	8.2	37
25	Partial characterisation of gelatinolytic activities in herring ( <i>Clupea harengus</i> ) and sardine ( <i>Sardina</i> ) Tj ETQq1 1 0.784314 rgBT /Overlock 10 T 675-683.	8.2	18
26	Myosin heavy chain degradation during post mortem storage of Atlantic cod ( <i>Gadus morhua</i> L.). Food Chemistry, 2009, 115, 1228-1233.	8.2	23
27	Proteolytic activities of ventral muscle and intestinal content of North Sea herring ( <i>Clupea</i> ) Tj ETQq1 1 0.784314 rgBT /Overlock 10 T	8.2	18
28	Identification of the farm origin of salmon by fatty acid and HR 13C NMR profiling. Food Chemistry, 2009, 116, 766-773.	8.2	24
29	Classification of Wild and Farmed Salmon Using Bayesian Belief Networks and Gas Chromatography-Derived Fatty Acid Distributions. Journal of Agricultural and Food Chemistry, 2009, 57, 7634-7639.	5.2	22
30	Analytical Methods to Differentiate Farmed from Wild Seafood. , 2009, , 215-232.		1
31	Non-destructive nuclear magnetic resonance image study of belly bursting in herring ( <i>Clupea</i> ) Tj ETQq1 1 0.784314 rgBT /Overlock 10 T	8.2	18
32	High resolution two-dimensional electrophoresis as a tool to differentiate wild from farmed cod ( <i>Gadus morhua</i> ) and to assess the protein composition of klipfish. Food Chemistry, 2007, 102, 504-510.	8.2	38
33	The genetic structure of <i>Pandalus borealis</i> in the Northeast Atlantic determined by RAPD analysis. ICES Journal of Marine Science, 2006, 63, 840-850.	2.5	14
34	Bioactive Compounds in Cod ( <i>Gadus morhua</i> ) Products and Suitability of 1H NMR Metabolite Profiling for Classification of the Products Using Multivariate Data Analyses. Journal of Agricultural and Food Chemistry, 2005, 53, 6889-6895.	5.2	53
35	High-resolution 1H magnetic resonance spectroscopy of whole fish, fillets and extracts of farmed Atlantic salmon ( <i>Salmo salar</i> ) for quality assessment and compositional analyses. Aquaculture, 2005, 250, 445-457.	3.5	50
36	Application of proteome analysis to seafood authentication. Proteomics, 2004, 4, 347-354.	2.2	98

#	ARTICLE	IF	CITATIONS
37	Genetic variability among isolates of <i>Listeria monocytogenes</i> from food products, clinical samples and processing environments, estimated by RAPD typing. <i>International Journal of Food Microbiology</i> , 2003, 84, 285-297.	4.7	46
38	Destructive and non-destructive analytical techniques for authentication and composition analyses of foodstuffs. <i>Trends in Food Science and Technology</i> , 2003, 14, 489-498.	15.1	87
39	Species identification of formed fishery products and high pressure-treated fish by electrophoresis: a collaborative study. <i>Food Chemistry</i> , 2001, 72, 105-112.	8.2	45
40	Post mortem muscle protein degradation during ice-storage of Arctic ( <i>Pandalus borealis</i> ) and tropical ( <i>Penaeus japonicus</i> and <i>Penaeus monodon</i> ) shrimps: a comparative electrophoretic and immunological study. <i>Journal of the Science of Food and Agriculture</i> , 2001, 81, 1199-1208.	3.5	54
41	Requirements for the application of protein sodium dodecyl sulfate-polyacrylamide gel electrophoresis and randomly amplified polymorphic DNA analyses to product speciation. <i>Electrophoresis</i> , 2001, 22, 1526-1533.	2.4	18
42	Identification of marine mammal species in food products. <i>Journal of the Science of Food and Agriculture</i> , 2000, 80, 527-533.	3.5	30
43	Species identification of smoked and gravad fish products by sodium dodecylsulphate polyacrylamide gel electrophoresis, urea isoelectric focusing and native isoelectric focusing: a collaborative study. <i>Food Chemistry</i> , 2000, 71, 1-7.	8.2	69
44	Identification of Fish Species after Cooking by SDS-PAGE and Urea IEF: A Collaborative Study. <i>Journal of Agricultural and Food Chemistry</i> , 2000, 48, 2653-2658.	5.2	94
45	RAPD-typing of Central and Eastern North Atlantic and Western North Pacific minke whales, <i>Balaenoptera acutorostrata</i> . <i>ICES Journal of Marine Science</i> , 1999, 56, 640-651.	2.5	11
46	Species identification of cooked fish by urea isoelectric focusing and sodium dodecylsulfate polyacrylamide gel electrophoresis. <i>Food Chemistry</i> , 1999, 67, 333-339.	8.2	44
47	Development of a sodium dodecyl sulfate-polyacrylamide gel electrophoresis reference method for the analysis and identification of fish species in raw and heat-processed samples: A collaborative study. <i>Electrophoresis</i> , 1999, 20, 1425-1432.	2.4	92
48	RAPD and scnDNA analyses of polar cod, <i>Boreogadus saida</i> (Pisces, Galidae), in the north Atlantic. <i>Sarsia</i> , 1999, 84, 99-103.	0.5	17
49	Species identification in meat products by RAPD analysis. <i>Food Research International</i> , 1998, 31, 459-466.	6.2	77
50	Seafood: Fulfilling Market Demands. <i>Outlook on Agriculture</i> , 1997, 26, 107-114.	3.4	3
51	Sample preparation and DNA extraction procedures for polymerase chain reaction identification of <i>Listeria monocytogenes</i> in seafoods. <i>International Journal of Food Microbiology</i> , 1997, 35, 275-280.	4.7	47
52	Effects of T3 and rearing temperature on growth and skeletal myosin heavy chain isoform transition during early development in the salmonid <i>Salvelinus alpinus</i> (L.). <i>Comparative Biochemistry and Physiology - B Biochemistry and Molecular Biology</i> , 1995, 112, 717-725.	1.6	23
53	Myofibrillar proteins in developing white muscle of the Arctic charr, <i>Salvelinus alpinus</i> (L.). <i>Comparative Biochemistry and Physiology Part B: Comparative Biochemistry</i> , 1994, 107, 11-20.	0.2	5
54	Myofibrillar proteins in skeletal muscles of parr, smolt and adult atlantic salmon ( <i>Salmo salar</i> l.). Comparison with another salmonid, the arctic charr <i>Salvelinus alpinus</i> (l.). <i>Comparative Biochemistry and Physiology Part B: Comparative Biochemistry</i> , 1993, 106, 1021-1028.	0.2	11

#	ARTICLE	IF	CITATIONS
55	Comparison of myosin isoenzymes present in skeletal and cardiac muscles of the Arctic charr <i>Salvelinus alpinus</i> (L.). Sequential expression of different myosin heavy chains during development of the fast white skeletal muscle. <i>FEBS Journal</i> , 1991, 195, 743-753.	0.2	42
56	Water retention properties and solubility of the myofibrillar proteins: Interrelationships and possible value as indicators of the gel strength in cod surimi by a multivariate data analysis. <i>Journal of the Science of Food and Agriculture</i> , 1989, 46, 469-479.	3.5	8
57	Evolution of Shannon entropy in a fish system (European seabass, <i>Dicentrarchus labrax</i> )	1.0784314	10