

Wen-Ko Tseng

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/350897/publications.pdf>

Version: 2024-02-01

11
papers

128
citations

1684188

5
h-index

1372567

10
g-index

11
all docs

11
docs citations

11
times ranked

308
citing authors

#	ARTICLE	IF	CITATIONS
1	Pretreatment serum interleukin-1 β , interleukin-6, and tumor necrosis factor- α levels predict the progression of colorectal cancer. <i>Cancer Medicine</i> , 2016, 5, 426-433.	2.8	51
2	Clinical Significance of Serum Glutamine Level in Patients with Colorectal Cancer. <i>Nutrients</i> , 2019, 11, 898.	4.1	25
3	The Ratio of Hmox1/Nrf2 mRNA Level in the Tumor Tissue Is a Predictor of Distant Metastasis in Colorectal Cancer. <i>Disease Markers</i> , 2016, 2016, 1-6.	1.3	17
4	Relationship between pre-treatment nutritional status, serum glutamine, arginine levels and clinicopathological features in Taiwan colorectal cancer patients. <i>Asia Pacific Journal of Clinical Nutrition</i> , 2015, 24, 598-604.	0.4	10
5	Associations between the Nrf2/Keap1 pathway and mitochondrial functions in colorectal cancer are affected by metastasis. <i>Cancer Biomarkers</i> , 2020, 27, 163-171.	1.7	9
6	The ratio of thioredoxin/Keap1 protein level is a predictor of distant metastasis in colorectal cancer. <i>Biomarkers in Medicine</i> , 2017, 11, 1103-1111.	1.4	5
7	Correlation Between the Glasgow Prognostic Score and the Serum Cytokine Profile in Taiwanese Patients with Colorectal Cancer. <i>International Journal of Biological Markers</i> , 2021, 36, 40-49.	1.8	4
8	Colonoscopy-induced right superior rectal artery tear: A case report. <i>International Journal of Surgery Case Reports</i> , 2017, 41, 47-49.	0.6	3
9	The level of S-glutathionylated protein is a predictor for metastasis in colorectal cancer and correlated with those of Nrf2/Keap1 pathway. <i>Biomarkers</i> , 2021, 26, 780-787.	1.9	3
10	Pretreatment Nutrition-Inflammation Biomarkers Correlated with Differential Cytokine Profiles in Taiwanese Patients with Colorectal Cancer. <i>Nutrition and Cancer</i> , 2021, , 1-11.	2.0	1
11	The tumor/normal tissue ratio of Keap1 protein is a predictor for lymphovascular invasion in colorectal cancer: A correlation study between the Nrf2 and KRas pathways. <i>Biomarkers</i> , 0, , 1-12.	1.9	0