## MichaÅ, Saniewski

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/3519215/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Effect of drying, blanching, pickling and maceration on the fate of 40K, total K and 137Cs in bolete mushrooms and dietary intake. Environmental Science and Pollution Research, 2022, 29, 742-754.	5.3	12
2	Benthic macroinvertebrates as reference indicators for monitoring of anthropogenic isotope 137Cs contamination in the marine environment. Environmental Science and Pollution Research, 2022, 29, 13822-13834.	5.3	4
3	Radiocaesium in Tricholoma spp. from the Northern Hemisphere in 1971–2016. Science of the Total Environment, 2022, 802, 149829.	8.0	10
4	137Cs and 40K activity concentrations in edible wild mushrooms from China regions during the 2014–2016 period. Foods and Raw Materials, 2022, , 86-96.	2.1	5
5	Alimentary exposure and elimination routes of rare earth elements (REE) in marine mammals from the Baltic Sea and Antarctic coast. Science of the Total Environment, 2021, 754, 141947.	8.0	12
6	An evaluation of the occurrence and trends in 137Cs and 40K radioactivity in King Bolete Boletus edulis mushrooms in Poland during 1995–2019. Environmental Science and Pollution Research, 2021, 28, 32405-32415.	5.3	15
7	137Cs and 40K activities and total K distribution in the sclerotia of the Wolfiporia cocos fungus from China. Journal of Environmental Radioactivity, 2021, 231, 106549.	1.7	9
8	Geochronology of the southern Baltic Sea sediments derived from 210Pb dating. Quaternary Geochronology, 2020, 56, 101039.	1.4	4
9	137Caesium, 40Potassium and potassium in raw and deep-oil stir-fried mushroom meals from Yunnan in China. Journal of Food Composition and Analysis, 2020, 91, 103538.	3.9	9
10	Bryophytes and lichens as fallout originated radionuclide indicators in the Svalbard archipelago (High Arctic). Polar Science, 2020, 25, 100536.	1.2	13
11	137Cs, 40K, and K in raw and stir-fried mushrooms from the Boletaceae family from the Midu region in Yunnan, Southwest China. Environmental Science and Pollution Research, 2020, 27, 32509-32517.	5.3	17
12	Artificial (137Cs) and natural (40K) radioactivity and total potassium in medicinal fungi from Yunnan in China. Isotopes in Environmental and Health Studies, 2020, 56, 324-333.	1.0	7
13	Meteorological phenomenon as a key factor controlling variability of labile particulate mercury in rivers and its inflow into coastal zone of the sea. Environmental Research, 2020, 184, 109355.	7.5	5
14	Impact of distance from the glacier on the content of 137Cs and 90Sr in the lichen Cetrariella delisei. Chemosphere, 2020, 259, 127433.	8.2	9
15	Anthropogenic radioactive isotopes in Actiniaria from the Svalbard archipelago. Marine Pollution Bulletin, 2020, 157, 111369.	5.0	3
16	137Cs and 40K in gray seals Halichoerus grypus in the southern Baltic Sea. Environmental Science and Pollution Research, 2019, 26, 17418-17426.	5.3	0
17	Radiocaesium pollution of fly agaric Amanita muscaria in fruiting bodies decreases with developmental stage. Isotopes in Environmental and Health Studies, 2019, 55, 317-324.	1.0	24
18	Budget of 90Sr in the Gulf of Gdańsk (southern Baltic Sea). Oceanologia, 2018, 60, 256-263.	2.2	8

MichaÅ, Saniewski

#	Article	IF	CITATIONS
19	Impact of intense rains and flooding on mercury riverine input to the coastal zone. Marine Pollution Bulletin, 2018, 127, 593-602.	5.0	24
20	Artificial 137Cs and natural 40K in mushrooms from the subalpine region of the Minya Konka summit and Yunnan Province in China. Environmental Science and Pollution Research, 2018, 25, 615-627.	5.3	36
21	90 Sr and 137 Cs in Arctic echinoderms. Marine Pollution Bulletin, 2017, 124, 563-568.	5.0	7
22	90Sr in Zostera marina from the Gulf of Gdańsk (southern Baltic Sea). Oceanological and Hydrobiological Studies, 2017, 46, 24-29.	0.7	4
23	90Sr in fish from the southern Baltic Sea, coastal lagoons and freshwater lake. Journal of Environmental Radioactivity, 2016, 158-159, 38-46.	1.7	7
24	90Sr in King Bolete Boletus edulis and certain other mushrooms consumed in Europe and China. Science of the Total Environment, 2016, 543, 287-294.	8.0	33
25	Atmospheric deposition and riverine load of 90 Sr and 137 Cs to the Gulf of Gdańsk (southern Baltic) Tj ETQq1	0.78431 1.7	4 rgBT /Over
26	Macrophyta as a vector of contemporary and historical mercury from the marine environment to the trophic web. Environmental Science and Pollution Research, 2015, 22, 5228-5240.	5.3	37
27	The impact of land use and season on the riverine transport of mercury into the marine coastal zone. Environmental Monitoring and Assessment, 2014, 186, 7593-7604.	2.7	31
28	Mercury loads into the sea associated with extreme flood. Environmental Pollution, 2014, 191, 93-100.	7.5	57
29	Spatiotemporal Variations of the <sup><b>90</b></sup> Sr in the Southern Part of the Baltic Sea over the Period of 2005–2010. Scientific World Journal, The, 2013, 2013, 1-8.	2.1	4
30	MACROPHYTOBENTHOS AS AN INDICATOR OF THE ENVIRONMENTAL STATUS OF THE BALTIC SEA. Polish Hyperbaric Research, 2013, , .	0.1	1
31	Bioaccumulation of 137Cs by benthic plants and macroinvertebrates. Oceanological and Hydrobiological Studies, 2011, 40, 1-8.	0.7	14
32	Distribution of mercury in different environmental compartments in the aquatic ecosystem of the coastal zone of the Southern Baltic Sea. Journal of Environmental Sciences, 2010, 22, 1144-1150.	6.1	30