Charles E Savrda

List of Publications by Year
in descending order

32 papers	1,498 citations	471509 17 h-index	501196 28 g-index
32 all docs	32 docs citations	32 times ranked	779 citing authors

#	Article	IF	Citations
1	Firmground crustacean burrow systems (Glossifungites ichnofacies) in marine shelf deposits, Paleocene Clayton Formation, Alabama, USA. Lethaia, 2020, 53, 500-514.	1.4	2
2	Shipworm bioerosion of lithic substrates in a freshwater setting, Abatan River, Philippines: Ichnologic, paleoenvironmental and biogeomorphical implications. PLoS ONE, 2019, 14, e0224551.	2.5	5
3	Bioerosion of a modern bedrock stream bed by insect larvae (Conecuh River, Alabama): Implications for ichnotaxonomy, continental ichnofacies, and biogeomorphology. Palaeogeography, Palaeoclimatology, Palaeoecology, 2019, 513, 3-13.	2.3	13
4	REVISITING THE ORIGINS OF CLAYTON SAND BODIES AT THE Kâ€“PG TRANSITION, MOSCOW LANDING, WESTERN ALABAMA: STRATIGRAPHIC RELATIONS, SEDIMENTOLOGY, AND ICHNOLOGY. Palaios, 2018, 33, 555-567.	1.3	8
5	Chronostratigraphy and environment of Furnas Formation by trace fossil analysis: Calibrating the lower Paleozoic Gondwana realm in the ParanÃ¡ Basin (Brazil). Palaeogeography, Palaeoclimatology, Palaeoecology, 2017, 487, 307-320.	2.3	27
6	Rhizocorallium in estuarine Ingersoll shale (Upper Cretaceous Eutaw Formation, Eastern Alabama) Tj ETQq0 0 0 rgBT /Overlock 10 Tf 50	1.4	3
7	COMPOSITE ICHNOFABRICS: CATEGORIZATION BASED ON NUMBER OF ICHNOCOENOSES AND THEIR TEMPORAL INCONGRUENCE. Palaios, 2016, 31, 92-96.	1.3	16
8	Chalk and Related Deep-Marine Carbonates. Developments in Sedimentology, 2012, 64, 777-806.	0.5	18
9	A new Upper Cretaceous (Santonian) amber deposit from the Eutaw Formation of eastern Alabama, USA. Cretaceous Research, 2010, 31, 85-93.	1.4	22
10	The Prospect of Compact Estuarine LagerstÃtten. The Sedimentary Record, 2009, 7, 4-8.	0.6	8
11	Taphonomy of Trace Fossils. , 2007, , 92-109.		35
12	Trace Fossil Preservation by Native Copper, Corocoro, Bolivia. Rocks and Minerals, 2006, 81, 362-363.	0.1	1
13	Ichnology of fair-weather and storm deposits in an Upper Cretaceous estuary (Eutaw Formation,) Tj ETQq1 1 0.784314 rgBT /Overlock	2.3	38
14	Ichnofabrics of a Pleistocene slope succession, New Jersey margin: relations to climate and sea-level dynamics. Palaeogeography, Palaeoclimatology, Palaeoecology, 2001, 171, 41-61.	2.3	64
15	Taenidiumand Associated Ichnofossils in Fluvial Deposits, Cretaceous Tuscaloosa Formation, Eastern Alabama, Southeastern U.S.A. Ichnos, 2000, 7, 227-242.	0.5	40
16	Ichnology of Rhythmically Bedded Demopolis Chalk (Upper Cretaceous, Alabama): Implications for Paleoenvironment, Depositional Cycle Origins, and Tracemaker Behavior. Palaios, 1998, 13, 423.	1.3	51
17	Ichnofabrics, ichnocoenoses, and ichnofacies implications of an upper cretaceous tidalâ€inlet sequence (Eutaw formation, central Alabama). Ichnos, 1998, 6, 53-74.	0.5	14
18	Ichnofossils: Linkages to life habitats and environmentsâ€•An introduction to symposium papers in the current issue. Ichnos, 1998, 6, 1-3.	0.5	0

#	Article	IF	Citations
19	Ichnofossil tiering analysis of a rhythmically bedded chalkâ€marl sequence in the Upper Cretaceous of Alabama. Lethaia, 1998, 31, 311-321.	1.4	13
20	Morphological Variation in <u>Teredolites</u>: Implications for Wood-Boring Bivalve Behavior. The Paleontological Society Special Publications, 1996, 8, 341-341.	0.0	0
21	Behavioral implications of branching and tubeâ€lining inTeredolites. Ichnos, 1996, 4, 191-198.	0.5	24
22	Ichnologic Applications in Paleoceanographic, Paleoclimatic, and Sea-Level Studies. Palaios, 1995, 10, 565.	1.3	97
23	Logâ€ground andTeredoliteslagerstatte in a Trangressive Sequence, Upper Cretaceous (Lower) Tj ETQq1 1 0.784314 rgBT /Ov	0.5	29
24	Trace Fossils and Benthic Oxygenation. Short Courses in Paleontology, 1992, 5, 172-196.	0.2	37
25	Ichnology in Sequence Stratigraphic Studies: An Example from the Lower Paleocene of Alabama. Palaios, 1991, 6, 39.	1.3	78
26	Teredolites, wood substrates, and sea-level dynamics. Geology, 1991, 19, 905.	4.4	64
27	Oxygen-related biofacies in marine strata: an overview and update. Geological Society Special Publication, 1991, 58, 201-219.	1.3	106
28	Trace-fossil model for reconstructing oxygenation histories of ancient marine bottom waters: Application to upper cretaceous niobrara formation, Colorado. Palaeogeography, Palaeoclimatology, Palaeoecology, 1989, 74, 49-74.	2.3	152
29	The exaerobic zone, a new oxygen-deficient marine biofacies. Nature, 1987, 327, 54-56.	27.8	137
30	Trace-fossil model for reconstruction of paleo-oxygenation in bottom waters. Geology, 1986, 14, 3.	4.4	333
31	Rhythmic bedding produced in Cretaceous pelagic carbonate environments: Sensitive recorders of climatic cycles. Paleoceanography, 1986, 1, 467-481.	3.0	53
32	Oxygen-Related Mudrock Biofacies. , 0, , 92-102.		10