Xinzhi Ni

List of Publications by Year in Descending Order

Source: https://exaly.com/author-pdf/3558145/xinzhi-ni-publications-by-year.pdf

Version: 2024-04-09

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

1,926 40 102 23 h-index g-index citations papers 4.64 2,404 109 3.2 avg, IF L-index ext. citations ext. papers

#	Paper	IF	Citations
102	A sorghum genome-wide association study (GWAS) identifies a WRKY transcription factor as a candidate gene underlying sugarcane aphid (Melanaphis sacchari) resistance <i>Planta</i> , 2022 , 255, 37	4.7	О
101	Spatio-temporal patterns of Aspergillus flavus infection and aflatoxin B biosynthesis on maize kernels probed by SWIR hyperspectral imaging and synchrotron FTIR microspectroscopy <i>Food Chemistry</i> , 2022 , 382, 132340	8.5	1
100	Melanaphisßorghi (Hemiptera: Aphididae) Clonal Diversity in the United States and Brazil. <i>Insects</i> , 2022 , 13, 416	2.8	1
99	Inductive and synergistic interactions between plant allelochemical flavone and Bt toxin Cry1Ac in Helicoverpa armigera. <i>Insect Science</i> , 2021 , 28, 1756-1765	3.6	
98	Extended investigation of field-evolved resistance of the corn earworm Helicoverpa zea(Lepidoptera: Noctuidae) to Bacillus thuringiensis Cry1A.105 and Cry2Ab2 proteins in thesoutheastern United States. <i>Journal of Invertebrate Pathology</i> , 2021 , 183, 107560	2.6	3
97	Combining host plant resistance and foliar insecticide application to manage Melanaphis sacchari (Hemiptera: Aphididae) in grain sorghum. <i>International Journal of Pest Management</i> , 2021 , 67, 10-19	1.5	4
96	A Sugarcane Aphid (Hemiptera: Aphididae) Buper-ClonelRemains on U.S. Sorghum and Johnsongrass and Feeds on Giant Miscanthus. <i>Journal of Entomological Science</i> , 2021 , 56, 43-52	0.4	4
95	Populations of (Boddie) in the Southeastern United States are Commonly Resistant to Cry1Ab, but Still Susceptible to Vip3Aa20 Expressed in MIR 162 Corn. <i>Toxins</i> , 2021 , 13,	4.9	4
94	Assessing spatio-temporal patterns of sugarcane aphid (Hemiptera: Aphididae) infestations on silage sorghum yield using unmanned aerial systems (UAS). <i>Crop Protection</i> , 2021 , 146, 105681	2.7	2
93	Spontaneous transposition of HzSINE1 into CYP321A2 is undetectable in the field populations of Helicoverpa zea. <i>Journal of Asia-Pacific Entomology</i> , 2021 , 24, 882-888	1.4	
92	Characterization of the First W-Specific Protein-Coding Gene for Sex Identification in. <i>Frontiers in Genetics</i> , 2020 , 11, 649	4.5	2
91	The sorghum epicuticular wax locus Bloomless2 reduces plant damage in P898012 caused by the sugarcane aphid 2020 , 3, e20008		0
90	Classifying maize kernels naturally infected by fungi using near-infrared hyperspectral imaging. <i>Infrared Physics and Technology</i> , 2020 , 105, 103242	2.7	15
89	Functional Biology and Molecular Mechanisms of Host-Pathogen Interactions for Aflatoxin Contamination in Groundnut (L.) and Maize (L.). <i>Frontiers in Microbiology</i> , 2020 , 11, 227	5.7	22
88	Parasitism of Melanaphis sacchari (Hemiptera: Aphididae) by Lysiphlebus testaceipes (Hymenoptera: Braconidae) in the Greenhouse and Field. <i>Journal of Entomological Science</i> , 2020 , 55, 14	0.4	2
87	Evaluation and classification of five cereal fungi on culture medium using Visible/Near-Infrared (Vis/NIR) hyperspectral imaging. <i>Infrared Physics and Technology</i> , 2020 , 105, 103206	2.7	11
86	Non-destructive discrimination of Illicium verum from poisonous adulterant using Vis/NIR hyperspectral imaging combined with chemometrics. <i>Infrared Physics and Technology</i> , 2020 , 111, 103509	9 ^{2.7}	2

(2018-2020)

85	Evaluation of strains of Beauveria bassiana and Isaria fumosorosea to control sugarcane aphids on grain sorghum 2020 , 3, e20047		3	
84	Genome Size Reversely Correlates With Host Plant Range in Species. <i>Frontiers in Physiology</i> , 2019 , 10, 29	4.6	3	
83	A long non-coding RNA regulates cadherin transcription and susceptibility to Bt toxin Cry1Ac in pink bollworm, Pectinophora gossypiella. <i>Pesticide Biochemistry and Physiology</i> , 2019 , 158, 54-60	4.9	14	
82	Molecular evolution of the plant ECERIFERUM1 and ECERIFERUM3 genes involved in aliphatic hydrocarbon production. <i>Computational Biology and Chemistry</i> , 2019 , 80, 1-9	3.6	7	
81	Field-evolved resistance of Helicoverpa zea (Boddie) to transgenic maize expressing pyramided Cry1A.105/Cry2Ab2 proteins in northeast Louisiana, the United States. <i>Journal of Invertebrate Pathology</i> , 2019 , 163, 11-20	2.6	41	
80	Evaluation of Elite Maize Inbred Lines for Reduced Aspergillus flavus Infection, Aflatoxin Accumulation, and Agronomic Traits. <i>Crop Science</i> , 2019 , 59, 2562-2571	2.4	2	
79	Decreased Cry1Ac activation by midgut proteases associated with Cry1Ac resistance in Helicoverpa zea. <i>Pest Management Science</i> , 2019 , 75, 1099-1106	4.6	13	
78	Monitoring of brown stink bug (Hemiptera: Pentatomidae) population dynamics in corn to predict its abundance using weather data. <i>Insect Science</i> , 2019 , 26, 536-544	3.6	3	
77	Deciphering drought-induced metabolic responses and regulation in developing maize kernels. <i>Plant Biotechnology Journal</i> , 2018 , 16, 1616	11.6	45	
76	The Environment Strongly Affects Estimates of Heterosis in Hybrid Sweet Sorghum. <i>Sugar Tech</i> , 2018 , 20, 261-274	1.9	6	
75	Cytotoxicity and binding profiles of activated Cry1Ac and Cry2Ab to three insect cell lines. <i>Insect Science</i> , 2018 , 25, 655-666	3.6	12	
74	Genome-Wide Association Mapping of Anthracnose () Resistance in the U.S. Sorghum Association Panel. <i>Plant Genome</i> , 2018 , 11, 170099	4.4	23	
73	Useful Bicistronic Reporter System for Studying Poly(A) Site-Defining cis Elements and Regulation of Alternative Polyadenylation. <i>International Journal of Molecular Sciences</i> , 2018 , 19,	6.3	3	
72	Growth Identification of Aspergillus flavus and Aspergillus parasiticus by Visible/Near-Infrared Hyperspectral Imaging. <i>Applied Sciences (Switzerland)</i> , 2018 , 8, 513	2.6	4	
71	Recent advancement in near infrared spectroscopy and hyperspectral imaging techniques for quality and safety assessment of agricultural and food products in the China Agricultural University. <i>NIR News</i> , 2018 , 29, 19-23	0.8	1	
70	Evaluation of growth characteristics of Aspergillus parasiticus inoculated in different culture media by shortwave infrared (SWIR) hyperspectral imaging. <i>Journal of Innovative Optical Health Sciences</i> , 2018 , 11, 1850031	1.2		
69	A Sugarcane Aphid Buper-ClonelPredominates on Sorghum and Johnsongrass from Four US States. <i>Crop Science</i> , 2018 , 58, 2533-2541	2.4	6	
68	Evaluation of Near-Infrared Hyperspectral Imaging for Detection of Peanut and Walnut Powders in Whole Wheat Flour. <i>Applied Sciences (Switzerland)</i> , 2018 , 8, 1076	2.6	17	

67	Evaluation of maize inbred lines for resistance to pre-harvest aflatoxin and fumonisin contamination in the field. <i>Crop Journal</i> , 2017 , 5, 259-264	4.6	14
66	The Genes Bm2 and Blmc that Affect Epicuticular Wax Deposition in Sorghum are Allelic. <i>Crop Science</i> , 2017 , 57, 1552-1556	2.4	7
65	Microsatellite Markers Reveal a Predominant Sugarcane Aphid (Homoptera: Aphididae) Clone is Found on Sorghum in Seven States and One Territory of the USA. <i>Crop Science</i> , 2017 , 57, 2064-2072	2.4	28
64	Identification of Resistance to Aflatoxin Accumulation and Yield Potential in Maize Hybrids in the Southeast Regional Aflatoxin Trials (SERAT). <i>Crop Science</i> , 2017 , 57, 202-215	2.4	11
63	F2 screen for resistance to Bacillus thuringiensis Cry2Ab2-maize in field populations of Spodoptera frugiperda (Lepidoptera: Noctuidae) from the southern United States. <i>Journal of Invertebrate Pathology</i> , 2016 , 138, 66-72	2.6	16
62	Structure-reactivity relationships between the fluorescent chromophores and antioxidant activity of grain and sweet sorghum seeds. <i>Food Science and Nutrition</i> , 2016 , 4, 811-817	3.2	4
61	APN1 is a functional receptor of Cry1Ac but not Cry2Ab in Helicoverpa zea. <i>Scientific Reports</i> , 2016 , 6, 19179	4.9	17
60	Diurnal Activities of the Brown Stink Bug (Hemiptera: Pentatomidae) in and near Tasseling Corn Fields. <i>Journal of Entomological Science</i> , 2016 , 51, 226-237	0.4	2
59	Feasibility of detecting Aflatoxin B1 in single maize kernels using hyperspectral imaging. <i>Journal of Food Engineering</i> , 2015 , 166, 182-192	6	45
58	Maize death acids, 9-lipoxygenase-derived cyclopente(a)nones, display activity as cytotoxic phytoalexins and transcriptional mediators. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2015 , 112, 11407-12	11.5	83
57	A re-examination of corn (Zea mays L.) ear volatiles. <i>Phytochemistry Letters</i> , 2015 , 14, 280-286	1.9	4
56	Near-infrared hyperspectral imaging for detecting Aflatoxin B1 of maize kernels. <i>Food Control</i> , 2015 , 51, 347-355	6.2	39
55	Evaluation of Whorl Damage by Fall Armyworm (Lepidoptera: Noctuidae) on Field- and Greenhouse-Grown Sweet Sorghum Plants. <i>Journal of Entomological Science</i> , 2015 , 50, 14-27	0.4	1
54	Feasibility of detecting aflatoxin B1 on inoculated maize kernels surface using Vis/NIR hyperspectral imaging. <i>Journal of Food Science</i> , 2015 , 80, M116-22	3.4	35
53	Stress Sensitivity Is Associated with Differential Accumulation of Reactive Oxygen and Nitrogen Species in Maize Genotypes with Contrasting Levels of Drought Tolerance. <i>International Journal of Molecular Sciences</i> , 2015 , 16, 24791-819	6.3	26
52	Influence of host plant nitrogen fertilization on hemolymph protein profiles of herbivore Spodoptera exigua and development of its endoparasitoid Cotesia marginiventris. <i>Biological Control</i> , 2014 , 70, 9-16	3.8	5
51	Influence of brown stink bug feeding, planting date and sampling time on common smut infection of maize. <i>Insect Science</i> , 2014 , 21, 564-71	3.6	3
50	Changes in life history parameters of Rhopalosiphum maidis (Homoptera: Aphididae) under four different elevated temperature and CO2 combinations. <i>Journal of Economic Entomology</i> , 2014 , 107, 14	17-8	23

(2010-2014)

49	Evaluation of fall armyworm resistance in maize germplasm lines using visual leaf injury rating and predator survey. <i>Insect Science</i> , 2014 , 21, 541-55	3.6	19
48	Integrated pest management is the lucrative bridge connecting the ever emerging knowledge islands of genetics and ecology. <i>Insect Science</i> , 2014 , 21, 537-40	3.6	2
47	Effects of photoperiod and temperature on diapause induction in Conogethes punctiferalis (Lepidoptera: Pyralidae). <i>Insect Science</i> , 2014 , 21, 556-63	3.6	8
46	Environmental influences on maize-Aspergillus flavus interactions and aflatoxin production. <i>Frontiers in Microbiology</i> , 2014 , 5, 40	5.7	71
45	Evaluation of spatial and temporal patterns of insect damage and aflatoxin level in the pre-harvest corn fields to improve management tactics. <i>Insect Science</i> , 2014 , 21, 572-83	3.6	2
44	Identification of aflatoxin B1 on maize kernel surfaces using hyperspectral imaging. <i>Food Control</i> , 2014 , 42, 78-86	6.2	63
43	Potential shortfall of pyramided transgenic cotton for insect resistance management. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2013 , 110, 5806-11	11.5	123
42	Evaluation of corn germplasm lines for multiple ear-colonizing insect and disease resistance. <i>Journal of Economic Entomology</i> , 2012 , 105, 1457-64	2.2	12
41	Crop Stress and Aflatoxin Contamination: Perspectives and Prevention Strategies 2012, 399-427		2
40	Potential adaptation of a Q biotype whitefly population from poinsettia to field crops. <i>Insect Science</i> , 2011 , 18, 719-728	3.6	8
39	Impact of applying edible oils to silk channels on ear pests of sweet corn. <i>Journal of Economic Entomology</i> , 2011 , 104, 956-64	2.2	4
38	Novel acidic sesquiterpenoids constitute a dominant class of pathogen-induced phytoalexins in maize. <i>Plant Physiology</i> , 2011 , 156, 2082-97	6.6	156
37	Foliar Resistance to Fall Armyworm in Corn Germplasm Lines that Confer Resistance to Root- and Ear-Feeding Insects*. <i>Florida Entomologist</i> , 2011 , 94, 971-981	1	16
36	Identity, regulation, and activity of inducible diterpenoid phytoalexins in maize. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2011 , 108, 5455-60	11.5	179
35	Spatial patterns of aflatoxin levels in relation to ear-feeding insect damage in pre-harvest corn. <i>Toxins</i> , 2011 , 3, 920-31	4.9	26
34	Metamorphosis of Cisgenic Insect Resistance Research in the Transgenic Crop Era 2011 , 258-279		1
33	Aflatoxin Accumulation in BT and Non-BT Maize Testcrosses. <i>Journal of Crop Improvement</i> , 2010 , 24, 392-399	1.4	17
32	Impact of brown stink bug (Heteroptera: Pentatomidae) feeding on corn grain yield components and quality. <i>Journal of Economic Entomology</i> , 2010 , 103, 2072-9	2.2	32

Using Nutrient Solutions to Trap the Almond Moth (Lepidoptera: Pyralidae) in a Peanut Shelling

Aphid (Hemiptera: Aphididae) resistance in wheat near-isogenic lines. Journal of Economic

and Storage Facility. Journal of Entomological Science, 2006, 41, 285-291

0.4

2.2

6

Entomology, **2004**, 97, 646-53

15

LIST OF PUBLICATIONS

13	Enzymatic chlorophyll degradation in wheat near-isogenic lines elicited by cereal aphid (Homoptera: Aphididae) feeding. <i>Journal of Economic Entomology</i> , 2004 , 97, 661-7	2.2	9
12	Possible roles of esterase, glutathione S-transferase, and superoxide dismutase activities in understanding aphidereal interactions. <i>Entomologia Experimentalis Et Applicata</i> , 2003 , 108, 187-195	2.1	22
11	Dynamic change in photosynthetic pigments and chlorophyll degradation elicited by cereal aphid feeding. <i>Entomologia Experimentalis Et Applicata</i> , 2002 , 105, 43-53	2.1	35
10	Comparison of Hindwing Hamuli from Five Species of Cereal Aphids (Hemiptera: Aphididae). <i>Annals of the Entomological Society of America</i> , 2002 , 95, 109-114	2	4
9	In vitro enzymatic chlorophyll catabolism in wheat elicited by cereal aphid feeding. <i>Entomologia Experimentalis Et Applicata</i> , 2001 , 101, 159-166	2.1	22
8	Oxidative responses of resistant and susceptible cereal leaves to symptomatic and nonsymptomatic cereal aphid (Hemiptera: Aphididae) feeding. <i>Journal of Economic Entomology</i> , 2001 , 94, 743-51	2.2	77
7	Comparison of DIMBOA concentrations among wheat isolines and corresponding plant introduction lines. <i>Entomologia Experimentalis Et Applicata</i> , 2000 , 96, 275-279	2.1	10
6	Hydrolase and Oxido-Reductase Activities in Diuraphis noxia and Rhopalosiphum padi (Hemiptera: Aphididae). <i>Annals of the Entomological Society of America</i> , 2000 , 93, 595-601	2	19
5	Identification and quantification of hydroxamic acids in maize seedling root tissue and impact on western corn rootworm (Coleoptera: Chrysomelidae) larval development. <i>Journal of Economic Entomology</i> , 2000 , 93, 989-92	2.2	15
4	Physiological and Growth Tolerance in Wheat to Russian Wheat Aphid (Homoptera: Aphididae) Injury. <i>Environmental Entomology</i> , 1999 , 28, 787-794	2.1	66
3	Influence of cereal leaf epicuticular wax on Diuraphis noxia probing behavior and nymphoposition. <i>Entomologia Experimentalis Et Applicata</i> , 1998 , 89, 111-118	2.1	9
2	Distribution of Russian Wheat Aphid (Homoptera: Aphididae) Salivary Sheaths in Resistant and Susceptible Wheat Leaves. <i>Journal of Economic Entomology</i> , 1997 , 90, 848-853	2.2	12
1	Effect of Wheat Leaf Epicuticular Structure on Host Selection and Probing Rhythm of Russian Wheat Aphid (Homoptera: Aphididae). <i>Journal of Economic Entomology</i> , 1997 , 90, 1400-1407	2.2	20