
Stefan Taubert

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/3560228/publications.pdf Version: 2024-02-01



#	Article	IF	CITATIONS
1	Nuclear hormone receptor NHR-49 acts in parallel with HIF-1 to promote hypoxia adaptation in Caenorhabditis elegans. ELife, 2022, 11, .	6.0	14
2	cSurvival: a web resource for biomarker interactions in cancer outcomes and in cell lines. Briefings in Bioinformatics, 2022, 23, .	6.5	7
3	Beyond Proteostasis: Lipid Metabolism as a New Player in ER Homeostasis. Metabolites, 2021, 11, 52.	2.9	30
4	Xenobiotic metabolism and transport in <i>Caenorhabditis elegans</i> . Journal of Toxicology and Environmental Health - Part B: Critical Reviews, 2021, 24, 51-94.	6.5	51
5	eVITTA: a web-based visualization and inference toolbox for transcriptome analysis. Nucleic Acids Research, 2021, 49, W207-W215.	14.5	45
6	NHR-49/PPAR-Î \pm and HLH-30/TFEB cooperate for C. elegans host defense via a flavin-containing monooxygenase. ELife, 2021, 10, .	6.0	37
7	Stress sensor Ire1 deploys a divergent transcriptional program in response to lipid bilayer stress. Journal of Cell Biology, 2020, 219, .	5.2	48
8	MDT-15/MED15 permits longevity at low temperature via enhancing lipidostasis and proteostasis. PLoS Biology, 2019, 17, e3000415.	5.6	51
9	Epigenetic regulator G9a provides glucose as a sweet key to stress resistance. PLoS Biology, 2019, 17, e3000236.	5.6	1
10	Mediator subunit MDT-15/MED15 and Nuclear Receptor HIZR-1/HNF4 cooperate to regulate toxic metal stress responses in Caenorhabditis elegans. PLoS Genetics, 2019, 15, e1008508.	3.5	20
11	<scp>NHR</scp> â€49/ <scp>HNF</scp> 4 integrates regulation of fatty acid metabolism with a protective transcriptional response to oxidative stress and fasting. Aging Cell, 2018, 17, e12743.	6.7	75
12	Activity of translation regulator eukaryotic elongation factor-2 kinase is increased in Parkinson disease brain and its inhibition reduces alpha synuclein toxicity. Acta Neuropathologica Communications, 2018, 6, 54.	5.2	48
13	Genomic and Cytogenetic Characterization of a Balanced Translocation Disrupting <i>NUP98</i> . Cytogenetic and Genome Research, 2017, 152, 117-121.	1.1	1
14	The R148.3 Gene Modulates <i>Caenorhabditis elegans</i> Lifespan and Fat Metabolism. G3: Genes, Genomes, Genetics, 2017, 7, 2739-2747.	1.8	5
15	eEF2K inhibition blocks Aβ42 neurotoxicity by promoting an NRF2 antioxidant response. Acta Neuropathologica, 2017, 133, 101-119.	7.7	48
16	Gain-of-Function Alleles in Caenorhabditis elegans Nuclear Hormone Receptor nhr-49 Are Functionally Distinct. PLoS ONE, 2016, 11, e0162708.	2.5	26
17	Caenorhabditis elegans Gets Metabolic Network Models. Cell Systems, 2016, 2, 293-294.	6.2	0
18	The Mediator Kinase Module Restrains Epidermal Growth Factor Receptor Signaling and Represses Vulval Cell Fate Specification in <i>Caenorhabditis elegans</i> . Genetics, 2016, 202, 583-599.	2.9	19

STEFAN TAUBERT

#	Article	IF	CITATIONS
19	The Mediator complex of Caenorhabditis elegans: insights into the developmental and physiological roles of a conserved transcriptional coregulator. Nucleic Acids Research, 2015, 43, 2442-2453.	14.5	39
20	s-Adenosylmethionine Levels Govern Innate Immunity through Distinct Methylation-Dependent Pathways. Cell Metabolism, 2015, 22, 633-645.	16.2	105
21	Activation of the endoplasmic reticulum unfolded protein response by lipid disequilibrium without disturbed proteostasis in vivo. Proceedings of the National Academy of Sciences of the United States of America, 2014, 111, E2271-80.	7.1	152
22	Membrane lipids and the endoplasmic reticulum unfolded protein response: An interesting relationship. Worm, 2014, 3, e962405.	1.0	8
23	The conserved Mediator subunit <scp>MDT</scp> â€15 is required for oxidative stress responses in <i><scp>C</scp>aenorhabditis elegans</i> . Aging Cell, 2014, 13, 70-79.	6.7	56
24	The C. elegans CDK8 Mediator module regulates axon guidance decisions in the ventral nerve cord and during dorsal axon navigation. Developmental Biology, 2013, 377, 385-398.	2.0	13
25	Conserved Mediator subunit MDTâ€15 assures metabolic homeostasis. FASEB Journal, 2013, 27, 822.14.	0.5	0
26	Repression of a Potassium Channel by Nuclear Hormone Receptor and TGF-β Signaling Modulates Insulin Signaling in Caenorhabditis elegans. PLoS Genetics, 2012, 8, e1002519.	3.5	16
27	Coordinate Regulation of Lipid Metabolism by Novel Nuclear Receptor Partnerships. PLoS Genetics, 2012, 8, e1002645.	3.5	86
28	Function and Regulation of Lipid Biology in Caenorhabditis elegans Aging. Frontiers in Physiology, 2012, 3, 143.	2.8	41
29	Somatic Differentiation and MR Imaging of Magnetically Labeled Human Embryonic Stem Cells. Cell Transplantation, 2012, 21, 2555-2567.	2.5	27
30	Nuclear hormone receptors in nematodes: Evolution and function. Molecular and Cellular Endocrinology, 2011, 334, 49-55.	3.2	84
31	Functional modularity of nuclear hormone receptors in a <i>Caenorhabditis elegans</i> metabolic gene regulatory network. Molecular Systems Biology, 2010, 6, 367.	7.2	93
32	SET(BP1)â€ing the stage for a better understanding of Schinzel—Giedion syndrome. Clinical Genetics, 2010, 78, 348-349.	2.0	2
33	The Mediator Subunit MDT-15 Confers Metabolic Adaptation to Ingested Material. PLoS Genetics, 2008, 4, e1000021.	3.5	100
34	Lifespan extension by conditions that inhibit translation in Caenorhabditis elegans. Aging Cell, 2007, 6, 95-110.	6.7	784
35	A Mediator subunit, MDT-15, integrates regulation of fatty acid metabolism by NHR-49-dependent and -independent pathways in C. elegans. Genes and Development, 2006, 20, 1137-1149.	5.9	220
36	E2F-Dependent Histone Acetylation and Recruitment of the Tip60 Acetyltransferase Complex to Chromatin in Late G 1. Molecular and Cellular Biology, 2004, 24, 4546-4556.	2.3	194

STEFAN TAUBERT

#	Article	IF	CITATIONS
37	MYC recruits the TIP60 histone acetyltransferase complex to chromatin. EMBO Reports, 2003, 4, 575-580.	4.5	331
38	Function of the c-Myc oncoprotein in chromatin remodeling and transcription. Biochimica Et Biophysica Acta: Reviews on Cancer, 2001, 1471, M135-M145.	7.4	102
39	Binding of c-Myc to chromatin mediates mitogen-induced acetylation of histone H4 and gene activation. Genes and Development, 2001, 15, 2069-2082.	5.9	441
40	Pleiotropic effects of cAMP on germination, antibiotic biosynthesis and morphological development inStreptomyces coelicolor. Molecular Microbiology, 1998, 30, 33-46.	2.5	115
41	Bacterial diet affects vulval organogenesis in Caenorhabditis elegans Mediator kinase module mutants . Matters, 0, , .	1.0	0