

Margaret K Mak

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3574347/publications.pdf>

Version: 2024-02-01

10
papers

581
citations

1464605

7
h-index

1637695

9
g-index

10
all docs

10
docs citations

10
times ranked

992
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|--|-----|-----------|
| 1 | Single session transcranial direct current stimulation to the primary motor cortex fails to enhance early motor sequence learning in Parkinson's disease. <i>Behavioural Brain Research</i> , 2022, 418, 113624. | 1.2 | 6 |
| 2 | Intelligent wearable system with accurate detection of abnormal gait and timely cueing for mobility enhancement of people with Parkinson's disease. <i>Wearable Technologies</i> , 2022, 3, . | 1.6 | 2 |
| 3 | The effect of transcranial direct current stimulation on upper limb motor performance in Parkinson's disease: a systematic review. <i>Journal of Neurology</i> , 2020, 267, 3479-3488. | 1.8 | 26 |
| 4 | Long-term effects of exercise and physical therapy in people with Parkinson disease. <i>Nature Reviews Neurology</i> , 2017, 13, 689-703. | 4.9 | 318 |
| 5 | The potential synergism by combining external counterpulsation with intermittent theta burst stimulation in post-stroke motor function recovery. <i>Medical Hypotheses</i> , 2016, 93, 140-142. | 0.8 | 0 |
| 6 | Task- and Context-Specific Balance Training Program Enhances Dynamic Balance and Functional Performance in Parkinsonian Nonfallers: A Randomized Controlled Trial With Six-Month Follow-Up. <i>Archives of Physical Medicine and Rehabilitation</i> , 2015, 96, 2103-2111. | 0.5 | 29 |
| 7 | Impaired Executive Function Can Predict Recurrent Falls in Parkinson's Disease. <i>Archives of Physical Medicine and Rehabilitation</i> , 2014, 95, 2390-2395. | 0.5 | 48 |
| 8 | Effect of cued training on motor evoked potential and cortical silent period in people with Parkinson's disease. <i>Clinical Neurophysiology</i> , 2013, 124, 545-550. | 0.7 | 10 |
| 9 | Validation of the Chinese Translated Activities-Specific Balance Confidence Scale. <i>Archives of Physical Medicine and Rehabilitation</i> , 2007, 88, 496-503. | 0.5 | 94 |
| 10 | Mediolateral sway in single-leg stance is the best discriminator of balance performance for Tai-Chi practitioners. <i>Archives of Physical Medicine and Rehabilitation</i> , 2003, 84, 683-686. | 0.5 | 48 |