

Ruth Schmidt-Ullrich

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3579536/publications.pdf>

Version: 2024-02-01

16
papers

1,508
citations

687363

13
h-index

940533

16
g-index

16
all docs

16
docs citations

16
times ranked

2695
citing authors

#	ARTICLE	IF	CITATIONS
1	Reciprocal Requirements for EDA/EDAR/NF- κ B and Wnt/ β 2-Catenin Signaling Pathways in Hair Follicle Induction. <i>Developmental Cell</i> , 2009, 17, 49-61.	7.0	310
2	Requirement of NF- κ B/Rel for the development of hair follicles and other epidermal appendices. <i>Development (Cambridge)</i> , 2001, 128, 3843-3853.	2.5	187
3	Requirement of Nuclear Factor- κ B in Angiotensin II- and Isoproterenol-Induced Cardiac Hypertrophy In Vivo. <i>Circulation</i> , 2005, 111, 2319-2325.	1.6	169
4	NF- κ B transmits Eda A1/EdaR signalling to activate Shh and cyclin D1 expression, and controls post-initiation hair placode down growth. <i>Development (Cambridge)</i> , 2006, 133, 1045-1057.	2.5	153
5	Proteasome-mediated degradation of I κ B \pm and processing of p105 in Crohn disease and ulcerative colitis. <i>Journal of Clinical Investigation</i> , 2006, 116, 3195-3203.	8.2	146
6	Tubular Epithelial NF- κ B Activity Regulates Ischemic AKI. <i>Journal of the American Society of Nephrology: JASN</i> , 2016, 27, 2658-2669.	6.1	138
7	Vascular Endothelial Cell-Specific NF- κ B Suppression Attenuates Hypertension-Induced Renal Damage. <i>Circulation Research</i> , 2007, 101, 268-276.	4.5	128
8	LHX2 is a direct NF- κ B target gene that promotes primary hair follicle placode down-growth. <i>Development (Cambridge)</i> , 2016, 143, 1512-22.	2.5	53
9	Central immune tolerance depends on crosstalk between the classical and alternative NF- κ B pathways in medullary thymic epithelial cells. <i>Journal of Autoimmunity</i> , 2017, 81, 56-67.	6.5	51
10	NF- κ B Activation Protects Oligodendrocytes against Inflammation. <i>Journal of Neuroscience</i> , 2017, 37, 9332-9344.	3.6	43
11	Canonical BMP signaling in tubular cells mediates recovery after acute kidney injury. <i>Kidney International</i> , 2019, 95, 108-122.	5.2	40
12	Transcriptional repression of <i>NFKBIA</i> triggers constitutive IKK and proteasome-independent p65/RelA activation in senescence. <i>EMBO Journal</i> , 2021, 40, e104296.	7.8	34
13	NF- κ B Participates in Mouse Hair Cycle Control and Plays Distinct Roles in the Various Pelage Hair Follicle Types. <i>Journal of Investigative Dermatology</i> , 2018, 138, 256-264.	0.7	23
14	Deficiency in <i>IκB\pm</i> in the intestinal epithelium leads to spontaneous inflammation and mediates apoptosis in the gut. <i>Journal of Pathology</i> , 2020, 251, 160-174.	4.5	14
15	NF- κ B Activity Is Required for Anagen Maintenance in Human Hair Follicles In Vitro. <i>Journal of Investigative Dermatology</i> , 2014, 134, 2036-2038.	0.7	12
16	NF- κ B determines Paneth versus goblet cell fate decision in the small intestine. <i>Development (Cambridge)</i> , 2021, 148, .	2.5	7