Mozhgan Dorkhan

List of Publications by Year in descending order

Source: https://exaly.com/author-pdf/3582167/publications.pdf

Version: 2024-02-01

19 papers 4,716 citations

12 h-index 18 g-index

20 all docs

20 docs citations

times ranked

20

11322 citing authors

#	Article	IF	CITATIONS
1	Large-scale association analysis provides insights into the genetic architecture and pathophysiology of type 2 diabetes. Nature Genetics, 2012, 44, 981-990.	9.4	1,748
2	Novel subgroups of adult-onset diabetes and their association with outcomes: a data-driven cluster analysis of six variables. Lancet Diabetes and Endocrinology,the, 2018, 6, 361-369.	5 . 5	1,430
3	Genome-wide trans-ancestry meta-analysis provides insight into the genetic architecture of type 2 diabetes susceptibility. Nature Genetics, 2014, 46, 234-244.	9.4	959
4	Genetic fine mapping and genomic annotation defines causal mechanisms at type 2 diabetes susceptibility loci. Nature Genetics, 2015, 47, 1415-1425.	9.4	365
5	Sweetened beverage intake and risk of latent autoimmune diabetes in adults (LADA) and type 2 diabetes. European Journal of Endocrinology, 2016, 175, 605-614.	1.9	35
6	Treatment with a thiazolidinedione increases eye protrusion in a subgroup of patients with type 2 diabetes. Clinical Endocrinology, 2006, 65, 35-39.	1.2	33
7	Prevalence and risk factors for diabetic retinopathy at diagnosis (DRAD) in patients recently diagnosed with type 2 diabetes (T2D) or latent autoimmune diabetes in the adult (LADA). Journal of Diabetes and Its Complications, 2016, 30, 1456-1461.	1.2	26
8	Effect of pioglitazone versus insulin glargine on cardiac size, function, and measures of fluid retention in patients with type 2 diabetes. Cardiovascular Diabetology, 2009, 8, 15.	2.7	24
9	Differences in effects of insulin glargine or pioglitazone added to oral anti-diabetic therapy in patients with type 2 diabetes. Diabetes Research and Clinical Practice, 2008, 82, 340-345.	1.1	18
10	Alcohol and the risk for latent autoimmune diabetes in adults: results based on Swedish ESTRID study. European Journal of Endocrinology, 2014, 171, 535-543.	1.9	17
11	Age at first childbirth and oral contraceptive use are associated with risk of androgen receptor-negative breast cancer: the Malmö Diet and Cancer Cohort. Cancer Causes and Control, 2014, 25, 945-957.	0.8	17
12	Gender differences in non-glycemic responses to improved insulin sensitivity by pioglitazone treatment in patients with type 2 diabetes. Acta Diabetologica, 2014, 51, 185-192.	1.2	12
13	Clinical Effectiveness of Liraglutide vs Sitagliptin on Glycemic Control and Body Weight in Patients with Type 2 Diabetes: A Retrospective Assessment in Sweden. Diabetes Therapy, 2016, 7, 321-333.	1.2	7
14	Education, immigration and income as risk factors for hemoglobin A1c >70 mmol/mol when diagnosed with type 2 diabetes or latent autoimmune diabetes in adult: a population-based cohort study. BMJ Open Diabetes Research and Care, 2017, 5, e000346.	1.2	6
15	Genotypes of HLA, TCF7L2, and FTO as potential modifiers of the association between sweetened beverage consumption and risk of LADA and type 2 diabetes. European Journal of Nutrition, 2020, 59, 127-135.	1.8	6
16	Long-term exposure to insulin and volumetric mammographic density: observational and genetic associations in the Karma study. Breast Cancer Research, 2018, 20, 93.	2.2	5
17	Treatment with pioglitazone induced significant, reversible mitral regurgitation. Cardiovascular Diabetology, 2008, 7, 12.	2.7	4
18	A review of pioglitazone HCL and glimepiride in the treatment of type 2 diabetes. Vascular Health and Risk Management, 2007, 3, 721-31.	1.0	4

ARTICLE

Abstract A126: Oral contraceptives and late first childbirth increase the risk of androgen receptor-negative breast cancer: The Malmö Diet and Cancer cohort., 2013,,...

O

CITATIONS