

# Seika Kamohara

## List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3584318/publications.pdf>

Version: 2024-02-01

21  
papers

838  
citations

687220

13  
h-index

713332

21  
g-index

21  
all docs

21  
docs citations

21  
times ranked

1636  
citing authors

#	ARTICLE	IF	CITATIONS
1	Variations in the FTO gene are associated with severe obesity in the Japanese. <i>Journal of Human Genetics</i> , 2008, 53, 546-553.	1.1	219
2	Association between obesity and polymorphisms in SEC16B, TMEM18, GNPDA2, BDNF, FAIM2 and MC4R in a Japanese population. <i>Journal of Human Genetics</i> , 2009, 54, 727-731.	1.1	115
3	Oxidative DNA damage (8-hydroxydeoxyguanosine) and body iron status: a study on 2507 healthy people. <i>Free Radical Biology and Medicine</i> , 2003, 35, 826-832.	1.3	89
4	Association of variations in the FTO, SCG3 and MTMR9 genes with metabolic syndrome in a Japanese population. <i>Journal of Human Genetics</i> , 2011, 56, 647-651.	1.1	69
5	Association of single-nucleotide polymorphisms in MTMR9 gene with obesity. <i>Human Molecular Genetics</i> , 2007, 16, 3017-3026.	1.4	51
6	INSIG2 gene rs7566605 polymorphism is associated with severe obesity in Japanese. <i>Journal of Human Genetics</i> , 2008, 53, 857-862.	1.1	43
7	Functional Single-Nucleotide Polymorphisms in the Secretogranin III (SCG3) Gene that Form Secretory Granules with Appetite-Related Neuropeptides Are Associated with Obesity. <i>Journal of Clinical Endocrinology and Metabolism</i> , 2007, 92, 1145-1154.	1.8	40
8	Genetic variations in the CYP17A1 and NT5C2 genes are associated with a reduction in visceral and subcutaneous fat areas in Japanese women. <i>Journal of Human Genetics</i> , 2012, 57, 46-51.	1.1	38
9	Polymorphisms in NRXN3, TFAP2B, MSRA, LYPLAL1, FTO and MC4R and their effect on visceral fat area in the Japanese population. <i>Journal of Human Genetics</i> , 2010, 55, 738-742.	1.1	36
10	Computed tomography analysis of the association between the SH2B1 rs7498665 single-nucleotide polymorphism and visceral fat area. <i>Journal of Human Genetics</i> , 2011, 56, 716-719.	1.1	24
11	Association between type 2 diabetes genetic susceptibility loci and visceral and subcutaneous fat area as determined by computed tomography. <i>Journal of Human Genetics</i> , 2012, 57, 305-310.	1.1	23
12	&lt;i>rs206936</i> is associated with body mass index in obese Japanese women. <i>Endocrine Journal</i> , 2013, 60, 991-1000.	0.7	16
13	Replication Study of 15 Recently Published Loci for Body Fat Distribution in the Japanese Population. <i>Journal of Atherosclerosis and Thrombosis</i> , 2013, 20, 336-350.	0.9	16
14	An evidence-based review: Anti-obesity effects of <i>Coleus forskohlii</i> . <i>Personalized Medicine Universe</i> , 2016, 5, 16-20.	0.1	11
15	A New Japanese Vegetarian Food Guide. <i>Asia-Pacific Journal of Public Health</i> , 2009, 21, 160-169.	0.4	10
16	A <i>Coleus forskohlii</i> extract improves body composition in healthy volunteers: An open-label trial. <i>Personalized Medicine Universe</i> , 2013, 2, 25-27.	0.1	10
17	Bridging conventional medicine and complementary and alternative medicine. <i>IEEE Engineering in Medicine and Biology Magazine</i> , 2005, 24, 30-34.	1.1	8
18	Screening of 336 single-nucleotide polymorphisms in 85 obesity-related genes revealed McKusick's "Kaufman syndrome gene variants are associated with metabolic syndrome. <i>Journal of Human Genetics</i> , 2009, 54, 230-235.	1.1	7

#	ARTICLE	IF	CITATIONS
19	Safety of a <i>Coleus forskohlii</i> formulation in healthy volunteers. <i>Personalized Medicine Universe</i> , 2015, 4, 63-65.	0.1	7
20	<i></i>ADIPOQ</i> polymorphisms are associated with insulin resistance in Japanese women. <i>Endocrine Journal</i> , 2015, 62, 513-521.	0.7	5
21	ANTIOXIDANT SUPPLEMENTATION DECREASES THE AMOUNT OF URINARY 8-OHdG EXCRETION INDUCED BY A SINGLE BOUT OF EXERCISE. <i>Japanese Journal of Physical Fitness and Sports Medicine</i> , 2006, 55, S251-S256.	0.0	1