Martin Brunovsky

List of Publications by Citations

Source: https://exaly.com/author-pdf/3584514/martin-brunovsky-publications-by-citations.pdf

Version: 2024-04-19

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

56
papers

1,239
citations

20
h-index
g-index

89
ext. papers

20
34
g-index

3.8
L-index

#	Paper	IF	Citations
56	Early reduction in prefrontal theta QEEG cordance value predicts response to venlafaxine treatment in patients with resistant depressive disorder. <i>European Psychiatry</i> , 2008 , 23, 350-5	6	101
55	Changes in QEEG prefrontal cordance as a predictor of response to antidepressants in patients with treatment resistant depressive disorder: a pilot study. <i>Journal of Psychiatric Research</i> , 2007 , 41, 319-25	5.2	88
54	Guidelines for the recording and evaluation of pharmaco-EEG data in man: the International Pharmaco-EEG Society (IPEG). <i>Neuropsychobiology</i> , 2012 , 66, 201-20	4	84
53	Effect of low-frequency rTMS on electromagnetic tomography (LORETA) and regional brain metabolism (PET) in schizophrenia patients with auditory hallucinations. <i>Neuropsychobiology</i> , 2007 , 55, 132-42	4	78
52	The change of prefrontal QEEG theta cordance as a predictor of response to bupropion treatment in patients who had failed to respond to previous antidepressant treatments. <i>European Neuropsychopharmacology</i> , 2010 , 20, 459-66	1.2	67
51	Electroencephalographic spectral and coherence analysis of ketamine in rats: correlation with behavioral effects and pharmacokinetics. <i>Neuropsychobiology</i> , 2011 , 63, 202-18	4	60
50	Latent toxoplasmosis reduces gray matter density in schizophrenia but not in controls: voxel-based-morphometry (VBM) study. <i>World Journal of Biological Psychiatry</i> , 2012 , 13, 501-9	3.8	58
49	Objective assessment of the degree of dementia by means of EEG. <i>Neuropsychobiology</i> , 2003 , 48, 19-26	4	56
48	EEG source analysis in obsessive-compulsive disorder. <i>Clinical Neurophysiology</i> , 2011 , 122, 1735-43	4.3	46
47	Low frequency (1-Hz), right prefrontal repetitive transcranial magnetic stimulation (rTMS) compared with venlafaxine ER in the treatment of resistant depression: a double-blind, single-centre, randomized study. <i>Journal of Affective Disorders</i> , 2009 , 118, 94-100	6.6	40
46	Sleep disturbances in patients treated for panic disorder. <i>Sleep Medicine</i> , 2005 , 6, 149-53	4.6	34
45	Guidelines for the recording and evaluation of pharmaco-sleep studies in man: the International Pharmaco-EEG Society (IPEG). <i>Neuropsychobiology</i> , 2013 , 67, 127-67	4	33
44	Subanesthetic dose of ketamine decreases prefrontal theta cordance in healthy volunteers: implications for antidepressant effect. <i>Psychological Medicine</i> , 2010 , 40, 1443-51	6.9	33
43	Prediction of treatment response and the effect of independent component neurofeedback in obsessive-compulsive disorder: a randomized, sham-controlled, double-blind study. <i>Neuropsychobiology</i> , 2013 , 67, 210-23	4	31
42	Behavioral, neurochemical and pharmaco-EEG profiles of the psychedelic drug 4-bromo-2,5-dimethoxyphenethylamine (2C-B) in rats. <i>Psychopharmacology</i> , 2013 , 225, 75-93	4.7	31
41	Standardized low-resolution brain electromagnetic tomography (sLORETA) in the prediction of response to cholinesterase inhibitors in patients with Alzheimer disease 2008, 7,		29
40	QEEG Theta Cordance in the Prediction of Treatment Outcome to Prefrontal Repetitive Transcranial Magnetic Stimulation or Venlafaxine ER in Patients With Major Depressive Disorder. <i>Clinical EEG and Neuroscience</i> , 2015 , 46, 73-80	2.3	26

39	LORETA functional imaging in antipsychotic-naive and olanzapine-, clozapine- and risperidone-treated patients with schizophrenia. <i>Neuropsychobiology</i> , 2008 , 58, 1-10	4	26
38	Subjective and objective evaluation of alertness and sleep quality in depressed patients. <i>BMC Psychiatry</i> , 2004 , 4, 14	4.2	26
37	The effectiveness of prefrontal theta cordance and early reduction of depressive symptoms in the prediction of antidepressant treatment outcome in patients with resistant depression: analysis of naturalistic data. <i>European Archives of Psychiatry and Clinical Neuroscience</i> , 2015 , 265, 73-82	5.1	21
36	The change of QEEG prefrontal cordance as a response predictor to antidepressive intervention in bipolar depression. A pilot study. <i>Journal of Psychiatric Research</i> , 2012 , 46, 219-25	5.2	20
35	Discontinuation of hypnotics during cognitive behavioural therapy for insomnia. <i>BMC Psychiatry</i> , 2008 , 8, 80	4.2	17
34	Augmentation of antidepressants with bright light therapy in patients with comorbid depression and borderline personality disorder. <i>Biomedical Papers of the Medical Faculty of the University Palacky&#x0301;, Olomouc, Czechoslovakia,</i> 2010 , 154, 355-61	1.7	16
33	Electrophysiological correlates of bilateral and unilateral repetitive transcranial magnetic stimulation in patients with bipolar depression. <i>Psychiatry Research</i> , 2016 , 240, 364-375	9.9	16
32	Psilocybin disrupts sensory and higher order cognitive processing but not pre-attentive cognitive processing-study on P300 and mismatch negativity in healthy volunteers. <i>Psychopharmacology</i> , 2018 , 235, 491-503	4.7	15
31	Role of Glutamatergic System in Obsessive-Compulsive Disorder with Possible Therapeutic Implications. <i>Pharmacopsychiatry</i> , 2018 , 51, 229-242	2	15
30	Physical Comorbidities in Depression Co-Occurring with Anxiety: A Cross Sectional Study in the Czech Primary Care System. <i>International Journal of Environmental Research and Public Health</i> , 2015 , 12, 15728-38	4.6	15
29	Trazodone improves the results of cognitive behaviour therapy of primary insomnia in non-depressed patients. <i>Neuroendocrinology Letters</i> , 2008 , 29, 895-901	0.3	15
28	Sad mood induction has an opposite effect on amygdala response to emotional stimuli in euthymic patients with bipolar disorder and healthy controls. <i>Journal of Psychiatry and Neuroscience</i> , 2015 , 40, 134-42	4.5	12
27	Early stages of pediatric bipolar disorder: retrospective analysis of a Czech inpatient sample. <i>Neuropsychiatric Disease and Treatment</i> , 2015 , 11, 2855-64	3.1	11
26	Standardized low-resolution electromagnetic tomography in obsessive-compulsive disordera replication study. <i>Neuroscience Letters</i> , 2013 , 548, 185-9	3.3	10
25	EEG abnormalities in dementia reflect the parietal lobe syndrome. <i>Clinical Neurophysiology</i> , 2001 , 112, 1001-5	4.3	10
24	Cognitive deficits in patients with obsessive-compulsive disorder - electroencephalography correlates. <i>Neuropsychiatric Disease and Treatment</i> , 2016 , 12, 1119-25	3.1	10
23	The Comparison of Effectiveness of Various Potential Predictors of Response to Treatment With SSRIs in Patients With Depressive Disorder. <i>Journal of Nervous and Mental Disease</i> , 2017 , 205, 618-626	1.8	9
22	The effect of ((-)-2-oxa-4-aminobicyclo[3.1.0]hexane-2,6-dicarboxylic acid (LY379268), an mGlu2/3 receptor agonist, on EEG power spectra and coherence in ketamine model of psychosis. <i>Pharmacology Biochemistry and Behavior</i> , 2014 , 122, 212-21	3.9	9

21	The Antidepressant Effect of Ketamine Is Dampened by Concomitant Benzodiazepine Medication. <i>Frontiers in Psychiatry</i> , 2020 , 11, 844	5	9
20	Early change of prefrontal theta cordance and occipital alpha asymmetry in the prediction of responses to antidepressants. <i>International Journal of Psychophysiology</i> , 2019 , 143, 1-8	2.9	7
19	EEG correlates of induced anxiety in obsessive-compulsive patients: comparison of autobiographical and general anxiety scenarios. <i>Neuropsychiatric Disease and Treatment</i> , 2018 , 14, 2165	- 2:1 74	7
18	Objective measurement of the alertness level in dementia. <i>Dementia and Geriatric Cognitive Disorders</i> , 2003 , 15, 212-7	2.6	6
17	Transcranial Direct-Current Stimulation (tDCS) Versus Venlafaxine ER In The Treatment Of Depression: A Randomized, Double-Blind, Single-Center Study With Open-Label, Follow-Up. <i>Neuropsychiatric Disease and Treatment</i> , 2019 , 15, 3003-3014	3.1	4
16	Latent Schizencephaly With Psychotic Phenotype or Schizophrenia With Schizencephaly? A Case Report and Review of the Literature. <i>Clinical EEG and Neuroscience</i> , 2019 , 50, 13-19	2.3	3
15	Frames of reference and their neural correlates within navigation in a 3D environment. <i>Visual Neuroscience</i> , 2012 , 29, 183-91	1.7	3
14	Predicting Sex From EEG: Validity and Generalizability of Deep-Learning-Based Interpretable Classifier. <i>Frontiers in Neuroscience</i> , 2020 , 14, 589303	5.1	3
13	The Effects of Daytime Psilocybin Administration on Sleep: Implications for Antidepressant Action. <i>Frontiers in Pharmacology</i> , 2020 , 11, 602590	5.6	3
12	Artifacts in Simultaneous hdEEG/fMRI Imaging: A Nonlinear Dimensionality Reduction Approach. <i>Sensors</i> , 2019 , 19,	3.8	3
11	Baseline Difference in Quantitative Electroencephalography Variables Between Responders and Non-Responders to Low-Frequency Repetitive Transcranial Magnetic Stimulation in Depression. <i>Frontiers in Psychiatry</i> , 2020 , 11, 83	5	2
10	Predictive value of heart rate in treatment of major depression with ketamine in two controlled trials. <i>Clinical Neurophysiology</i> , 2021 , 132, 1339-1346	4.3	2
9	Electrical Source Imaging in Rats: Cortical EEG Performance and Limitations 2018,		1
8	Is short-term memory capacity ($7\square 2$) really predicted by theta to gamma cycle length ratio?. <i>Behavioural Brain Research</i> , 2021 , 414, 113465	3.4	1
7	Nonlinear Dimensionality Reduction and Feature Analysis for Artifact Component Identification in hdEEG Datasets. <i>IFMBE Proceedings</i> , 2019 , 415-419	0.2	
6	P.1.b.009 Default mode network activity decrement interrelated with anxious-like and dreadful experiences during acute cannabis intoxication. <i>European Neuropsychopharmacology</i> , 2014 , 24, S176-S17	77 ²	
5	Prefrontal Theta Cordance in the Prediction of Antidepressant Response to Various Classes of Antidepressants in Patients with Depressive Disorder. <i>European Psychiatry</i> , 2017 , 41, s140-s140	6	
4	Different impact of t.gondii seropositivity on the brain morphometry in schizophrenia and healthy control groups. <i>European Psychiatry</i> , 2011 , 26, 2124-2124	6	

LIST OF PUBLICATIONS

- P.3.020 How ketamine changes the neurophysiology of depressive patients Tbrains: a randomised controlled trial. *European Neuropsychopharmacology*, **2011**, 21, S75
- P.1.c.067 Effect of group II metabotropic glutamate receptor agonist on behaviour and electroencephalography in a model of psychosis. *European Neuropsychopharmacology*, **2010**, 20, S273-S274
- The effectiveness of various potential predictors of response to treatment with SSRIs in patients with depressive disorder. *European Psychiatry*, **2016**, 33, S406-S406