

Audny Anke

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3588418/publications.pdf>

Version: 2024-02-01

42
papers

2,458
citations

471509

17
h-index

265206

42
g-index

47
all docs

47
docs citations

47
times ranked

3514
citing authors

#	ARTICLE	IF	CITATIONS
1	Traumatic brain injury: integrated approaches to improve prevention, clinical care, and research. <i>Lancet Neurology</i> , The, 2017, 16, 987-1048.	10.2	1,571
2	Incidence of Hospital-Admitted Severe Traumatic Brain Injury and In-Hospital Fatality in Norway: A National Cohort Study. <i>Neuroepidemiology</i> , 2012, 38, 259-267.	2.3	87
3	Efficacy of antibiotic treatment in patients with chronic low back pain and Modic changes (the AIM) Tj ETQq1 1 0.784314 rgBT /Overl 6.0 77	6.0	77
4	Factors Affecting Caregiver Burden 1 Year After Severe Traumatic Brain Injury. <i>Journal of Head Trauma Rehabilitation</i> , 2015, 30, 411-423.	1.7	70
5	Alterations in cognitive outcome between 3 and 12 months in survivors of out-of-hospital cardiac arrest. <i>Resuscitation</i> , 2016, 105, 92-99.	3.0	47
6	Patterns of change and stability in caregiver burden and life satisfaction from 1 to 2 years after severe traumatic brain injury: A Norwegian longitudinal study. <i>NeuroRehabilitation</i> , 2017, 40, 211-222.	1.3	42
7	Functional Recovery and Life Satisfaction in the First Year After Severe Traumatic Brain Injury. <i>Journal of Head Trauma Rehabilitation</i> , 2015, 30, E38-E49.	1.7	38
8	A meta-analysis of constraint-induced movement therapy after stroke. <i>Journal of Rehabilitation Medicine</i> , 2014, 46, 833-842.	1.1	37
9	Family needs after brain injury: A cross cultural study. <i>NeuroRehabilitation</i> , 2015, 36, 203-214.	1.3	36
10	Determinants of cognitive outcome in survivors of out-of-hospital cardiac arrest. <i>Resuscitation</i> , 2014, 85, 1462-1468.	3.0	34
11	Experiences of quality of life the first year after stroke in Denmark and Norway. A qualitative analysis. <i>International Journal of Qualitative Studies on Health and Well-being</i> , 2019, 14, 1659540.	1.6	34
12	Efficacy of Constraint-Induced Movement Therapy in Early Stroke Rehabilitation. <i>Neurorehabilitation and Neural Repair</i> , 2015, 29, 517-525.	2.9	33
13	Return to work after severe traumatic brain injury: a national study with a one-year follow-up of neurocognitive and behavioural outcomes. <i>Neuropsychological Rehabilitation</i> , 2020, 30, 281-297.	1.6	29
14	Severe traumatic brain injury in Norway: Impact of age on outcome. <i>Journal of Rehabilitation Medicine</i> , 2013, 45, 734-740.	1.1	25
15	Mortality and One-Year Functional Outcome in Elderly and Very Old Patients with Severe Traumatic Brain Injuries: Observed and Predicted. <i>Behavioural Neurology</i> , 2015, 2015, 1-7.	2.1	23
16	Olfactory identification and its relationship to executive functions, memory, and disability one year after severe traumatic brain injury. <i>Neuropsychology</i> , 2016, 30, 98-108.	1.3	23
17	Antibiotic treatment In patients with chronic low back pain and Modic changes (the AIM study): study protocol for a randomised controlled trial. <i>Trials</i> , 2017, 18, 596.	1.6	21
18	Stroke-Specific Quality of Life one-year post-stroke in two Scandinavian country-regions with different organisation of rehabilitation services: a prospective study. <i>Disability and Rehabilitation</i> , 2021, 43, 3810-3820.	1.8	19

#	ARTICLE	IF	CITATIONS
19	Validity, reliability and Norwegian adaptation of the Stroke-Specific Quality of Life (SS-QOL) scale. SAGE Open Medicine, 2018, 6, 205031211775203.	1.8	18
20	Family members and health care workers' perspectives on motivational factors of participation in physical activity for people with intellectual disability: A qualitative study. Journal of Intellectual Disability Research, 2020, 64, 259-270.	2.0	17
21	Life satisfaction in subjects with long-term musculoskeletal pain in relation to pain intensity, pain distribution and coping. Journal of Rehabilitation Medicine, 2013, 45, 277-285.	1.1	15
22	Memory performance, global cerebral volumes and hippocampal subfield volumes in long-term survivors of Out-of-Hospital Cardiac Arrest. Resuscitation, 2018, 126, 21-28.	3.0	15
23	A quality indicator set for use in rehabilitation team care of people with rheumatic and musculoskeletal diseases; development and pilot testing. BMC Health Services Research, 2019, 19, 265.	2.2	15
24	Rehabilitation pathways and functional independence one year after severe traumatic brain injury. European Journal of Physical and Rehabilitation Medicine, 2016, 52, 650-661.	2.2	15
25	Norwegian constraint-induced therapy multisite trial: Adherence to treatment protocol applied early after stroke. Journal of Rehabilitation Medicine, 2015, 47, 816-823.	1.1	13
26	Family members'™ experience with in-hospital health care after severe traumatic brain injury: a national multicentre study. BMC Health Services Research, 2018, 18, 951.	2.2	13
27	Early versus late-applied constraint-induced movement therapy: A multisite, randomized controlled trial with a 12-month follow-up. Physiotherapy Research International, 2018, 23, e1689.	1.5	11
28	Family needs at one and two years after severe traumatic brain injury: a prospective study of changes and predictors. Brain Injury, 2020, 34, 89-97.	1.2	11
29	Physical Activity With Tailored mHealth Support for Individuals With Intellectual Disabilities: Protocol for a Randomized Controlled Trial. JMIR Research Protocols, 2020, 9, e19213.	1.0	11
30	Clinical effect modifiers of antibiotic treatment in patients with chronic low back pain and Modic changes - secondary analyses of a randomised, placebo-controlled trial (the AIM study). BMC Musculoskeletal Disorders, 2020, 21, 458.	1.9	9
31	How do multimorbidity and lifestyle factors impact the perceived health of adults with intellectual disabilities?. Journal of Intellectual Disability Research, 2021, 65, 772-783.	2.0	9
32	Can the health related quality of life measure QOLIBRI- overall scale (OS) be of use after stroke? A validation study. BMC Neurology, 2018, 18, 98.	1.8	6
33	Cortical Thickness and Cognitive Performance After Out-of-Hospital Cardiac Arrest. Neurorehabilitation and Neural Repair, 2019, 33, 296-306.	2.9	5
34	Rehabilitation Needs, Service Provision, and Costs in the First Year Following Traumatic Injuries: Protocol for a Prospective Cohort Study. JMIR Research Protocols, 2021, 10, e25980.	1.0	5
35	eHealth Approach for Motivating Physical Activities of People with Intellectual Disabilities. IFIP Advances in Information and Communication Technology, 2020, , 31-41.	0.7	5
36	Factors associated with non-completion of and scores on physical capability tests in health surveys: The North Health in Intellectual Disability Study. Journal of Applied Research in Intellectual Disabilities, 2021, , .	2.0	4

#	ARTICLE	IF	CITATIONS
37	mHealth Support to Stimulate Physical Activity in Individuals With Intellectual Disability: Protocol for a Mixed Methods Pilot Study. <i>JMIR Research Protocols</i> , 2022, 11, e37849.	1.0	4
38	The Family Needs Questionnaire-Revised: a Rasch analysis of measurement properties in the chronic phase after traumatic brain injury. <i>Brain Injury</i> , 2020, 34, 1375-1383.	1.2	3
39	Factors associated with met and unmet rehabilitation needs after stroke: A multicentre cohort study in Denmark and Norway. <i>Journal of Rehabilitation Medicine</i> , 2021, .	1.1	3
40	Motivating Physical Activity for Individuals with Intellectual Disability through Indoor Bike Cycling and Exergaming. <i>International Journal of Environmental Research and Public Health</i> , 2022, 19, 2914.	2.6	3
41	The family experiences of in-hospital care questionnaire in severe traumatic brain injury (FECQ-TBI): a validation study. <i>BMC Health Services Research</i> , 2016, 16, 675.	2.2	1
42	Persistent pain associated with socioeconomic and personal factors in a Sami and Non-Sami population in Norway: an analysis of SAMINOR 2 survey data. <i>International Journal of Circumpolar Health</i> , 2020, 79, 1787022.	1.2	1