

Alexander Kiani

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3589924/publications.pdf>

Version: 2024-02-01

64
papers

4,359
citations

257357

24
h-index

161767

54
g-index

65
all docs

65
docs citations

65
times ranked

6159
citing authors

#	ARTICLE	IF	CITATIONS
1	FOLFIRI plus cetuximab versus FOLFIRI plus bevacizumab as first-line treatment for patients with metastatic colorectal cancer (FIRE-3): a randomised, open-label, phase 3 trial. <i>Lancet Oncology</i> , The, 2014, 15, 1065-1075.	5.1	1,479
2	Addition of sorafenib versus placebo to standard therapy in patients aged 60 years or younger with newly diagnosed acute myeloid leukaemia (SORAML): a multicentre, phase 2, randomised controlled trial. <i>Lancet Oncology</i> , The, 2015, 16, 1691-1699.	5.1	347
3	FOLFIRI plus cetuximab versus FOLFIRI plus bevacizumab for metastatic colorectal cancer (FIRE-3): a post-hoc analysis of tumour dynamics in the final RAS wild-type subgroup of this randomised open-label phase 3 trial. <i>Lancet Oncology</i> , The, 2016, 17, 1426-1434.	5.1	336
4	Consensus Conference on Clinical Practice in Chronic GVHD: Second-Line Treatment of Chronic Graft-versus-Host Disease. <i>Biology of Blood and Marrow Transplantation</i> , 2011, 17, 1-17.	2.0	311
5	Manipulating Immune Responses with Immunosuppressive Agents that Target NFAT. <i>Immunity</i> , 2000, 12, 359-372.	6.6	267
6	Consensus Conference on Clinical Practice in Chronic Graft-versus-Host Disease (GVHD): First-Line and Topical Treatment of Chronic GVHD. <i>Biology of Blood and Marrow Transplantation</i> , 2010, 16, 1611-1628.	2.0	226
7	Down-Regulation of IL-4 Gene Transcription and Control of Th2 Cell Differentiation by a Mechanism Involving NFAT1. <i>Immunity</i> , 1997, 7, 849-860.	6.6	161
8	Regulation of interferon- γ gene expression by nuclear factor of activated T cells. <i>Blood</i> , 2001, 98, 1480-1488.	0.6	116
9	Impact of Subsequent Therapies on Outcome of the FIRE-3/AIO KRK0306 Trial: First-Line Therapy With FOLFIRI Plus Cetuximab or Bevacizumab in Patients With <i>KRAS</i> Wild-Type Tumors in Metastatic Colorectal Cancer. <i>Journal of Clinical Oncology</i> , 2015, 33, 3718-3726.	0.8	112
10	Monitoring of donor chimerism in sorted CD34+ peripheral blood cells allows the sensitive detection of imminent relapse after allogeneic stem cell transplantation. <i>Haematologica</i> , 2009, 94, 1613-1617.	1.7	98
11	Does time from diagnosis to treatment affect the prognosis of patients with newly diagnosed acute myeloid leukemia?. <i>Blood</i> , 2020, 136, 823-830.	0.6	85
12	FOLFIRI plus cetuximab or bevacizumab for advanced colorectal cancer: final survival and per-protocol analysis of FIRE-3, a randomised clinical trial. <i>British Journal of Cancer</i> , 2021, 124, 587-594.	2.9	79
13	Minimal residual disease-directed preemptive treatment with azacitidine in patients with NPM1-mutant acute myeloid leukemia and molecular relapse. <i>Haematologica</i> , 2011, 96, 1568-1570.	1.7	67
14	Prophylactic transfer of BCR-ABL ⁻ , PR1 ⁻ , and WT1-reactive donor T cells after T cell-depleted allogeneic hematopoietic cell transplantation in patients with chronic myeloid leukemia. <i>Blood</i> , 2011, 117, 7174-7184.	0.6	48
15	Pharmacokinetics of gemcitabine in a patient with end-stage renal disease: effective clearance of its main metabolite by standard hemodialysis treatment. <i>Cancer Chemotherapy and Pharmacology</i> , 2003, 51, 266-270.	1.1	46
16	Cidofovir and Foscarnet for Treatment of Human Herpesvirus 6 Encephalitis in a Neutropenic Stem Cell Transplant Recipient. <i>Clinical Infectious Diseases</i> , 2007, 44, e118-e120.	2.9	46
17	Expression and regulation of NFAT (nuclear factors of activated T cells) in human CD34+ cells: down-regulation upon myeloid differentiation. <i>Journal of Leukocyte Biology</i> , 2004, 76, 1057-1065.	1.5	43
18	Minimal Residual Disease (MRD) Based Preemptive 5-azacytidine Treatment Can Prevent or Delay Imminent Relapse In Patients with High-Risk MDS or AML After Allogeneic HSCT – Results of the RELAZA Trial. <i>Blood</i> , 2010, 116, 679-679.	0.6	41

#	ARTICLE	IF	CITATIONS
19	Exploring the effect of primary tumor sidedness on therapeutic efficacy across treatment lines in patients with metastatic colorectal cancer: analysis of FIRE-3 (AIOKRK0306). <i>Oncotarget</i> , 2017, 8, 105749-105760.	0.8	41
20	Sorafenib or placebo in patients with newly diagnosed acute myeloid leukaemia: long-term follow-up of the randomized controlled SORAML trial. <i>Leukemia</i> , 2021, 35, 2517-2525.	3.3	40
21	T cell-mediated graft-versus-leukemia reactions after allogeneic stem cell transplantation. <i>Cancer Immunology, Immunotherapy</i> , 2005, 54, 1043-1058.	2.0	28
22	Velcade, Intravenous Cyclophosphamide and Dexamethasone (VCD) Induction for Previously Untreated Multiple Myeloma (German DSMM XIa Trial).. <i>Blood</i> , 2009, 114, 131-131.	0.6	27
23	Expression analysis of nuclear factor of activated T cells (NFAT) during myeloid differentiation of CD34+ cells: regulation of Fas ligand gene expression in megakaryocytes. <i>Experimental Hematology</i> , 2007, 35, 757-770.	0.2	26
24	Gemtuzumab Ozogamicin as Part of Reduced-Intensity Conditioning for Allogeneic Hematopoietic Cell Transplantation in Patients with Relapsed Acute Myeloid Leukemia. <i>Clinical Cancer Research</i> , 2008, 14, 5585-5593.	3.2	26
25	Ponatinib in the Treatment of Chronic Myeloid Leukemia and Philadelphia Chromosome-Positive Acute Leukemia: Recommendations of a German Expert Consensus Panel with Focus on Cardiovascular Management. <i>Acta Haematologica</i> , 2020, 143, 217-231.	0.7	26
26	¹⁸⁸ Re anti-CD66 radioimmunotherapy combined with reduced-intensity conditioning and <i>in vivo</i> T cell depletion in elderly patients undergoing allogeneic haematopoietic cell transplantation. <i>British Journal of Haematology</i> , 2010, 148, 910-917.	1.2	21
27	The Addition of Sorafenib to Standard AML Treatment Results in a Substantial Reduction in Relapse Risk and Improved Survival. Updated Results from Long-Term Follow-up of the Randomized-Controlled Soraml Trial. <i>Blood</i> , 2017, 130, 721-721.	0.6	20
28	Osteomyelosclerosis, anemia and extramedullary hematopoiesis in mice lacking the transcription factor NFATc2. <i>Haematologica</i> , 2011, 96, 1580-1588.	1.7	19
29	Amphiregulin Expression Is a Predictive Biomarker for EGFR Inhibition in Metastatic Colorectal Cancer: Combined Analysis of Three Randomized Trials. <i>Clinical Cancer Research</i> , 2020, 26, 6559-6567.	3.2	17
30	Time from Diagnosis to Treatment Does Not Affect Outcome in Intensively Treated Patients with Newly Diagnosed Acute Myeloid Leukemia. <i>Blood</i> , 2019, 134, 13-13.	0.6	16
31	Relation of early tumor shrinkage (ETS) observed in first-line treatment to efficacy parameters of subsequent treatment in FIRE-3 (AIOKRK0306). <i>International Journal of Cancer</i> , 2017, 140, 1918-1925.	2.3	15
32	Regulation of Down Syndrome Critical Region 1 expression by Nuclear Factor of Activated T cells in megakaryocytes. <i>British Journal of Haematology</i> , 2009, 144, 395-408.	1.2	14
33	Third-party mesenchymal stem cells as part of the management of graft-failure after haploidentical stem cell transplantation. <i>Leukemia Research</i> , 2009, 33, e215-e217.	0.4	14
34	Regulation of fas/fas ligand-mediated apoptosis by nuclear factor of activated T cells in megakaryocytes. <i>British Journal of Haematology</i> , 2012, 156, 523-534.	1.2	14
35	Normal intrinsic Th1/Th2 balance in patients with chronic phase chronic myeloid leukemia not treated with interferon-alpha or imatinib. <i>Haematologica</i> , 2003, 88, 754-61.	1.7	13
36	Relevance of liver-limited disease in metastatic colorectal cancer: Subgroup findings of the FIRE-3/AIO KRK0306 trial. <i>International Journal of Cancer</i> , 2018, 142, 1047-1055.	2.3	12

#	ARTICLE	IF	CITATIONS
37	UBTF::ATXN7L3 gene fusion defines novel B cell precursor ALL subtype with CDX2 expression and need for intensified treatment. <i>Leukemia</i> , 2022, 36, 1676-1680.	3.3	12
38	Early weight loss is an independent risk factor for shorter survival and increased side effects in patients with metastatic colorectal cancer undergoing first-line treatment within the randomized Phase III trial FIRE-3 (AIO KKRK0306). <i>International Journal of Cancer</i> , 2022, 150, 112-123.	2.3	10
39	Interference of Ha-ras with inositol trisphosphate-mediated Ca ²⁺ -release. <i>FEBS Letters</i> , 1991, 291, 113-116.	1.3	5
40	Efficacy of FOLFIRI plus cetuximab vs FOLFIRI plus bevacizumab in 1st-line treatment of older patients with RAS wild-type metastatic colorectal cancer: an analysis of the randomised trial FIRE-3. <i>British Journal of Cancer</i> , 2022, 127, 836-843.	2.9	5
41	Azacitidine Followed By Intensive Induction/Consolidation Chemotherapy in Older Patients with Acute Myeloid Leukemia (AML): Results from the Randomized AML-AZA Trial of the Study Alliance Leukemias (SAL). <i>Blood</i> , 2014, 124, 946-946.	0.6	4
42	Preinfection laboratory parameters may predict COVID-19 severity in tumor patients. <i>Cancer Medicine</i> , 2021, 10, 4424-4436.	1.3	3
43	Mutational profiles of metastatic colorectal cancer treated with FOLFIRI plus cetuximab or bevacizumab before and after secondary resection (AIO KKRK 0306; FIRE-3). <i>International Journal of Cancer</i> , 2021, 149, 1935-1943.	2.3	3
44	5-azacitidine Treatment of Imminent Relapse Defined by Decreasing Donor CD34+ Progenitor Subset Chimerism in Patients with CD34+ High-Risk Myelodysplastic Syndromes (MDS) or Acute Myeloid Leukemia (AML) after Allogeneic Stem Cell Transplantation. <i>Blood</i> , 2008, 112, 2143-2143.	0.6	3
45	Step-in Dosing in the Bosutinib Dose Optimization Study (BODO) Failed to Reduce Gastrointestinal (GI) Toxicity in Patients Failing Second Generation TKI (2G-TKI) in Chronic Phase Chronic Myeloid Leukemia (CML) but Suggests Promising Molecular Response. <i>Blood</i> , 2021, 138, 3608-3608.	0.6	3
46	Reduced-Intensity Conditioning (RIC) with Busulfan, Fludarabine and Campath-1H Is Complicated by a High Rate of Graft Failure and Severe Viral Complications in Patients with CLL. <i>Blood</i> , 2004, 104, 5080-5080.	0.6	2
47	Flow Cytometric Detection of Minimal Residual Disease One Year Post Allogeneic Stem Cell Transplantation Predicts Outcome in Patients with B-CLL. <i>Blood</i> , 2009, 114, 202-202.	0.6	2
48	Midazolam: Significant Pain Reduction in Patients Undergoing Bone Marrow Puncture - a Clinical Trial. <i>Blood</i> , 2004, 104, 90-90.	0.6	2
49	Prior Treatment with Alemtuzumab Interferes with T-Cell Engraftment After Allogeneic Stem Cell Transplantation in Patients with Chronic Lymphocytic Leukemia. <i>Blood</i> , 2009, 114, 3351-3351.	0.6	2
50	Treatment-Free Remission (TFR) after Two Different Durations of Nilotinib Consolidation in Patients with Chronic Myeloid Leukemia (CML) Previously Treated with Imatinib: Enestpath Study Results. <i>Blood</i> , 2021, 138, 635-635.	0.6	2
51	Graft Versus Host Disease Prophylaxis with Everolimus and Tacrolimus in Patients with Myelodysplastic Syndromes (MDS) and Acute Myeloid Leukaemia (AML) Receiving Allogeneic Peripheral Blood Stem Cell Transplantation (PBSCT). <i>Blood</i> , 2006, 108, 2886-2886.	0.6	1
52	Monitoring of Donor Chimerism in CD34+ Peripheral Blood Progenitors Allows to Detect Minimal Residual Disease after Allogeneic Stem Cell Transplantation -Results of a Randomized Trial. <i>Blood</i> , 2008, 112, 340-340.	0.6	1
53	Graft Versus Host Disease (GVHD) Prophylaxis with Everolimus and Tacrolimus Is Associated with a High Incidence of Sinusoidal Obstruction Syndrome and Microangiopathy - Results of the EVTAC Trial. <i>Blood</i> , 2008, 112, 1172-1172.	0.6	1
54	Gemtuzumab Ozogamicin (Mylotarg®) as Part of Reduced-Intensity Conditioning for Allogeneic Hematopoietic Cell Transplantation in Patients with Relapsed Acute Myeloid Leukemia. <i>Blood</i> , 2004, 104, 1245-1245.	0.6	0

#	ARTICLE	IF	CITATIONS
55	The Campath-Level at the Day of Transplant Predicts T-Cell Engraftment after Allogeneic Stem Cell Transplantation.. Blood, 2006, 108, 2951-2951.	0.6	0
56	Nuclear Factor of Activated T Cells Regulates Down Syndrome Critical Region 1 Gene Expression in Megakaryocytes.. Blood, 2006, 108, 1193-1193.	0.6	0
57	Specific Regulation of NFAT (Nuclear Factors of Activated T Cells) Expression in CD34+ Cells Differentiating into Diverse Hematopoietic Lineages.. Blood, 2006, 108, 4213-4213.	0.6	0
58	The Calcineurin/NFAT Signaling Pathway Is a Target for the Collagen Type I-Induced Calcium Response in Megakaryocytes and Mediates Expression of Megakaryocytic Genes.. Blood, 2007, 110, 3647-3647.	0.6	0
59	Reciprocal Regulation of DSCR1 (Down Syndrome Critical Region 1) Expression and NFAT (Nuclear Tj ETQq1 1 0.784314 rgBJ /Overlo	0.6	0
60	Osteomyelosclerosis, Anemia and Extramedullary Hematopoiesis in Mice Deficient for the Transcription Factor NFAT (Nuclear Factor of Activated T Cells) c2. Blood, 2008, 112, 3726-3726.	0.6	0
61	Regulation of Fas/Fas Ligand-Mediated Apoptosis in Megakaryocytes by Nuclear Factor of Activated T Cells.. Blood, 2009, 114, 4023-4023.	0.6	0
62	Lack of the Transcription Factor NFAT (Nuclear Factor of Activated T cells) c2 in Hematopoietic Progenitor Cells Results in Profound Hematological Abnormalities in Mice. Blood, 2011, 118, 1296-1296.	0.6	0
63	Improved Safety with the Use of Subcutaneous Bortezomib in Combination with Panobinostat and Dexamethasone: Preliminary Data from a Panobinostat Global Expanded Treatment Protocol. Blood, 2016, 128, 5692-5692.	0.6	0
64	Enestpath Leukemic Stem Cell (LSC) Sub-Study: Analyzing Characteristics of LSC-Positive Patients and Impact of Switch from Imatinib to Nilotinib Therapy on LSCs in Patients with Chronic Myeloid Leukemia. Blood, 2019, 134, 4160-4160.	0.6	0