

Abdolali Moshfe

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3590069/publications.pdf>

Version: 2024-02-01

18
papers

257
citations

1040056

9
h-index

940533

16
g-index

18
all docs

18
docs citations

18
times ranked

396
citing authors

| # | ARTICLE | IF | CITATIONS |
|----|---|-----|-----------|
| 1 | Authors'™ response. Comparative Immunology, Microbiology and Infectious Diseases, 2021, 76, 101645. | 1.6 | 0 |
| 2 | Neospora caninum Infection in Cattle in the Province of Kohgiluyeh and Boyer Ahmad, Southwest of Iran: Seroprevalence and Molecular Assessment. Journal of Parasitology Research, 2021, 2021, 1-6. | 1.2 | 1 |
| 3 | Molecular Genotyping of Toxoplasma gondii in Sheep Aborted Fetuses Reveals Predominance of Type I Infection in Southwest of Iran. Iranian Journal of Parasitology, 2020, 15, 374-382. | 0.6 | 5 |
| 4 | Clinical Features, Diagnosis and Management of Patients with Suspicion of Fascioliasis in Kohgiluyeh and Boyer-Ahmad Province, Southwestern Iran. Iranian Journal of Parasitology, 2020, 15, 84-90. | 0.6 | 3 |
| 5 | Molecular genotyping and serological evaluation of Toxoplasma gondii in mothers and their spontaneous aborted fetuses in Southwest of Iran. Comparative Immunology, Microbiology and Infectious Diseases, 2019, 66, 101342. | 1.6 | 14 |
| 6 | Prevalence of bovine fascioliasis in a new-emerging focus of human fascioliasis in BoyerAhmad district, southwest of Iran. Comparative Immunology, Microbiology and Infectious Diseases, 2019, 66, 101350. | 1.6 | 3 |
| 7 | Seroepidemiological study of cystic echinococcosis in nomadic communities in the southwest of Iran: A population-based study. Journal of Immunoassay and Immunochemistry, 2019, 40, 183-192. | 1.1 | 15 |
| 8 | Role of environmental, climatic risk factors and livestock animals on the occurrence of cutaneous leishmaniasis in newly emerging focus in Iran. Journal of Infection and Public Health, 2018, 11, 425-433. | 4.1 | 33 |
| 9 | Toxoplasma gondii in Blood Donors: A Study in Boyer-Ahmad County, Southwest Iran. Interdisciplinary Perspectives on Infectious Diseases, 2018, 2018, 1-5. | 1.4 | 14 |
| 10 | Activity of Seeds Against a Clinical Strain of Genotype T4. Iranian Journal of Pharmaceutical Research, 2018, 17, 661-667. | 0.5 | 5 |
| 11 | Production of Monoclonal Antibody Against Excretory-Secretory Antigen of Fasciola hepatica and Evaluation of Its Efficacy in the Diagnosis of Fascioliasis. Monoclonal Antibodies in Immunodiagnosis and Immunotherapy, 2017, 36, 8-14. | 1.6 | 11 |
| 12 | Helminth Infections of Rodents and Their Zoonotic Importance in Boyer-Ahmad District, Southwestern Iran. Iranian Journal of Parasitology, 2017, 12, 572-579. | 0.6 | 9 |
| 13 | Prevalence and risk factors of intestinal protozoan infections: a population-based study in rural areas of Boyer-Ahmad district, Southwestern Iran. BMC Infectious Diseases, 2016, 16, 703. | 2.9 | 48 |
| 14 | Molecular Evaluation of a Case of Visceral Leishmaniasis Due to Leishmania tropica in Southwestern Iran. Iranian Journal of Parasitology, 2016, 11, 126-30. | 0.6 | 13 |
| 15 | Clinical and Molecular Evaluation of a Case of Giant Primary Splenic Hydatid Cyst: A Case Report. Iranian Journal of Parasitology, 2016, 11, 585-590. | 0.6 | 8 |
| 16 | Epidemiology of Human Fascioliasis and Intestinal Helminthes in Rural Areas of Boyer-Ahmad Township, Southwest Iran; A Population Based Study. Iranian Journal of Public Health, 2015, 44, 1520-5. | 0.5 | 21 |
| 17 | Molecular and Morphological Characterization of Fasciola spp. Isolated from Different Host Species in a Newly Emerging Focus of Human Fascioliasis in Iran. Veterinary Medicine International, 2014, 2014, 1-10. | 1.5 | 42 |
| 18 | A Consistent PCR-RFLP Assay Based on ITS-2 Ribosomal DNA for Differentiation of Fasciola Species. Iranian Journal of Basic Medical Sciences, 2013, 16, 1266-9. | 1.0 | 12 |