

Stephen A Duncan

List of Publications by Year in descending order

Source: <https://exaly.com/author-pdf/3596422/publications.pdf>

Version: 2024-02-01

107
papers

11,664
citations

38660

50
h-index

33814

99
g-index

112
all docs

112
docs citations

112
times ranked

14243
citing authors

#	ARTICLE	IF	CITATIONS
1	Advancements in Disease Modeling and Drug Discovery Using iPSC-Derived Hepatocyte-like Cells. <i>Genes</i> , 2022, 13, 573.	1.0	5
2	Small molecules targeting the NADH-binding pocket of VDAC modulate mitochondrial metabolism in hepatocarcinoma cells. <i>Biomedicine and Pharmacotherapy</i> , 2022, 150, 112928.	2.5	6
3	Chromatin remodeling is restricted by transient GATA6 binding during iPSC differentiation to definitive endoderm. <i>IScience</i> , 2022, 25, 104300.	1.9	3
4	GATA6 defines endoderm fate by controlling chromatin accessibility during differentiation of human-induced pluripotent stem cells. <i>Cell Reports</i> , 2021, 35, 109145.	2.9	32
5	Adipose expression of CREB3L3 modulates body weight during obesity. <i>Scientific Reports</i> , 2021, 11, 19400.	1.6	2
6	Generation of isogenic Propionyl-CoA carboxylase beta subunit (PCCB) deficient induced pluripotent stem cell lines. <i>Stem Cell Research</i> , 2020, 48, 101953.	0.3	1
7	FoxA factors: the chromatin key and doorstop essential for liver development and function. <i>Genes and Development</i> , 2020, 34, 1003-1004.	2.7	7
8	Enhanced genome editing in human iPSCs with CRISPR-CAS9 by co-targeting <i>ATP1a1</i> . <i>PeerJ</i> , 2020, 8, e9060.	0.9	10
9	iPSC-Derived Hepatocytes as a Platform for Disease Modeling and Drug Discovery. <i>Frontiers in Medicine</i> , 2019, 6, 265.	1.2	90
10	HNF4A Regulates the Formation of Hepatic Progenitor Cells from Human iPSC-Derived Endoderm by Facilitating Efficient Recruitment of RNA Pol II. <i>Genes</i> , 2019, 10, 21.	1.0	33
11	The Use of Human Pluripotent Stem Cells for Modeling Liver Development and Disease. <i>Hepatology</i> , 2019, 69, 1306-1316.	3.6	18
12	A Screen Using iPSC-Derived Hepatocytes Reveals NAD ⁺ as a Potential Treatment for mtDNA Depletion Syndrome. <i>Cell Reports</i> , 2018, 25, 1469-1484.e5.	2.9	36
13	Using Human Induced Pluripotent Stem Cell-derived Hepatocyte-like Cells for Drug Discovery. <i>Journal of Visualized Experiments</i> , 2018, , .	0.2	8
14	Lack of MTTP Activity in Pluripotent Stem Cell-Derived Hepatocytes and Cardiomyocytes Abolishes apoB Secretion and Increases Cell Stress. <i>Cell Reports</i> , 2017, 19, 1456-1466.	2.9	36
15	A small molecule screen reveals that HSP90 ¹ promotes the conversion of iPSC-derived endoderm to a hepatic fate and regulates HNF4A turnover. <i>Development (Cambridge)</i> , 2017, 144, 1764-1774.	1.2	24
16	ATP-Binding Cassette Transporter A1 Deficiency in Human Induced Pluripotent Stem Cell-Derived Hepatocytes Abrogates HDL Biogenesis and Enhances Triglyceride Secretion. <i>EBioMedicine</i> , 2017, 18, 139-145.	2.7	23
17	A Drug Screen using Human iPSC-Derived Hepatocyte-like Cells Reveals Cardiac Glycosides as a Potential Treatment for Hypercholesterolemia. <i>Cell Stem Cell</i> , 2017, 20, 478-489.e5.	5.2	92
18	Large, Diverse Population Cohorts of hiPSCs and Derived Hepatocyte-like Cells Reveal Functional Genetic Variation at Blood Lipid-Associated Loci. <i>Cell Stem Cell</i> , 2017, 20, 558-570.e10.	5.2	138

#	ARTICLE	IF	CITATIONS
19	N-glycoprotein surfaceome of human induced pluripotent stem cell derived hepatic endoderm. <i>Proteomics</i> , 2017, 17, 1600397.	1.3	19
20	GATA6 is essential for endoderm formation from human pluripotent stem cells. <i>Biology Open</i> , 2017, 6, 1084-1095.	0.6	45
21	Modeling Inborn Errors of Hepatic Metabolism Using Induced Pluripotent Stem Cells. <i>Arteriosclerosis, Thrombosis, and Vascular Biology</i> , 2017, 37, 1994-1999.	1.1	17
22	Mapping the Cell-Surface N-Glycoproteome of Human Hepatocytes Reveals Markers for Selecting a Homogeneous Population of iPSC-Derived Hepatocytes. <i>Stem Cell Reports</i> , 2016, 7, 543-556.	2.3	44
23	Liver Capsule: Multipotent stem cells and their lineage restriction to hepatocytes. <i>Hepatology</i> , 2016, 64, 1330-1330.	3.6	1
24	Design of a Vitronectin-Based Recombinant Protein as a Defined Substrate for Differentiation of Human Pluripotent Stem Cells into Hepatocyte-Like Cells. <i>PLoS ONE</i> , 2015, 10, e0136350.	1.1	13
25	FGF2 mediates hepatic progenitor cell formation during human pluripotent stem cell differentiation by inducing the WNT antagonist NKD1. <i>Genes and Development</i> , 2015, 29, 2463-2474.	2.7	32
26	Sall4 overexpression blocks murine hematopoiesis in a dose-dependent manner. <i>Experimental Hematology</i> , 2015, 43, 53-64.e8.	0.2	20
27	Generation of iPSCs as a Pooled Culture Using Magnetic Activated Cell Sorting of Newly Reprogrammed Cells. <i>PLoS ONE</i> , 2015, 10, e0134995.	1.1	30
28	Restoration of Liver Function Following Transplantation of Healthy Hepatocytes into the Fah ^{-/-} IL2rg ^{-/-} Rat Model. <i>FASEB Journal</i> , 2015, 29, LB681.	0.2	0
29	Epicardial GATA factors regulate early coronary vascular plexus formation. <i>Developmental Biology</i> , 2014, 386, 204-215.	0.9	10
30	Aneuploidy is permissive for hepatocyte-like cell differentiation from human induced pluripotent stem cells. <i>BMC Research Notes</i> , 2014, 7, 437.	0.6	14
31	Engineering liver tissue from induced pluripotent stem cells: A first step in generating new organs for transplantation?. <i>Hepatology</i> , 2013, 58, 2198-2201.	3.6	8
32	Identification of small molecules for human hepatocyte expansion and iPSC differentiation. <i>Nature Chemical Biology</i> , 2013, 9, 514-520.	3.9	230
33	Differentiation of Hepatocytes from Pluripotent Stem Cells. <i>Current Protocols in Stem Cell Biology</i> , 2013, 26, 1G.4.1-1G.4.13.	3.0	96
34	Generation of Hepatocyte-Like Cells from Human Pluripotent Stem Cells. , 2013, , 139-147.		4
35	A Cell Surfaceome Map for Immunophenotyping and Sorting Pluripotent Stem Cells. <i>Molecular and Cellular Proteomics</i> , 2012, 11, 303-316.	2.5	58
36	Induction of Cardiomyogenesis in Human Embryonic Stem Cells by Human Embryonic Stem Cell-Derived Definitive Endoderm. <i>Stem Cells and Development</i> , 2012, 21, 987-994.	1.1	3

#	ARTICLE	IF	CITATIONS
37	Modeling hepatitis C virus infection using human induced pluripotent stem cells. Proceedings of the National Academy of Sciences of the United States of America, 2012, 109, 2544-2548.	3.3	197
38	Comparison of Cardiomyogenic Potential among Human ESC and iPSC Lines. Cell Transplantation, 2012, 21, 2523-2530.	1.2	29
39	JD induced pluripotent stem cell-derived hepatocytes faithfully recapitulate the pathophysiology of familial hypercholesterolemia. Hepatology, 2012, 56, 2163-2171.	3.6	120
40	Hepatocyte-like cells differentiated from human induced pluripotent stem cells: Relevance to cellular therapies. Stem Cell Research, 2012, 9, 196-207.	0.3	95
41	Endoplasmic reticulum-tethered transcription factor cAMP responsive element-binding protein, hepatocyte specific, regulates hepatic lipogenesis, fatty acid oxidation, and lipolysis upon metabolic stress in mice. Hepatology, 2012, 55, 1070-1082.	3.6	163
42	Pancreas-specific deletion of mouse Gata4 and Gata6 causes pancreatic agenesis. Journal of Clinical Investigation, 2012, 122, 3516-3528.	3.9	138
43	GATA Factors Regulate Proliferation, Differentiation, and Gene Expression in Small Intestine of Mature Mice. Gastroenterology, 2011, 140, 1219-1229.e2.	0.6	91
44	HNF4A is essential for specification of hepatic progenitors from human pluripotent stem cells. Development (Cambridge), 2011, 138, 4143-4153.	1.2	178
45	The transcription factor cyclic AMP-responsive element-binding protein H regulates triglyceride metabolism. Nature Medicine, 2011, 17, 812-815.	15.2	174
46	Loss of intestinal GATA4 prevents diet-induced obesity and promotes insulin sensitivity in mice. American Journal of Physiology - Endocrinology and Metabolism, 2011, 300, E478-E488.	1.8	17
47	HNF4A is essential for specification of hepatic progenitors from human pluripotent stem cells. Journal of Cell Science, 2011, 124, e1-e1.	1.2	1
48	Laboratory-Scale Purification of a Recombinant E-Cadherin-IgG Fc Fusion Protein That Provides a Cell Surface Matrix for Extended Culture and Efficient Subculture of Human Pluripotent Stem Cells. Springer Protocols, 2011, , 25-35.	0.1	0
49	Isoflurane Preconditioning Elicits Competent Endogenous Mechanisms of Protection from Oxidative Stress in Cardiomyocytes Derived from Human Embryonic Stem Cells. Anesthesiology, 2010, 113, 906-916.	1.3	41
50	Highly efficient generation of human hepatocyte-like cells from induced pluripotent stem cells. Hepatology, 2010, 51, 297-305.	3.6	1,081
51	Culture of human pluripotent stem cells using completely defined conditions on a recombinant E-cadherin substratum. BMC Developmental Biology, 2010, 10, 60.	2.1	169
52	Generation of human induced pluripotent stem cells by simple transient transfection of plasmid DNA encoding reprogramming factors. BMC Developmental Biology, 2010, 10, 81.	2.1	191
53	The Transcription Factor GATA-6 Regulates Pathological Cardiac Hypertrophy. Circulation Research, 2010, 107, 1032-1040.	2.0	88
54	Foxa1 Functions as a Pioneer Transcription Factor at Transposable Elements to Activate Afp during Differentiation of Embryonic Stem Cells. Journal of Biological Chemistry, 2010, 285, 16135-16144.	1.6	65

#	ARTICLE	IF	CITATIONS
55	Transcriptional Control of Hepatocyte Differentiation. Progress in Molecular Biology and Translational Science, 2010, 97, 79-101.	0.9	13
56	Organogenesis and Development of the Liver. Developmental Cell, 2010, 18, 175-189.	3.1	649
57	hESC-Derived Definitive Endoderm Induces Cardiomyogenesis in Human Embryonic Stem Cells.. FASEB Journal, 2010, 24, 175.2.	0.2	0
58	ER Stress Controls Iron Metabolism Through Induction of Heparin. Science, 2009, 325, 877-880.	6.0	278
59	Hepatocyte expression of serum response factor is essential for liver function, hepatocyte proliferation and survival, and postnatal body growth in mice. Hepatology, 2009, 49, 1645-1654.	3.6	27
60	Hepatocyte nuclear factor 4 β is implicated in endoplasmic reticulum stress-induced acute phase response by regulating expression of cyclic adenosine monophosphate responsive element binding protein H. Hepatology, 2008, 48, 1242-1250.	3.6	88
61	GATA4 Is Essential for Jejunal Function in Mice. Gastroenterology, 2008, 135, 1676-1686.e1.	0.6	80
62	Loss of both GATA4 and GATA6 blocks cardiac myocyte differentiation and results in acardia in mice. Developmental Biology, 2008, 317, 614-619.	0.9	193
63	Design of the Artificial Acellular Feeder Layer for the Efficient Propagation of Mouse Embryonic Stem Cells. Journal of Biological Chemistry, 2008, 283, 26468-26476.	1.6	37
64	Cardiomyocyte GATA4 functions as a stress-responsive regulator of angiogenesis in the murine heart. Journal of Clinical Investigation, 2008, 118, 387-387.	3.9	0
65	Junctional Adhesion Molecule-A Is Critical for the Formation of Pseudocanaliculi and Modulates E-cadherin Expression in Hepatic Cells. Journal of Biological Chemistry, 2007, 282, 28137-28148.	1.6	42
66	Light Chain 1 of Microtubule-associated Protein 1B Can Negatively Regulate the Action of Pes1. Journal of Biological Chemistry, 2007, 282, 11308-11316.	1.6	7
67	Development of the mammalian liver and ventral pancreas is dependent on GATA4. BMC Developmental Biology, 2007, 7, 37.	2.1	165
68	Cited2, a coactivator of HNF4 β , is essential for liver development. EMBO Journal, 2007, 26, 4445-4456.	3.5	70
69	Cardiomyocyte GATA4 functions as a stress-responsive regulator of angiogenesis in the murine heart. Journal of Clinical Investigation, 2007, 117, 3198-3210.	3.9	212
70	Transcriptional regulation of the human hepatic lipase (LIPC) gene promoter. Journal of Lipid Research, 2006, 47, 1463-1477.	2.0	38
71	Hepatocyte Nuclear Factor 4 β Is Essential for Embryonic Development of the Mouse Colon. Gastroenterology, 2006, 130, 19.e1-19.e.	0.6	143
72	Essential function of PTP-PEST during mouse embryonic vascularization, mesenchyme formation, neurogenesis and early liver development. Mechanisms of Development, 2006, 123, 869-880.	1.7	54

#	ARTICLE	IF	CITATIONS
73	Generation of mice harbouring a conditional loss-of-function allele of Gata6. <i>BMC Developmental Biology</i> , 2006, 6, 19.	2.1	67
74	Improved cardiac function in infarcted mice after treatment with pluripotent embryonic stem cells. <i>The Anatomical Record Part A: Discoveries in Molecular, Cellular, and Evolutionary Biology</i> , 2006, 288A, 1216-1224.	2.0	57
75	A threshold of GATA4 and GATA6 expression is required for cardiovascular development. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2006, 103, 11189-11194.	3.3	170
76	Cardiac-Specific Deletion of Gata4 Reveals Its Requirement for Hypertrophy, Compensation, and Myocyte Viability. <i>Circulation Research</i> , 2006, 98, 837-845.	2.0	384
77	Hepatocyte nuclear factor 4 $\hat{\pm}$ orchestrates expression of cell adhesion proteins during the epithelial transformation of the developing liver. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2006, 103, 8419-8424.	3.3	225
78	Robert H. Costa. <i>Journal of Clinical Investigation</i> , 2006, 116, 3086-3086.	3.9	0
79	Generation of embryos directly from embryonic stem cells by tetraploid embryo complementation reveals a role for GATA factors in organogenesis. <i>Biochemical Society Transactions</i> , 2005, 33, 1534.	1.6	13
80	Embryonic development of the liver. <i>Hepatology</i> , 2005, 41, 956-967.	3.6	274
81	GATA6 Is Essential for Embryonic Development of the Liver but Dispensable for Early Heart Formation. <i>Molecular and Cellular Biology</i> , 2005, 25, 2622-2631.	1.1	216
82	The MODY1 gene HNF-4 $\hat{\pm}$ regulates selected genes involved in insulin secretion. <i>Journal of Clinical Investigation</i> , 2005, 115, 1006-1015.	3.9	195
83	GATA4 is essential for formation of the proepicardium and regulates cardiogenesis. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 2004, 101, 12573-12578.	3.3	316
84	Conserved Enhancer in the Serum Response Factor Promoter Controls Expression During Early Coronary Vasculogenesis. <i>Circulation Research</i> , 2004, 94, 1059-1066.	2.0	21
85	Progression of HCC in mice is associated with a downregulation in the expression of hepatocyte nuclear factors. <i>Hepatology</i> , 2004, 39, 1038-1047.	3.6	192
86	HNF4: A central regulator of hepatocyte differentiation and function. <i>Hepatology</i> , 2003, 37, 1249-1253.	3.6	245
87	Hepatocyte nuclear factor 4 $\hat{\pm}$ controls the development of a hepatic epithelium and liver morphogenesis. <i>Nature Genetics</i> , 2003, 34, 292-296.	9.4	530
88	The orphan nuclear receptor HNF4 $\hat{\pm}$ determines PXR- and CAR-mediated xenobiotic induction of CYP3A4. <i>Nature Medicine</i> , 2003, 9, 220-224.	15.2	418
89	Mechanisms controlling early development of the liver. <i>Mechanisms of Development</i> , 2003, 120, 19-33.	1.7	160
90	Pescadillo Is Essential for Nucleolar Assembly, Ribosome Biogenesis, and Mammalian Cell Proliferation. <i>Journal of Biological Chemistry</i> , 2002, 277, 45347-45355.	1.6	106

#	ARTICLE	IF	CITATIONS
91	Generation of a conditionally null allele of <i>hnf4b</i> . <i>Genesis</i> , 2002, 32, 130-133.	0.8	41
92	Gene Targeting in the Mouse: Advances in Introduction of Transgenes into the Genome by Homologous Recombination. <i>Endocrine</i> , 2002, 19, 229-238.	2.2	42
93	Generation of single-copy transgenic mouse embryos directly from ES cells by tetraploid embryo complementation. <i>BMC Biotechnology</i> , 2001, 1, 12.	1.7	27
94	Dynamic expression of a glutamate decarboxylase gene in multiple non-neural tissues during mouse development. , 2001, 1, 1.		45
95	An efficient method to successively introduce transgenes into a given genomic locus in the mouse. <i>BMC Developmental Biology</i> , 2001, 1, 10.	2.1	4
96	The pancreas and its heartless beginnings. <i>Nature Genetics</i> , 2001, 27, 355-356.	9.4	1
97	BMPs on the road to hepatogenesis. <i>Genes and Development</i> , 2001, 15, 1879-1884.	2.7	31
98	The Murine <i>Pes1</i> Gene Encodes a Nuclear Protein Containing a BRCT Domain. <i>Genomics</i> , 2000, 70, 201-210.	1.3	27
99	Transcriptional regulation of liver development. , 2000, 219, 131.		3
100	Transcriptional regulation of liver development. <i>Developmental Dynamics</i> , 2000, 219, 131-142.	0.8	93
101	Mammalian hepatocyte differentiation requires the transcription factor HNF4 β . <i>Genes and Development</i> , 2000, 14, 464-474.	2.7	398
102	In Situ Hybridization with ³³ P-Labeled RNA Probes for Determination of Cellular Expression Patterns of Liver Transcription Factors in Mouse Embryos. <i>Methods</i> , 1998, 16, 29-41.	1.9	25
103	The maturity-onset diabetes of the young (MODY1) transcription factor HNF4 β regulates expression of genes required for glucose transport and metabolism. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1997, 94, 13209-13214.	3.3	374
104	Normal cerebellar development but susceptibility to seizures in mice lacking G protein-coupled, inwardly rectifying K ⁺ channel GIRK2. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1997, 94, 923-927.	3.3	312
105	STAT signaling is active during early mammalian development. , 1997, 208, 190-198.		83
106	Disruption of the <i>MacMARCKS</i> gene prevents cranial neural tube closure and results in anencephaly. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1996, 93, 6275-6279.	3.3	90
107	Expression of transcription factor HNF-4 in the extraembryonic endoderm, gut, and nephrogenic tissue of the developing mouse embryo: HNF-4 is a marker for primary endoderm in the implanting blastocyst. <i>Proceedings of the National Academy of Sciences of the United States of America</i> , 1994, 91, 7598-7602.	3.3	333