

# Marta Staniszewska

## List of Publications by Citations

**Source:** <https://exaly.com/author-pdf/3600694/marta-staniszewska-publications-by-citations.pdf>

**Version:** 2024-04-28

This document has been generated based on the publications and citations recorded by exaly.com. For the latest version of this publication list, visit the link given above.

The third column is the impact factor (IF) of the journal, and the fourth column is the number of citations of the article.

31  
papers

513  
citations

14  
h-index

22  
g-index

31  
ext. papers

609  
ext. citations

5.7  
avg, IF

3.89  
L-index

| #  | Paper   | IF  | Citations |
|----|---|-----|-----------|
| 31 | Bisphenol A, 4-tert-octylphenol, and 4-nonylphenol in the Gulf of Gdańsk (Southern Baltic). <i>Archives of Environmental Contamination and Toxicology</i> , <b>2014</b> , 67, 335-47  | 3.2 | 104       |
| 30 | Occurrence and distribution of bisphenol A and alkylphenols in the water of the Gulf of Gdansk (Southern Baltic). <i>Marine Pollution Bulletin</i> , <b>2015</b> , 91, 372-9  | 6.7 | 51        |
| 29 | Mercury and Chlorinated Pesticides on the Highest Level of the Food Web as Exemplified by Herring from the Southern Baltic and African Penguins from the Zoo. <i>Water, Air, and Soil Pollution</i> , <b>2013</b> , 224, 1549   | 2.6 | 32        |
| 28 | Alkylphenols in Surface Sediments of the Gulf of Gdansk (Baltic Sea). <i>Water, Air, and Soil Pollution</i> , <b>2014</b> , 225, 2040   | 2.6 | 28        |
| 27 | The relationship between the black carbon and bisphenol A in sea and river sediments (Southern Baltic). <i>Journal of Environmental Sciences</i> , <b>2016</b> , 41, 24-32  | 6.4 | 25        |
| 26 | The fate of bisphenol A, 4-tert-octylphenol and 4-nonylphenol leached from plastic debris into marine water--experimental studies on biodegradation and sorption on suspended particulate matter and nano-TiO <sub>2</sub> . <i>Chemosphere</i> , <b>2016</b> , 145, 535-42 | 8.4 | 25        |
| 25 | The role of phytoplankton composition, biomass and cell volume in accumulation and transfer of endocrine disrupting compounds in the Southern Baltic Sea (The Gulf of Gdansk). <i>Environmental Pollution</i> , <b>2015</b> , 207, 319-28                                   | 9.3 | 23        |
| 24 | Changes of concentrations and possibility of accumulation of bisphenol A and alkylphenols, depending on biomass and composition, in zooplankton of the Southern Baltic (Gulf of Gdansk). <i>Environmental Pollution</i> , <b>2016</b> , 213, 489-501                        | 9.3 | 23        |
| 23 | Human Hair, Baltic Grey Seal ( <i>Halichoerus grypus</i> ) Fur and Herring Gull ( <i>Larus argentatus</i> ) Feathers as Accumulators of Bisphenol A and Alkylphenols. <i>Archives of Environmental Contamination and Toxicology</i> , <b>2017</b> , 72, 552-561             | 3.2 | 22        |
| 22 | The Microtox <sup>®</sup> biological test: Application in toxicity evaluation of surface waters and sediments in Poland. <i>Oceanological and Hydrobiological Studies</i> , <b>2007</b> , 36, 151-163   | 0.8 | 18        |
| 21 | Factors determining accumulation of bisphenol A and alkylphenols at a low trophic level as exemplified by mussels <i>Mytilus trossulus</i> . <i>Environmental Pollution</i> , <b>2017</b> , 220, 1147-1159  | 9.3 | 17        |
| 20 | Factors controlling benzo(a)pyrene concentration in aerosols in the urbanized coastal zone. A case study: Gdynia, Poland (Southern Baltic Sea). <i>Environmental Science and Pollution Research</i> , <b>2013</b> , 20, 4154-63   | 5.1 | 17        |
| 19 | Reconstruction of the pollution history of alkylphenols (4-tert-octylphenol, 4-nonylphenol) in the Baltic Sea. <i>Environmental Science and Pollution Research</i> , <b>2016</b> , 23, 11598-610  | 5.1 | 16        |
| 18 | Analytical methods and problems related to the determination of organotin compounds in marine sediments. <i>International Journal of Environmental Analytical Chemistry</i> , <b>2008</b> , 88, 747-774   | 1.8 | 15        |
| 17 | Air quality at two stations (Gdynia and Rumia) located in the region of Gulf of Gdansk during periods of intensive smog in Poland. <i>Air Quality, Atmosphere and Health</i> , <b>2019</b> , 12, 879-890  | 5.6 | 13        |
| 16 | Benzo(a)pyrene parallel measurements in PM <sub>10</sub> and PM <sub>2.5</sub> in the coastal zone of the Gulf of Gdansk (Baltic Sea) in the heating and non-heating seasons. <i>Environmental Science and Pollution Research</i> , <b>2018</b> , 25, 19458-19469           | 5.1 | 12        |
| 15 | Transfer of mercury and phenol derivatives across the placenta of Baltic grey seals ( <i>Halichoerus grypus grypus</i> ). <i>Environmental Pollution</i> , <b>2017</b> , 231, 1005-1012   | 9.3 | 11        |

|    |  |      |    |
|----|--|------|----|
| 14 | Maternal transfer of phenol derivatives in the Baltic grey seal <i>Halichoerus grypus grypus</i> . <i>Environmental Pollution</i> , <b>2018</b> , 242, 1642-1651   | 9.3  | 11 |
| 13 | Residue of chlorinated pesticides in fish caught in the Southern Baltic. <i>Oceanological and Hydrobiological Studies</i> , <b>2013</b> , 42, 251-259  | 0.8  | 9  |
| 12 | Sources, deposition flux and carcinogenic potential of PM2.5-bound polycyclic aromatic hydrocarbons in the coastal zone of the Baltic Sea (Gdynia, Poland). <i>Air Quality, Atmosphere and Health</i> , <b>2019</b> , 12, 1291-1301                              | 5.6  | 8  |
| 11 | Inhalation - Route of EDC exposure in seabirds ( <i>Larus argentatus</i> ) from the Southern Baltic. <i>Marine Pollution Bulletin</i> , <b>2017</b> , 117, 111-117   | 6.7  | 7  |
| 10 | Nonylphenol and 4-tert-octylphenol in the Gulf of Gdansk coastal zone. <i>Oceanological and Hydrobiological Studies</i> , <b>2011</b> , 40, 49-56  | 0.8  | 7  |
| 9  | Gastrointestinal and respiratory exposure of water birds to endocrine disrupting phenolic compounds. <i>Science of the Total Environment</i> , <b>2021</b> , 754, 142435   | 10.2 | 6  |
| 8  | Could biotransport be an important pathway in the transfer of phenol derivatives into the coastal zone and aquatic system of the Southern Baltic?. <i>Environmental Pollution</i> , <b>2020</b> , 262, 114358  | 9.3  | 3  |
| 7  | Dangerous compounds in the dredged material from the sea - Assessment of the current approach to the evaluation of contaminations based on the data from the Polish coastal zone (the Baltic Sea). <i>Marine Pollution Bulletin</i> , <b>2018</b> , 130, 324-334 | 6.7  | 3  |
| 6  | Concentrations of bisphenol a (BPA) in fresh pork loin meat under standard stock-farming conditions and after oral exposure - A preliminary study.. <i>Chemosphere</i> , <b>2022</b> , 295, 133816   | 8.4  | 2  |
| 5  | Analytical methods for determination of bisphenol A, 4--octylphenol and 4-nonylphenol in herrings and physiological fluids of the grey seal. <i>MethodsX</i> , <b>2018</b> , 5, 1124-1128  | 1.9  | 2  |
| 4  | The Influence of Transport on PAHs and Other Carbonaceous Species[(OC, EC) Concentration in Aerosols in the Coastal Zone of the Gulf of Gdansk (Gdynia). <i>Atmosphere</i> , <b>2021</b> , 12, 1005  | 2.7  | 2  |
| 3  | Distribution paths of endocrine disrupting phenolic compounds in waterbirds ( <i>Mergus merganser</i> , <i>Alca torda</i> , <i>Clangula hyemalis</i> ) from the Southern Baltic. <i>Science of the Total Environment</i> , <b>2021</b> , 793, 148556             | 10.2 | 1  |
| 2  | Bioaccumulation of phenolic endocrine disruptors in the clam <i>Rangia cuneata</i> : Storage in shells and influence of size and sex. <i>Environmental Research</i> , <b>2021</b> , 197, 111181  | 7.9  | 0  |
| 1  | Cs and K in gray seals <i>Halichoerus grypus</i> in the southern Baltic Sea. <i>Environmental Science and Pollution Research</i> , <b>2019</b> , 26, 17418-17426   | 5.1  |    |